

# **Appendix G**

## **Non-fishing Impacts to Essential Fish Habitat and Recommended Conservation Measures**

**Prepared by**

**National Marine Fisheries Service**

**January 2004**

This appendix to the Alaska Essential Fish Habitat Environmental Impact Statement was adapted from a document developed jointly by the National Marine Fisheries Service Alaska Region, Northwest Region, and Southwest Region, with contributions from the following people (listed in alphabetical order): Lt. Mark Boland, Eric Chavez, Bryant Chesney, Brian Cluer, Tracy Collier, Natalie Consentino-Manning, Joe Dillion, Bob Donnelly, Jeanne Hanson, Mark Helvey, Ron A. Heintz, Bob Hoffman, Thom Hooper, DeAnee Kirkpatrick, Brian Lance, Stacey Li, Marc Liverman, Matt Longenbaugh, Jon Mann, Leah Mahan, Ben Meyer, Adam Moles, Nancy Munn, Ken Phippen, Maggie Sommer, John Stadler, Dan Tonnes, and Susan Walker.

# TABLE OF CONTENTS

<b>1.0 INTRODUCTION</b>	G-1
<b>2.0 UPLAND ACTIVITIES</b>	G-8
2.1 Nonpoint Source Pollution	G-8
2.1.1 Agricultural/Nursery Runoff	G-8
2.1.2 Silviculture/Timber Harvest	G-11
2.1.3 Pesticide Application	G-13
2.2 Urban/Suburban Development	G-14
2.3 Road Building and Maintenance	G-16
<b>3.0 RIVERINE ACTIVITIES</b>	G-18
3.1 Mining	G-18
3.1.1 Mineral Mining	G-19
3.1.2 Sand and Gravel Mining	G-20
3.2 Debris Removal	G-22
3.2.1 Organic Debris	G-22
3.2.2 Inorganic Debris	G-23
3.3 Dam Operation	G-24
3.4 Commercial and Domestic Water Use	G-25
<b>4.0 ESTUARINE ACTIVITIES</b>	G-26
4.1 Dredging	G-26
4.2 Disposal/Landfills	G-28
4.2.1 Disposal of Dredged Material	G-28
4.2.2 Fill Material	G-29
4.3 Vessel Operations/Transportation/Navigation	G-29
4.4 Introduction of Exotic Species	G-31
4.5 Pile Installation and Removal	G-33
4.5.1 Pile Driving	G-33
4.5.2 Pile Removal	G-35
4.6 Overwater Structures	G-36
4.7 Flood Control/Shoreline Protection	G-39
4.8 Water Control Structures	G-40
4.9 Log Transfer Facilities/In-water Log Storage	G-42
4.10 Utility Line/Cables/Pipeline Installation	G-43
4.11 Commercial Utilization of Habitat	G-45
<b>5.0 COASTAL/MARINE ACTIVITIES</b>	G-47
5.1 Point Source Discharge	G-47
5.2 Fish Processing Waste—Shoreside and Vessel Operation	G-49
5.3 Water Intake Structures/Discharge Plumes	G-51
5.4 Oil/Gas Exploration/Development/Production	G-53
5.5 Habitat Restoration/Enhancement	G-56
5.6 Marine Mining	G-58
5.7 Persistent Organic Pollutants (North Pacific and Alaska Specific)	G-59
<b>REFERENCES</b>	G-63

## Tables

<b>Table 1.</b> Recommendations from the 1999 EFH EA Regarding Non-fishing Threats . . . . .	G-5
<b>Table 2.</b> Summary of Non-fishing Threats to EFH from the 1999 EFH EA . . . . .	G-6
<b>Table 3.</b> The Twelve Persistent Organic Pollutants Regulated by the POPs Treaty . . . . .	G-60

## ACRONYMS AND ABBREVIATIONS

AAPA	American Association of Port Authorities
ACZA	ammoniacal copper zinc arsenate
AFS	American Fisheries Society
Ag	silver
AI	Aleutian Islands
AMAP	Arctic Mapping and Assessment Program
As	arsenic
ATTF	Alaska Timber Task Force
BMPs	best management practices
BOD	biochemical oxygen demand
BTA	best technology available
CCA	chromated copper arsenate
Cd	cadmium
CERCLA	Comprehensive Environmental Response, Compensation, and Liability Act
Cr	chromium
Council	North Pacific Fishery Management Council
CSREEs	Cooperative State Research, Education, and Extension
CWA	Clean Water Act
dB	decibel
DDT	dichloro-diphenyl-trichloroethane
DDE	dichlorodiphenyl dichloroethylene
DoN	Department of the Navy
EBS	Bering Sea
EFH	Essential Fish Habitat
EPA	U.S. Environmental Protection Agency
ESA	Endangered Species Act
FC	fecal coliform (bacteria)
FERC	Federal Energy Regulatory Commission
FIFRA	Federal Insecticide, Fungicide, and Rodenticide Act
FL	fork length
FMCs	Fishery Management Councils
FREP	Fertilizer Research and Education Program
GIS	geographical information system
GOA	Gulf of Alaska
HCH	hexachloro cyclohexanes
Hg	mercury
HHB	hexachloro benzene
Hz	Hertz
IPM	integrated pest management
LTF	log transfer facilities
LWD	large woody debris
mm	millimeter
m/s <sup>2</sup>	meters per second squared
Magnuson-Stevens Act	Magnuson-Stevens Fishery Conservation and Management Act

MARPOL	International Convention for the Prevention of Pollution from Ships
NAWQA	National Water Quality Assessment
NEPA	National Environmental Policy Act
nm	nautical mile
NMDMP	National Marine Debris Monitoring Program
NMFS	National Marine Fishery Service
NOAA	National Oceanic and Atmospheric Administration
NPDES	National Pollution Discharge Elimination System
NPPC	Northwest Power Planning Council
NRC	National Research Council
OCS	outer coastal shelf
OGTAD	Oil and Gas Technologies for the Arctic and Deep Water
OWRRI	Oregon Water Resources Research Institute
PAH	polyaromatic hydrocarbon
Pb	lead
PBDE	polybrominated diphenyl ether
PCB	polychlorinated biphenyls
PFMC	Pacific Fishery Management Council
PNPCC	Pacific Northwest Pollution Control Council
POPs	persistent organic pollutants
RPWAST	Rich Passage Wave Action Study Team
SAIC	Science Applications International Corporation
SCS	Soil Conservation Service
SPL	sound pressure levels
SSC	suspended sediment concentration
TSS	total suspended solids
USACE	U.S. Army Corps of Engineers
USDA	U.S. Department of Agriculture
USFWS	U.S. Fish and Wildlife Service
USGS	U.S. Geological Survey
WCS	water control structure
WDFW	Washington State Department of Fish and Wildlife
ZOD	zone of deposit

## 1.0 INTRODUCTION

### Background on Essential Fish Habitat

In 1996, the United States Congress added new habitat conservation provisions to the Magnuson-Stevens Fishery Conservation and Management Act (Magnuson-Stevens Act), the federal law that governs United States marine fisheries management. The renamed Magnuson-Stevens Act mandated the identification of Essential Fish Habitat<sup>1</sup> (EFH) for federally managed species and consideration of measures to conserve and enhance the habitat necessary for these species to carry out their life cycles.

The Magnuson-Stevens Act requires federal agencies to consult with the National Marine Fisheries Service (NMFS) on all actions or proposed actions permitted, funded, or undertaken by the agency that may adversely affect<sup>2</sup> EFH. Federal agencies initiate consultation by preparing and submitting a written assessment of the effects of the proposed federal action on EFH to NMFS. If a federal action agency determines that an action will not adversely affect EFH, no consultation is required. To promote efficiency and avoid duplication, EFH consultation is usually integrated into existing environmental review procedures under other laws such as the National Environmental Policy Act (NEPA), Endangered Species Act (ESA), or Fish and Wildlife Coordination Act.

The Magnuson-Stevens Act requires NMFS to recommend conservation measures to federal and state agencies regarding actions that would adversely affect EFH. These EFH conservation recommendations are advisory, not mandatory, and may include measures to avoid, minimize, mitigate, or otherwise offset the adverse effects to EFH. Within 30 days of receiving NMFS' conservation recommendations, federal action agencies must provide a detailed response in writing. The response must include measures proposed for avoiding, mitigating, or offsetting the impact of a proposed activity on EFH. State agencies are not required to respond to EFH conservation recommendations. If a federal action agency chooses not to adopt NMFS' conservation recommendations, it must provide an explanation. Examples of federal action agencies that permit or undertake activities that may trigger EFH consultation include, but are not limited to, the U.S. Army Corps of Engineers (USACE), Environmental Protection Agency (EPA), Federal Energy Regulatory Commission (FERC), and Department of the Navy (DoN). Fishery Management Councils (FMCs) may also choose to comment on proposed actions that may adversely affect EFH.

### Significance of Essential Fish Habitat

The waters and substrate that comprise EFH designations under the jurisdiction of the FMCs are diverse and widely distributed. They are also closely interconnected with other aquatic and terrestrial environments.

---

<sup>1</sup> EFH is defined as "those waters and substrate necessary to fish for spawning, breeding, feeding, or growth to maturity." "Waters" include aquatic areas and their associated physical, chemical, and biological properties. Substrate includes sediment underlying the waters. "Necessary" means the habitat required to support a sustainable fishery and the managed species' contribution to a healthy ecosystem. "Spawning, breeding, feeding, or growth to maturity" covers all habitat types utilized by a species throughout its life cycle.

<sup>2</sup> An adverse effect is any impact that reduces the quality and/or quantity of EFH. Adverse effects may include direct or indirect physical, chemical, or biological alterations of the waters or substrate and loss of, or injury to, benthic organisms, prey species, and their habitat, as well as other ecosystem components. Adverse effects may be site-specific or habitat-wide impacts, including individual, cumulative, or synergistic consequences of actions [50 CFR 600.910(a)]

From a broad perspective, EFH is the geographic area where the species occurs at any time during its life. This area can be described in terms of ecological characteristics, location, and time. Ecologically, EFH includes waters and substrate that focus distribution (e.g., migration corridors, spawning areas, rocky reefs, intertidal salt marshes, or submerged aquatic vegetation) and other characteristics that are less distinct (e.g., turbidity zones, salinity gradients). Spatially, habitats and their use may shift over time due to climate change, human activities, geologic events, and impacts. The type of habitat available, its attributes, and its functions are important for species productivity, diversity, health, and survival.

The following discussion addresses non-fishing activities that may adversely affect EFH. They are grouped into four different systems in which the activities usually occur: upland, river or riverine, estuary or estuarine, and coastal or marine. Riverine habitats provide important habitat that serves multiple purposes for anadromous species such as salmon. These purposes include migration, feeding, spawning, nursery, and rearing functions. Protecting these functions is key to providing for a productive system and a healthy fishery. The riparian corridor is an important component of a river system. The term “riparian” refers to the land directly adjacent to a stream, lake, or estuary. A healthy riparian area has vegetation harboring prey items (e.g., insects), contributes necessary nutrients, provides large woody debris (LWD) that creates channel structure and cover for fish, and provides shade, which controls stream temperatures (Bilby and Ward 1991). When vegetation is removed from riparian areas, waters are heated, and LWD is less common. This results in less refuge for fish, fundamental changes in channel structure (e.g., loss of pool habitats), instability of streambanks, and alteration of nutrient and prey sources within the river system.

Estuaries are the bays and inlets influenced by both the ocean and rivers, and they serve as the transition zone between freshwater and saltwater (Botkin et al. 1995). Estuaries support a community of plants and animals that are adapted to the zone where freshwater and saltwater mix (Zedler et al. 1992). Estuarine habitats fulfill fish and wildlife needs for reproduction, feeding, refuge, and other physiological necessities (Simenstad et al. 1991, Good 1987, Phillips 1984). Healthy estuaries include eelgrass beds that protect young fish from predators, provide habitat for fish and wildlife, improve water quality, and control sediments (Thayer et al. 1984, Hoss and Thayer 1993, Phillips 1984). In addition, mud flats, high salt marsh, and saltmarsh creeks also provide productive shallow-water habitat for epibenthic fishes and decapods (Sogard and Able 1991).

Coastal or marine habitats comprise a variety of broad habitat types for EFH-managed species, including sand bottoms, rocky reefs, and submarine canyons. When rock reefs support kelp stands, they become exceptionally productive. Relative to other habitats, including wetlands, shallow and deep sand bottoms, and rock bottom artificial reefs, giant kelp habitats are substantially more productive in the fish communities they support (Bond et al. 1999). Foster and Schiel (1985) reported that the net primary productivity of kelp beds may be the highest of any marine community. Lush kelp forest communities (e.g., giant kelp, bull kelp, elk kelp, and feather boa kelp) are found relatively close to shore along the open coast. These subtidal communities provide vertically structured habitat through the water column on the rocky shelf made up of a canopy of tangled stipes from the water line to a depth of 10 feet; a mid-kelp, water-column region; and the bottom, holdfast region. The stands provide nurseries, feeding grounds, and/or shelter to a variety of groundfish species and their prey (Feder et al. 1974, Ebeling et al. 1980).

### Non-fishing Impacts

The diversity, widespread distribution, and ecological linkages with other aquatic and terrestrial environments make the waters and substrates that comprise EFH susceptible to a wide array of human activities unrelated to fishing.

Non-fishing activities have the potential to adversely affect the quantity or quality of EFH in riverine, estuarine, and marine systems. Broad categories of such activities include, but are not limited to, mining, dredging, fill, impoundment, discharge, water diversions, thermal additions, actions that contribute to nonpoint source pollution and sedimentation, introduction of potentially hazardous materials, introduction of exotic species, and the conversion of aquatic habitat that may eliminate, diminish, or disrupt the functions of EFH. For each activity, known and potential adverse impacts to EFH are described in this document. The descriptions explain the mechanisms or processes that may cause the adverse effects and how these may affect habitat function.

Non-fishing activities discussed in this document are subject to a variety of regulations and restrictions under federal, state, and local laws designed to limit environmental impacts. Many of these existing requirements help to avoid or minimize adverse effects to aquatic habitats, including EFH. The conservation recommendations contained in this document are rather general and may overlap with certain existing standards for specific development activities. Nevertheless, the recommendations highlight practices that can help to avoid and minimize adverse effects to EFH. During EFH consultations between NMFS and other agencies, NMFS strives to provide reasonable and scientifically based recommendations that account for restrictions imposed under various state and federal laws by agencies with appropriate regulatory jurisdiction. NMFS will not recommend that state or federal agencies take actions beyond their statutory authority, and NMFS' EFH conservation recommendations are not binding.

The conservation measures discussed in this document should be viewed as options to avoid, minimize, or compensate for adverse impacts and promote the conservation and enhancement of EFH. Ideally, non-water-dependent actions should not be located in EFH if such actions may have adverse impacts on EFH. Activities that may result in significant adverse effects on EFH should be avoided where less environmentally harmful alternatives are available. If there are no alternatives, the impacts of these actions should be minimized. Environmentally sound engineering and management practices should be employed for all actions that may adversely affect EFH. If avoidance or minimization is not possible, or will not adequately protect EFH, compensatory mitigation to conserve and enhance EFH is recommended.

#### Purpose of the Document

NMFS biologists should review proposed projects under the EFH provisions to ensure that they provide appropriate EFH conservation recommendations. During consultation, it is challenging to consider all potential non-fishing impacts to EFH so that the appropriate mix of recommendations can be made. Because impacts that may adversely affect EFH can be direct, indirect, and cumulative, the biologist must consider and analyze these interrelated impacts. Consequently, it is not unusual for particular impacts to be overlooked or discounted during a consultation. This document was prepared to aid NMFS' biologists reviewing proposed projects as they consider potential impacts that may adversely affect EFH and to provide consistent and substantiated EFH conservation recommendations. The document should also be useful for federal action agencies undertaking EFH consultations, especially in preparing EFH assessments.

This document is organized by activities that may potentially affect EFH occurring in four discrete ecosystems: upland, riverine, estuarine, and coastal/marine. The separation of these ecosystems is artificial, and many of the impacts and related activities are not exclusive to one system. For instance, sand and gravel mining activities often occur in riverine systems, but also take place in estuarine systems. Thus, the activities are located in the ecosystem where they initially occur in a watershed progression, while secondary impacts may be addressed elsewhere. For example, pile driving creates its own set of unique impacts to EFH. While installing piles, however, other construction may cause activities such as



dredging to occur, which brings its own set of potential adverse impacts. In addition, some of the activities discussed may not be applicable to all three regions, either due to geographic location or because the industry associated with the impact does not occur in that region.

The conservation recommendations included with each activity present a series of site-specific measures that can be undertaken by the action agency to avoid, offset, or mitigate impacts to EFH. Not all of these suggested measures are necessarily applicable to any one project or activity that may adversely affect EFH. More specific or different measures based on the best and most current scientific information may be developed before, or during, EFH consultations and communicated to the appropriate agency. The conservation recommendations provided represent a short menu of general types of conservation actions that can contribute to the conservation, enhancement, and proper functioning of EFH.

#### Overall Approach and Comparison to Previous Analyses

The 1999 EFH Environmental Assessment (EA) limited the scope of the discussion of non-fishing threats to coastal activities with some references to possible offshore impacts from non-fishing activities. The EFH EA categorized the non-fishing impacts to EFH in Alaska by lumping together several types of activities that may or may not occur together and calling out specific activities that cause habitat alteration. For example, the EFH EA lumped together dredging, fill, and excavation, providing a combined narrative discussion and analysis of these activities as they relate to port construction and support activities. These activities often occur independently, and the possible impacts to EFH that may occur from one activity (e.g., dredging) differ from those that may associated with another (e.g., fill). In contrast, the EFH EA's discussion and analysis of possible adverse impacts from certain activities (e.g., mining) were limited to activities that occur in the marine environment without addressing the same activities in other areas with potential adverse effects on EFH (such as anadromous streams).

The format for discussion of non-fishing threats in the EFH EA summarized potential impacts from each activity and then provided an expanded discussion with general conservation recommendations for some of the activities. In addition, an attached worksheet provided a professional interpretative summary of the broad category of threats discussed in the Non-fishing Adverse Impacts section. Habitat conservation and enhancement recommendations were provided in tabular format starting with a broad category of habitats (i.e., near shore habitat and waters [0 to 3 nm], pelagic habitat and waters [312 nm], and offshore habitat and waters [more than 12 nm ]) with general recommendations as they relate to a particular area or habitat type and associated managed species. See Tables 1 and 2, which are reproduced from the EFH EA.

Comments received on the EFH EA indicated that impacts to EFH can be direct, indirect, and cumulative. While it is necessary to distinguish between activities to identify possible adverse impacts, it is equally important to consider and analyze these activities as they interrelate within habitats. Appendix G to the EFH EIS, therefore, takes more of an ecosystem perspective and provides more detail and a different format than the non-fishing impacts section of the EFH EA.

This document is organized by activities that may potentially impact EFH occurring in four discrete ecosystems. The separation of these ecosystems is artificial, and many of the impacts and their related activities are not exclusive to one system. For instance, as recognized in the discussion of the EFH EA, activities such as sand and gravel mining occur in riverine, estuarine, and marine systems. Because activities are discussed in the section corresponding to the primary ecosystem where they occur, readers should use the Master Index at the end of the document to identify other systems where such activities may also take place. Also, certain activities (e.g., pile driving) have specific potential impacts to EFH and may be associated with other construction activities (e.g., dredging) that have their own potential impacts. Readers should use the Master Index to ensure that all activities for a given project are considered.

**Table 1.** Recommendations from the 1999 EFH EA Regarding Non-fishing Threats

<b>Recommendation</b>	<b>Area</b>	<b>Species</b>
<b>Near Shore Habitat and Waters (0-3nm)</b>		
Minimize construction of structures such as causeways or breaches that would affect local flushing, water temperatures, water quality, lateral drift, and/or migration.	Sensitive areas, special aquatic and vegetation areas	groundfish, salmon, scallop, crab
Minimize construction of structures such as docks that ground on tidal lands during low water events.	Sensitive areas, special aquatic and vegetation areas	groundfish, salmon, crab
Minimize deposition of fill in tidelands.	Sensitive areas, special aquatic and vegetation areas	groundfish, salmon, crab
Stage rapid response equipment, and establish measures for accidental impacts such as oil and hazardous material spills.	ports, sensitive areas	groundfish, salmon, scallop, crab
Monitor point source pollution sites such as fish processing waste, sewage, and storm water runoff outfalls.	ports, vessel processors, communities	groundfish, salmon, scallop, crab
Minimize disposal or dumping of dredge spoils, drilling muds, and municipal and industrial wastes.	known concentration of bottom species and their habitats	groundfish, salmon, scallop, crab
Test dredge spoils prior to marine disposal.	port and upland sources	groundfish, salmon, scallop, crab
Establish monitoring that incorporates federal and state regulatory agency determinations, i.e., tracking database and GIS system.	areawide	groundfish, salmon, scallop, crab
<b>Pelagic Habitat and Waters (3-12nm)</b>		
Assess cumulative oil and gas production activities.	BSAI, Chukchi Sea, OCS, Cook Inlet, GOA	groundfish, salmon, scallop, crab
Identify marine disposal sites.	areawide	groundfish, salmon, scallop, crab
Establish monitoring that incorporates Federal and State regulatory agency determinations, i.e., tracking database and GIS system	areawide	groundfish, salmon, scallop, crab
Establish no-discharge zones for ballast waters to prevent introduction of nonindigenous species and chemical contaminants.	ports, known gyre areas	groundfish, salmon, scallop, crab
Minimize disposal or dumping of dredge spoils, drilling muds, and municipal and industrial wastes.	known concentration of bottom species and their habitats	groundfish, salmon, scallop, crab
<b>Offshore Habitat and Waters (&gt;12 nm)</b>		
Establish monitoring that incorporates federal and state regulatory agency determinations, i.e., tracking database and GIS system	areawide	groundfish, salmon, scallop, crab
Establish no-discharge zones for ballast waters to prevent introduction of nonindigenous species and chemical contaminants.	known offshore gyre areas	groundfish, salmon, scallop, crab
Minimize disposal or dumping of dredge spoils, drilling muds, and municipal and industrial wastes.	known concentration of bottom species and their habitats	groundfish, salmon, scallop, crab

**Table 2.** Summary of Non-fishing Threats to EFH from the 1999 EFH EA

Threats	HABITAT ALTERATION										TOPOGRAPHIC ALTERATION										ORGANISM ALTERATION										OCEANOGRAPHIC ALTERATION										WATER QUALITY ALTERATION									
	Alteration of Physical Habitat	Alteration of Biological Habitat	Alteration of Chemical Habitat	Alteration of Physical Habitat	Alteration of Biological Habitat	Alteration of Chemical Habitat	Alteration of Physical Habitat	Alteration of Biological Habitat	Alteration of Chemical Habitat	Alteration of Physical Habitat	Alteration of Biological Habitat	Alteration of Chemical Habitat	Alteration of Physical Habitat	Alteration of Biological Habitat	Alteration of Chemical Habitat	Alteration of Physical Habitat	Alteration of Biological Habitat	Alteration of Chemical Habitat	Alteration of Physical Habitat	Alteration of Biological Habitat	Alteration of Chemical Habitat	Alteration of Physical Habitat	Alteration of Biological Habitat	Alteration of Chemical Habitat	Alteration of Physical Habitat	Alteration of Biological Habitat	Alteration of Chemical Habitat	Alteration of Physical Habitat	Alteration of Biological Habitat	Alteration of Chemical Habitat	Alteration of Physical Habitat	Alteration of Biological Habitat	Alteration of Chemical Habitat																	
	Physical	Biological	Chemical	Physical	Biological	Chemical	Physical	Biological	Chemical	Physical	Biological	Chemical	Physical	Biological	Chemical	Physical	Biological	Chemical	Physical	Biological	Chemical	Physical	Biological	Chemical	Physical	Biological	Chemical	Physical	Biological	Chemical	Physical	Biological	Chemical																	
Excavation																																																		
Dredging		X			X	X	X	X	X			X	X	X		X	X					X		*	*	*		*	X	X	X																			
Dredge Material Disposal		X	X		X	X	X	X	X			X	X			X	X	X				X	X		*	*	*		*	X	X	X																		
Marine Mining		X	X		X			X	X			X	X	X		X	X					X	X		X	X	X		*	X	X	X																		
Nearshore Mining		X			X	X	X		X	X			X	X	X		X	X				X	X		*	*	*		*	X	X	X																		
Recreational Uses																																																		
Boating				X	X	X	X			X						X	X				X	X			*	*	*		*	*	X	X	X																	
Streambank Overusage		X						X	X	X			X	X	X		X	X				X	X						X	X	X																			
Fish Waste Processing																																																		
Shoreside Discharge		X			X	X	X		X	X			X	X								X	X		X	X			*	X	X	X																		
Vessel Discharge				X		X			X													X	X						*	X		X																		
Aquaculture					X		X		X								X	X	X	X	X		X	X	X			*	X	X	X																			
Petroleum Production																																																		
Production Facility		X	X		X	X	X	X	X			X	X	X			X	X				X	X		X	X	X				X	X	X																	
Exploration		X	X		X	X	X	X				X	X	X			X	X				X	X				X				X	X	X																	
Oil Spill		X	X		X	X	X	X	X							X	X		X			X	X			*			X		X	X	X																	
Hydrological																																																		
Hydroelectric Dams								X									X					X	X						X			X																		
Impoundments		X				X	X	X				X	X	X			X	X				X	X						X	X		X																		
Flood Erosion/Control		X			X		X	X	X			X	X	X			X	X				X							X	X																				
Agricultural																																																		
Agricultural/Farming		X			X		X	X	X	X			X	X	X				X			X			*	*			X	X	X	X																		
Insect Control					X		X	X								X	X					X	X								X		X																	
Forestry		X			X		X	X	X	X			X	X	X		X		X			X			X	*					X	X																		
Water Diversion/Withdrawal		X			X		X	X				X	X	X								X			*		X		X	X	X	X																		

**Table 2.** Summary of Non-fishing Threats to EFH from the 1999 EFH EA (continued)

Threats	HABITAT ALTERATION										TOPOGRAPHIC ALTERATION										ORGANISM ALTERATION										OCEANOGRAPHIC/WATER QUALITY ALTERATION									
	Alteration	Loss of Function	Loss of Structure	Loss of Species	Loss of Habitat	Loss of Function	Loss of Structure	Loss of Species	Loss of Habitat	Alteration	Loss of Function	Loss of Structure	Loss of Species	Loss of Habitat	Alteration	Loss of Function	Loss of Structure	Loss of Species	Loss of Habitat	Alteration	Loss of Function	Loss of Structure	Loss of Species	Loss of Habitat	Alteration	Loss of Function	Loss of Structure	Loss of Species	Loss of Habitat	Alteration	Loss of Function	Loss of Structure	Loss of Species	Loss of Habitat						
	Harbors/Ports/Marinas	Port Construction	Port Development	Artificial Reefs	Municipal and Industrial	Non-point Source	Coastal Urbanization	Sewage Treatment	Storm Water Runoff	Environmental	Climate Changes/Shifts	Toxic Algal Bloom	Introduction of Exotic Species	Marine Transportation	Vessel Groundings	Ballast Water	Marine Debris																							

\* - Short-term impact

Similar to the non-fishing impacts section of the EFH EA, this document is not meant to provide an exhaustive review. This document is, however, a result of a collaborative effort among the NMFS Alaska Region, Northwest Region, and Southwest Region and the respective Fisheries Science Centers, which provided a broader range of expertise to reach consensus regarding the general conservation recommendations.

The format selected for presenting the information in this document was chosen because it could be easily adapted for use by all three regions. It provides an introductory description of each activity, identification of potential adverse impacts, and suggested general conservation measures that would help minimize and avoid adverse effects of non-fishing activities on EFH. Section 3.4.4 of the EFH EIS recognizes that some of the activities discussed in Appendix G may not be applicable to Alaska. To better reflect the Alaska perspective, Table 3.4-36 of the EIS identifies the categories from Appendix G and correlates them with possible changes in physical, chemical, and biological parameters, and Table 3.4-37 takes the same categories from Appendix G and broadly interprets whether the effects from the activities in Alaska have been positive, insignificant, negative, or unknown.

## **2.0 UPLAND ACTIVITIES**

### **2.1 Nonpoint Source Pollution**

The information in this section is adapted from EPA 1993.

Nonpoint source pollution generally results from land runoff, precipitation, atmospheric deposition, seepage, or hydrologic modification. Technically, the term nonpoint source means anything that does not meet the legal definition of point source in Section 502(14) of the Clean Water Act (CWA), which refers to discernable, confined, and discrete conveyance from which pollutants are or may be discharged. The major categories of nonpoint pollution are agricultural runoff, urban runoff, including developed and developing areas (see Section 2.2), silvicultural (forestry) runoff (see Section 2.1.2), marinas and recreational boating, road construction, and channel and streambank modifications, including channelization, channel modifications (see Section 4.7), and streambank and shoreline erosion.

Nonpoint source pollution is usually lower in intensity than an acute point source event, but may be more damaging to fish habitat in the long term. Nonpoint source pollution is often difficult to detect. It may affect sensitive life stages and processes, and the impacts may go unnoticed for a long time. When severe population impacts are finally noticed, they may not be tied to any one event; hence, it may be difficult to correct, clean up, or mediate.

#### **2.1.1 Agricultural/Nursery Runoff**

Substantial portions of croplands and commercial nursery operations are connected to inland and coastal waters where nonpoint pollution can have a direct adverse effect on aquatic habitats. Tillage aerates the upper soil, but compacts fine textured soils just below the depth of tillage, thus altering infiltration. Use of farm machinery on cropland and adjacent roads causes further compaction, reducing infiltration and increasing surface runoff. Agricultural lands are also characterized by poorly maintained dirt roads and ditches that, along with drains, route sediments, nutrients, and pesticides directly into surface waters. Natural channels filter and process pollutants. In many instances, roads ditches and drains have replaced headwater streams, and these constructed systems deliver pollutants directly to surface waters (Larimore and Smith 1963).

Rangeland soils can also become compacted by livestock with similar effects on runoff (Platts 1991, Heady and Child 1994). Compaction of rangelands generally increases with grazing intensity, although

site-specific soil and vegetative conditions are important (Kauffman and Krueger 1984, Heady and Child 1994). Johnson (1992) reviewed studies related to grazing and hydrologic processes and concluded that heavy grazing nearly always decreases infiltration, reduces vegetative biomass, and increases bare soil. Primary runoff pollutants are nutrients, pesticides, sediment, salts, and animal wastes. Because the primary routes of pesticide transport to EFH include not only surface runoff events, but also direct application, aerial drift, and groundwater systems, pesticide contamination is addressed separately in Section 2.1.3.

#### Potential Adverse Impacts

Adverse impacts to EFH from agricultural and nursery runoff can result from (1) nutrient loading, (2) introduction of animal wastes, (3) erosion, and (4) sedimentation.

Nutrients are applied to agricultural land in several different forms and come from various sources, including commercial fertilizers; manure from animal production facilities (with bedding and other wastes added to the manure); municipal and industrial treatment plant effluent and sludge; legume and crop residues; irrigation water; and atmospheric deposition of nutrients, such as nitrogen and sulfur. Specifically, nitrogen and phosphorus are the two major nutrients from agricultural land that degrade water quality. Introduction of these nutrients into aquatic systems can dramatically increase aquatic plant productivity and decay (Waldichuk 1993). This process can increase turbidity, temperature, and the accumulation of dead organic material. It can also decrease light penetration, oxygen, and the growth of submerged aquatic vegetation. These alterations can result in the destruction of habitat for small or juvenile fish and severely impair biological food chains.

Animal waste (manure) includes fecal and urinary wastes of livestock and poultry; process water (such as from a milking parlor); and the feed, bedding, litter, and soil with which they become intermixed. Because riparian areas are favored by cattle, nutrients consumed elsewhere are often excreted as waste in riparian zones (Heady and Child 1994). Pollutants contained in manure and associated bedding materials can be transported into marine environments by runoff and process wastewater from rangelands, pastures, or confined animal facilities. These pollutants may include oxygen-demanding substances, such as nitrogen, phosphorus, and organic solids; salts; bacteria, viruses, and other microorganisms, as well as sediments that increase organic decomposition. Runoff of animal wastes can cause fish kills due to ammonia. Solids deposited into the marine environment can reduce productivity over extended periods of time due to the accelerated effects of cultural eutrophication. Runoff can be accelerated by grazing processes that remove or disturb riparian vegetation and soils.

Sediment is the result of erosion. Sheet, rill, and gully erosion all transport fine sediment, enriched with a wide variety of attached pollutants, from agricultural land into the aquatic environment. The presence of livestock in the riparian zone accelerates sediment transport rates by increasing both surface erosion and mass wasting (Platts 1991, Marcus et al. 1990, Heady and Child 1994). Likewise, grazing in uplands can result in increased sediment delivery through channelized flows. For example, the Soil Conservation Service (SCS) estimated that 92 percent of the total sediment yields in the Snake and Walla Walla River basins of southeastern Washington resulted from sheet and rill erosion from cropland accounting for only 43 percent of total land area (SCS et al. 1984). Increased sediment in aquatic systems can increase turbidity, reduce light penetration, smother fish spawning areas and food supplies, clog the filtering capacity of filter feeders, clog and harm the gills of fish, interfere with feeding behaviors, and significantly lower overall biological productivity.

Salts are a product of natural weathering of soil and geologic material. The movement and deposition of salts depend on the amount and distribution of rainfall and irrigation, the soil and underlying strata, evapotranspiration rates, and other environmental factors. Irrigation water, whether from ground or

surface water sources, has a natural base load of dissolved mineral salts. As water is consumed by plants or lost to the atmosphere by evaporation, the remaining salts become concentrated in the soil (the concentrating effect). Thus, the total salt load carried by irrigation return flow is the sum of the salts remaining in the applied water plus any additional salt picked up from the irrigated land. Irrigation return flows convey the salt to the receiving streams or groundwater reservoirs. If the amount of salt in the return flow is low in comparison to the total stream flow, water quality may not be degraded to the extent that EFH functions are impaired. If the process of water diversion and the return flow of saline drainage water is repeated many times along a stream or river, downstream habitat quality can become progressively degraded.

Groundwater is also susceptible to nutrient contamination in agricultural lands composed of sandy or other coarse-textured soil (Franco et al. 1994, USGS 1999). Nitrate, a highly soluble form of nitrogen, can leach rapidly through the soil profile and accumulate in groundwater, especially in shallow zones (Jordan and Weller 1996, Brady and Weil 1996). This groundwater can be a significant source of nutrients in surface waters when discharged through seeps, drains, or by direct subsurface flow to water bodies (Lee and Taylor 2000).

### Recommended Conservation Measures

The recommended conservation measures for agricultural/nursery runoff include the following:

1. Protect and restore soil quality with controls that affect the ability of the soil to grow crops, partition and regulate water flow, and act as an environmental filter (e.g., permeability, water holding capacity, nutrient availability, organic matter content, and biological activity). Relevant practices include cover cropping, crop sequence, conservation tillage, crop residue management, grazing management, and use of low-impact equipment (e.g., minimally sized, rubber tired).
2. Improve land use efficiencies for key agricultural inputs, including nitrogen, phosphorus, pesticides, and irrigation water. Relevant practices are agronomic nutrient applications based upon nutrient testing, including manure, during clear weather, use of integrated pest management, and irrigation management.
3. Increase resistance to soil erosion and runoff. Sediment basins, contour farming, and grazing management are examples of key practices.
4. Protect and restore rangelands using practices, such as rotational grazing systems or livestock distribution controls, exclusion from riparian and aquatic areas, livestock-specific erosion controls, reestablishment of vegetation, or extensive brush management correction.
5. Increase field and landscape buffers to provide cost-effective protection against the cumulative effects of many small, but unavoidable, pollutant discharges associated with an active agricultural enterprise and the kinds of catastrophic pollution that can be associated with the high energy flows and runoff associated with episodic storms. The full range of agricultural buffer practices (e.g., riparian forests, alley cropping, contour buffer strips, crosswind trap strips, field borders, filter strips, grassed waterways with vegetative filters, herbaceous wind barriers, vegetative barriers, and windbreak/shelterbelts) has to be systematically deployed, protected, and managed across the agricultural landscape or overall aquatic habitat improvements will be minimal.
6. Optimize siting of new confined animal facilities or expansion of existing facilities by placing them away from riparian areas, surface water, and areas with high leaching potential to surface or groundwater. Ensure that adequate nutrient and wastewater collection facilities are in place. Ensure that sufficient cropland is available for agronomic application of animal wastes.
7. Consider using restored wetlands to reduce contamination from a variety of sources, including nitrogen, phosphorus, suspended solids, biochemical oxygen demand (BOD), trace metals, trace organics, and pathogens. Larger wetland systems relative to the amount of land that is drained with longer retention times (at least 1 to 2 weeks) are most beneficial at improving water quality.

Wetlands located within riparian buffer strips provide the most effective pollution removal by combining different treatment methods.

### **2.1.2 Silviculture/Timber Harvest**

Recent revisions of Alaska's federal and state timber harvest regulations and best management practices (BMPs) have resulted in increased protection of EFH on federal, state, and private timber lands (Alaska Forest Resources and Practices Act [FRPA, AS 41.17], the Tongass Land Management Plan [<http://www.fs.fed.us/r10/tongass/management%20news/tlmp/tlmp.shtml>], and the Chugach Land and Resources Management Plan [<http://www.geographynetwork.com/chugach/>]). The Tongass and Chugach forest management plans provide for multiple uses on national forest lands and include protective measures on lands designated for timber production and on less intensively managed lands. The FRPA and its regulations set riparian buffers and establish mandatory BMPs for timber harvesting, road construction, road maintenance, and reforestation to protect water quality on state and privately owned timber production lands. The FRPA is also the standard for compliance with federal Coastal Zone and Clean Water Act requirements in Alaska.

Although current forest management practices are designed to avoid or minimize adverse effects to fish habitat, the harvest and cultivation of timber and other forestry products are major activities that can have both short- and long-term impacts throughout many coastal watersheds and estuaries if management practices are not fully implemented or effective. Timber harvest removes the dominant vegetation, converts mature and old-growth upland and riparian forests to tree stands or forests of early seral stage, reduces permeability of soils and increases the area of impervious surfaces, increases sedimentation from surface runoff and mass wasting processes, results in altered hydrologic regimes, and impairs fish passage through inadequate design, construction, and/or maintenance of stream crossings.

Deforestation associated with timber harvest can alter or impair instream habitat structure and watershed function. Timber harvest may result in inadequate or excessive surface and stream flows, increased stream bank and stream bed erosion, loss of complex instream habitats, sedimentation of riparian habitat, and increased surface runoff with associated contaminants (e.g., herbicides, fertilizers, fine sediments). Hydrologic characteristics, (e.g., water temperature, annual hydrograph) change, and greater variation in stream discharge is associated with timber harvest. Alterations in the supply of LWD and sediment can have negative effects on the formation and persistence of instream habitat features. Excess debris in the form of small wood and silt can smother benthic habitat and reduce dissolved oxygen levels.

#### Potential Adverse Impacts

Four major categories of activities can adversely affect EFH: 1) construction of logging roads, 2) creation of fish migration barriers, 3) removal of streamside vegetation, and 4) disturbance associated with log transfer facilities (LTFs) (see Section 4.9).

Logging road construction can destabilize slopes and increase erosion and sedimentation (see Road Building and Maintenance, Section 2.3). Two major types of erosion occur: mass wasting and surface erosion. Mass movement of soils, commonly referred to as landslides or debris slides, is associated with timber harvest and road building on high hazard soils and unstable slopes. Both frequency and size of debris slides are increased when logging roads are built on, or timber is harvested from, these unstable land forms. The result is increased erosion and sediment deposition in downslope waterways. Erosion from roadways is most severe when poor construction practices are employed that do not include properly located, sized, and installed culverts; proper ditching; and ditch blocker water bars (Furniss et al. 1991).



Stream crossings (bridges and culverts) on forest roads are often inadequately designed, installed, and maintained, and they frequently result in full or partial barriers to both the upstream and downstream migration of adult and juvenile fish. Perched and undersized culverts can accelerate stream flows to the point that these structures become velocity barriers for migrating fish. Blocked culverts result from installation of undersized culverts or inadequate maintenance to remove debris. Blocked culverts can result in displacement of the stream from the downstream channel to the roadway or roadside ditch, resulting in dewatering of the downstream channel and increased erosion of the roadway. Culverts and bridges deteriorate structurally over time. Failure to replace or remove them at the end of their useful life may cause partial or total blockage of fish passage. Caution should be used, however, when removing culverts. Channel incision can often occur downstream of a culvert and generally moves upstream. An existing culvert can act as a grade control, halting the upstream progression of a headcut and causing further channel regrade (Castro 2003). The unchecked upstream progression of a headcut can cause further damage to EFH.

Removing streamside vegetation increases the amount of solar radiation reaching the stream and can result in warmer water temperatures, especially in small, shallow streams of low velocity. In southeast Alaska, Meehan et al. (1969) found that maximum temperature in logged streams without riparian buffers exceeded that of unlogged streams by up to 5°C, but did not reach lethal temperatures. However, the increased water temperatures often exceeded optimum temperatures for pink and chum salmon (Reiser and Bjornn 1979). Logged streams have been associated with higher water temperatures, lower base flows and higher peak flows, and low oxygen levels that have resulted in significant mortalities of pink and chum salmon (Flanders and Cariello 2000). In cold climates, the removal of riparian vegetation can result in lower water temperatures during winter, increasing the formation of ice and damaging and delaying the development of incubating fish eggs and alevins.

By removing vegetation, timber harvest reduces transpiration losses from the landscape and decreases the absorptive capability of the groundcover. These changes result in increased surface runoff during periods of high precipitation and decreased base flows during dry periods. Reduced soil strength results in destabilized slopes and increased sediment and debris input to streams (Swanston 1974). Sediment deposition in streams can reduce benthic community production (Culp and Davies 1983), cause mortality of incubating salmon eggs and alevins, and reduce the amount of habitat available for juvenile salmon (Heifetz et al. 1996). Cumulative sedimentation from logging activities can significantly reduce the egg-to-fry survival of coho and chum salmon (Cederholm and Reid 1987, Myren and Ellis 1984.) Reductions in the supply of LWD also result when old-growth forests are removed, with resulting loss of habitat complexity that is critically important for successful salmonid spawning and rearing. (Bisson et al. 1988).

#### Recommended Conservation Measures

1. Implement best management practices (BMPs) for impacts affecting particular habitats and resulting from specific types of silviculture-related activities provided in the “Additional Resources” section.
2. Avoid timber operations to the extent practicable near streams with EFH. For the Alaska region, see the following links: Fish: Forest-Wide Standards and Guides:  
[http://www.fs.fed.us/r10/TLMP/F\\_PLAN/FPCHAP4.PDF](http://www.fs.fed.us/r10/TLMP/F_PLAN/FPCHAP4.PDF);  
<http://www.or.blm.gov/ForestPlan/newsandga.pdf>;  
<http://www.dnr.state.ak.us/forestry/pdfs/forpracregs.pdf>
3. Avoid timber operations to the extent practicable in wetlands contiguous with anadromous fish streams. See the following links: Wetlands: Forest-Wide Standards and Guides:  
[http://www.fs.fed.us/r10/TLMP/F\\_PLAN/FPCHAP4.PDF](http://www.fs.fed.us/r10/TLMP/F_PLAN/FPCHAP4.PDF)
4. Avoid timber operations to the extent practicable near estuary and beach habitats. See the following links: Beach and Estuary Fringe: Forest-Wide Standards and Guides:

[http://www.fs.fed.us/r10/TLMP/F\\_PLAN/FPCHAP4.PDF](http://www.fs.fed.us/r10/TLMP/F_PLAN/FPCHAP4.PDF);

[http://www.fs.fed.us/r10/chugach/forest\\_plan/forest\\_plan\\_web.pdf](http://www.fs.fed.us/r10/chugach/forest_plan/forest_plan_web.pdf)

5. Maintain riparian buffers along all streams. In the Alaska region, buffer width is site-specific and dependent on use by anadromous fish and stream process type. Stream process groups are described in the following link: [http://www.fs.fed.us/r10/TLMP/F\\_PLAN/APPEND\\_D.PDF](http://www.fs.fed.us/r10/TLMP/F_PLAN/APPEND_D.PDF). Standards and guidelines for riparian buffers for the Alaska region are described in the following links. Riparian Forest-Wide Standards and Guides: [http://www.fs.fed.us/r10/TLMP/F\\_PLAN/FPCHAP4.PDF](http://www.fs.fed.us/r10/TLMP/F_PLAN/FPCHAP4.PDF) and [http://www.fs.fed.us/r10/chugach/forest\\_plan/forest\\_plan\\_web.pdf](http://www.fs.fed.us/r10/chugach/forest_plan/forest_plan_web.pdf); FPRA riparian buffer regulations can be found at <http://www.dnr.state.ak.us/forestry/pdfs/fprachrt.pdf>
6. Incorporate watershed analysis into timber and silviculture projects. Particular attention should be given to the cumulative effects of past, present, and future timber sales within the watershed. See the following link on watershed analysis: [http://www.fs.fed.us/r10/TLMP/F\\_PLAN/APPEND\\_J.PDF](http://www.fs.fed.us/r10/TLMP/F_PLAN/APPEND_J.PDF)
7. Follow BMPs. See the following links on BMPs:  
[http://www.fs.fed.us/r10/TLMP/F\\_PLAN/APPEND\\_C.PDF](http://www.fs.fed.us/r10/TLMP/F_PLAN/APPEND_C.PDF);  
[http://www.fs.fed.us/r10/chugach/forest\\_plan/forest\\_plan\\_web.pdf](http://www.fs.fed.us/r10/chugach/forest_plan/forest_plan_web.pdf);  
<http://www.dnr.state.ak.us/forestry/pdfs/forpracregs.pdf>
8. For forest roads, see Section 2.3, Road Building and Maintenance. For the Alaska region, also see the following links: 1) transportation: forest-wide standards and guides  
[http://www.fs.fed.us/r10/TLMP/F\\_PLAN/FPCHAP4.PDF](http://www.fs.fed.us/r10/TLMP/F_PLAN/FPCHAP4.PDF) and 2) soils and water: forest-wide standards and guides: [http://www.fs.fed.us/r10/TLMP/F\\_PLAN/FPCHAP4.PDF](http://www.fs.fed.us/r10/TLMP/F_PLAN/FPCHAP4.PDF);  
[http://www.fs.fed.us/r10/chugach/forest\\_plan/forest\\_plan\\_web.pdf](http://www.fs.fed.us/r10/chugach/forest_plan/forest_plan_web.pdf);  
<http://www.dnr.state.ak.us/forestry/pdfs/forpracregs.pdf>

### **2.1.3 Pesticide Application**

More than 800 different pesticides are currently registered for use in the United States. Legal mandates covering pesticides are the CWA and the Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA). Water quality criteria for the protection of aquatic life have only been developed for a few of the currently used chemicals (EPA, Office of Pesticide Programs). Collectively, these substances are designed to repel, kill, or regulate the growth of undesirable biological organisms. This diverse group includes fungicides, herbicides, insecticides, nematocides, molluscicides, rodenticides, fumigants, disinfectants, repellents, wood preservatives, and antifoulants. The most common pesticides are insecticides, herbicides, and fungicides. These are used for pest control on forested lands, agricultural crops, tree farms and nurseries, highways and utility rights of way, parks and golf courses, and residences. Pesticides can enter the aquatic environment as single chemicals or complex mixtures. Direct applications, surface runoff, spray drift, agricultural return flows, and groundwater intrusions are all examples of transport processes that deliver pesticides to aquatic ecosystems.

Pesticides are frequently detected in freshwater and estuarine systems that provide EFH. Nationwide, the most comprehensive environmental monitoring efforts have been conducted by the USGS as part of the National Water Quality Assessment (NAWQA) Program. A variety of human activities, such as fire suppression on forested lands, forest site preparation, noxious weed control, right-of-way maintenance (roads, railroads, power lines, etc.), algae control in lakes and irrigation canals, various agricultural practices, riparian habitat restoration, and urban and residential pest control, result in contamination from these substances. It is important to note that the term pesticide is a collective description of hundreds of chemicals with different sources, different fates in the aquatic environment, and different toxic effects on fish and other aquatic organisms. Despite these variations, all current use pesticides are (1) specifically designed to kill, repel, or regulate the growth of biological organisms, and (2) intentionally released into the environment. Habitat alteration from pesticides is different from more conventional water quality parameters, such as temperature, suspended solids, or dissolved oxygen, because, unlike temperature or dissolved oxygen, the presence of pesticides can be difficult to detect due to limitation in proven

methodologies. This monitoring may also be expensive. However, as analytical methodologies have improved in recent years, the number of pesticides documented in fish and their habitats has increased.

### Potential Adverse Impacts

There are three basic ways that pesticides can adversely affect EFH. These are (1) a direct toxicological impact on the health or performance of exposed fish, (2) an indirect impairment of the productivity of aquatic ecosystems, and (3) a loss of aquatic vegetation that provides physical shelter for fish.

Fish kills are rare when pesticides are used according to their labels. For fish, the vast majority of effects from pesticide exposures are sublethal. Sublethal effects are a concern if they impair the physiological or behavioral performance of individual animals in ways that will decrease their growth or survival, alter migratory behavior, or reduce reproductive success. In addition to early development and growth, key physiological systems affected include the endocrine, immune, nervous, and reproductive systems. Many pesticides have been shown to impair one or more of these physiological processes in fish (Moore and Waring 2001). In general, however, the sublethal impacts of pesticides on fish health are poorly understood. Accordingly, this is a focus of recent and ongoing NOAA research (Scholz et al. 2000, Van Dolah et al. 1997).

The effects of pesticides on ecosystem structure and function can be a key factor in determining the cascading impacts of that chemical on fish and other aquatic organisms at higher trophic levels (Preston 2002). This includes impacts on primary producers (Hoagland et al. 1996) and aquatic microorganisms (DeLorenzo et al. 2001), as well as macroinvertebrates that are prey species for fish. For example, many pesticides are specifically designed to kill insects. Not surprisingly, these chemicals are relatively toxic to insects and crustaceans that inhabit river systems and estuaries. Overall, pesticides will have an adverse impact on fish habitat if they reduce the productivity of aquatic ecosystems. Finally, some herbicides are toxic to aquatic plants that provide shelter for various fish species. A loss of aquatic vegetation could damage nursery habitat or other sensitive habitats, such as eelgrass beds and emergent marshes.

### Recommended Conservation Measures

The recommended conservation measures for pesticide application include the following:

1. Incorporate integrated pest management (IPM) and BMPs as part of the authorization or permitting process to ensure the reduction of pesticide contamination in EFH (Scott et al. 1999).
2. Carefully review labels and ensure that application is consistent. Follow local, supplemental instructions such as county use bulletins where they are available.
3. Avoid the use of pesticides in and near EFH designated waters.
4. Refrain from areal spraying of pesticides on windy days.

## **2.2 Urban/Suburban Development**

The information in this section is adapted from NMFS 1998.

Urban growth and development in the United States continues to expand in coastal areas at a rate approximately four times greater than in other areas. The construction of urban, suburban, commercial, and industrial centers and corresponding infrastructure results in land use conversions typically resulting in vegetation removal and the creation of additional impervious surfaces. This runoff from impervious surfaces and storm sewers is the most widespread source of pollution into the Nation's waterways (EPA 1995).

## Potential Adverse Impacts

Development activities within watersheds and in coastal marine areas often impact the EFH of managed species on both long-term and short-term scales. Many of the impacts listed here are discussed in greater detail in other sections of this documents. The primary impacts include (1) the loss of riparian and shoreline habitat and vegetation, and (2) runoff. The removal of upland and shoreline vegetation removal can increase stream water temperatures, reduce supplies of LWD, and reduce sources of prey and nutrients to the water system. An increase in impervious surfaces, such as the addition of new roads (see also Section 2.3), roofs, bridges, and parking facilities, results in a decreased infiltration to groundwater and increased runoff volumes. This also has the potential to adversely affect water quality and water quantity/timing in downstream water bodies (i.e., estuaries and coastal waters).

The loss of riparian and shoreline habitat and vegetation can increase water temperatures and remove sources of cover. Such impacts can alter the structure of benthic and fish communities, resulting in an expected reduction in diversity and abundance of EFH species. Shoreline stabilization projects (see Section 4.7) that affect reflective wave energy can impede or accelerate natural movements of shoreline substrates, thereby affecting intertidal and sub-tidal habitats. Channelization of rivers cause loss of floodplain connectivity and simplification of habitat. The resulting sediment runoff can also restrict tidal flows and tidal elevations, resulting in losses of important fauna and flora (e.g., submerged aquatic vegetation).

Due to the intermittent nature of rainfall and runoff, the large variety of pollutant source types, and the variable nature of source loadings, urban runoff is difficult to control (Safavi 1996). The National Water Quality Inventory (EPA 2002) reports that runoff from urban areas is the leading source of impairment to surveyed estuaries and the third largest source of impairment to surveyed lakes. These include construction sediments, oil from autos, bacteria from failing septic systems, road salts, and heavy metals. Urban areas have an insidious pollution potential that one-time events such as oil spills do not. Pollutant increases gradually result in gradual declines in habitat quality.

Storm drains are often built to move water quickly away from roads, resulting in increased water input to streams. This greater volume and velocity erodes streambanks, increasing sediment loads and often temperatures. In a simulation model comparing an urban watershed with a forested watershed, Corbett et al. (1997) demonstrated that urban runoff volume and sediment yield were 5.5 times greater than forest runoff.

Waterborne polyaromatic hydrocarbon (PAH) levels have also been found to be significantly higher in an urbanized watershed when compared to a non-urbanized watershed (Fulton et al. 1993). Petroleum-based contaminants (such as fuel, oil, and some hydraulic fluids) contain PAHs, which can cause acute toxicity to EFH species and their prey at high levels of exposure and can also cause chronic lethal, as well as acute and chronic sublethal toxicity (Neff 1985).

Failing septic systems are an outgrowth of urban development. EPA estimates that 10 to 25 percent of all individual septic systems are failing at any one time, introducing excrement, detergents, endocrine disruptors, and chlorine into the environment. Even treated wastewater from urban areas can alter the physiology of intertidal organisms (Moles, A. and N. Hale. in press). Sewage discharge is a major source of coastal pollution, contributing 41 percent, 16 percent, 41 percent, and 6 percent of the total pollutant load for nutrients, bacteria, oils, and toxic metals, respectively (Kennish 1998). Nutrients such as phosphorus concentrations, in particular, are indicative of urban stormwater runoff (Holler 1990). Sewage wastes may also contain significant amounts of organic matter that exert a biochemical oxygen demand (Kennish 1998). Organic contamination contained within urban runoff can also cause immuno suppression (Arkoosh et al. 2001) (NMFS Draft 1998).

## Recommended Conservation Measures

The recommended conservation measures for urban/suburban development are provided below. For additional measures, see Section 2.3, Recommended Conservation Measures for Road Building and Maintenance.

1. Implement BMPs (EPA 1993) for sediment control during construction and maintenance operations. These can include avoiding ground disturbing activities during the wet season; minimizing exposure time of disturbed lands; using erosion prevention and sediment control methods; minimizing the spatial extent of vegetation disturbance; maintaining buffers of vegetation around wetlands, streams, and drainage ways; and avoiding building activities in areas of steep slopes and areas prone to mass wasting events with highly erodible soils. Use methods such as sediment ponds, sediment traps, bioswales, or other facilities designed to slow water runoff and trap sediment and nutrients.
2. Avoid using hard engineering structures for shoreline stabilization and channelization when possible. Use bioengineering approaches (i.e., using vegetation approaches with principles of geomorphology, ecology, and hydrology) to protect shorelines and river banks. Naturally stable shorelines and river banks should not be altered (see Section 4.7).
3. Encourage comprehensive planning for watershed protection so as to avoid filling and building in floodplain areas affecting EFH. Development sites should be planned to minimize clearing and grading, cut-and-fill, and new impervious surfaces.
4. Where feasible, remove impervious surfaces such as abandoned parking lots and buildings from riparian and shoreline areas, and re-establish wetlands and native vegetation.
5. Protect and restore vegetated buffer zones of appropriate width along all streams, lakes, and wetlands that include or influence EFH.
6. Manage stormwater to duplicate the natural hydrologic cycle, maintaining natural infiltration, and runoff rates to the maximum extent practicable.
7. Where in-stream flows are insufficient to maintain water quality and quantity needed for EFH, establish conservation guidelines for water use permits and encourage the purchase or lease of water rights and the use of water to conserve or augment instream flows in accordance with state and federal water law.
8. Encourage municipalities to use the best available technologies in upgrading their wastewater systems to avoid combined sewer overflow problems and chlorinated sewage discharges into rivers, estuaries, and the ocean.
9. On-site disposal systems should be properly designed and installed. They should be located away from open waters, wetlands, and floodplains.

### **2.3 Road Building and Maintenance**

The building and maintenance of roads can affect aquatic habitats by increasing rates of natural processes such as debris slides or landslides and sedimentation, introducing exotic species, and degrading water quality and chemical contamination (e.g., petroleum-based contaminants; see Section 2.2). Paved and dirt roads introduce an impervious or semi-pervious surface into the landscape. This surface intercepts rain and creates runoff carrying soil, sand and other sediments, and oil-based materials quickly downslope. If roads are built near streams, wetlands, or other sensitive areas, these may be affected by the increased sedimentation that occurs both from maintenance and use and during storm and snowmelt events. Even carefully designed and constructed roads can become sources of sediment and pollutants if they are not properly maintained.

## Potential Adverse Impacts

The effects of roads on aquatic habitat can be profound and include (1) increased deposition of fine sediments, (2) changes in water temperature, (3) elimination or introduction of migration barriers such as culverts, (4) changes in streamflow, (5) introduction of non-native plant species, and (6) changes in channel configuration.

Poorly surfaced roads can substantially increase surface erosion, and the rate of erosion is primarily a function of storm intensity, surfacing material, road slope, and traffic levels. This surface erosion results in an increase in fine sediment deposition (Bilby et al. 1989, MacDonald et al. 2001, Ziegler et al. 2001). An increase of fine-sediment deposition in stream gravels has been linked to decreased fry emergence, decreased juvenile densities, loss of winter carrying capacity, and increased predation of fishes (Koski 1981). Increased fines can reduce benthic production or alter the composition of the benthic community. For example, embryo-to-emergent fry survival of incubating salmonids is negatively affected by increases in fine sediments in spawning gravels (Chapman 1988, Everest et al. 1987, Scrivener and Brownlee 1989, Weaver and Fraley 1993, Young et al. 1991).

Roads built adjacent to streams result in changes in water temperature and increased sunlight reaching the stream as riparian vegetation is removed and/or altered in composition. Beschta et al. (1987) and Hicks et al. (1991) document some of the negative effects of road construction on fish habitat, including elevation of stream temperatures beyond the range of preferred rearing, inhibition of upstream migrations, increased disease susceptibility, reduced metabolic efficiency, and shifts in species assemblages.

Roads can also degrade aquatic habitat through improperly placed culverts at road-stream crossings that reduce or eliminate fish passage (Belford and Gould 1989, Clancy and Reichmuth 1990, Evans and Johnston 1980, Furniss et al. 1991). In a large river basin in Washington, 13 percent of the historical coho habitat was lost due to improper culvert design and placement (Beechie et al. 1994). Road crossings also affect benthic communities of stream invertebrates. Roads have a negative effect on the biotic integrity of both terrestrial and aquatic ecosystems (Trombulak and Frissell 2000). Studies indicate that populations of non-insect invertebrates tend to increase the more they are away from a road (Luce and Crowe 2001).

Roads may be the first point of entry into a virgin landscape for non-native grass species that are seeded along road cuts or introduced from seeds transported by tires and shoes. Roads can serve as corridors for such species allowing plants to move further into the landscape (Greenberg et al. 1997, Lonsdale and Lane 1994). Some non-native plants may be able to move away from the roadside and into aquatic sites of suitable habitat, where they may out-compete native species and have significant biological and ecological effects on the structure and function of the ecosystem.

Roads have three primary effects on hydrologic processes. First, they intercept rainfall directly on the road surface, in road cutbanks, and as subsurface water moving down the hillslope. Second, they concentrate flow, either on the road surfaces or in adjacent ditches or channels. Last, they divert or reroute water from flowpaths that would otherwise be taken if the road were not present (Furniss et al. 1991).

Road drainage and transport of water and debris, especially during heavy rains and snow melt periods, are primary reasons why roads fail, often with major structural, ecological, economic, or other social consequences. The effects of roads on peak streamflow depend strongly on the size of the watershed and the density of roads. Two of the effects are (1) changes in flood flows (Wemple et al. 1996) but mainly in smaller basins and for smaller floods (Beschta et al. 2000), and (2) increases in channel erosion and mass wasting (Montgomery 1994, Madej 2001, Wemple et al. 2001). For example, capture and rerouting of

water can dewater one small stream and cause major channel adjustments in the stream receiving the additional water. In large watersheds with low road density, properly located and maintained roads may constitute a small proportion of the land surface and have relatively insignificant effects on peak flow.

Roads can lead to increased rates of natural processes such as debris or landslides and sedimentation when slopes are destabilized and surface erosion and soil mass movement increases. Erosion is most severe when poor construction practices are allowed, combined with inadequate attention to proper road drainage and maintenance practices. Mass movement risks increase when roads are constructed on high-hazard soils and overly steep slopes. In steep areas prone to landslides, rates of mass soil movements affected by roads include shallow debris slides, deep-seated slumps and earthflows, and debris flows. Accelerated erosion rates from roads because of debris slides range from 30 to 300 times the natural rate in forested areas, but vary with terrain in the Pacific Northwest (Sidle et al. 1985). The magnitude of road-related mass erosion varies by climate, geology, road age, construction practices, and storm history. Road-related mass failures result from various causes, including improper placement and construction of road fills and stream crossings; inadequate culvert sizes to pass water, sediment, and wood during floods; poor road siting; modification of surface or subsurface drainage by the road surface or prism; and diversion of water into unstable parts of the landscape (Burroughs et al. 1976, Clayton 1983, Hammond et al. 1988, Furniss et al. 1991, Larsen and Parks 1997).

### Recommended Conservation Measures

The recommended conservation measures for road building and maintenance include the following:

1. Avoid locating roads near fish-bearing streams. Roads should be sited to avoid sensitive areas such as streams, wetlands, and steep slopes.
2. Incorporate erosion control and stabilization measures into road construction plans to reduce erosion potential.
3. Build bridges when possible. If culverts are to be used, they should be sized, constructed, and maintained to match the gradient and width of the stream, so as to accommodate 100-year flood flows, but equally to provide for migratory passage of adult and juvenile fishes. Use guidelines provided in "Guidelines for Salmonid Passage at Stream Crossing," NMFS, Southwest Region, October 2001 (<http://swr.nmfs.noaa.gov/hcd/NMFSSCG.PDF>).
4. Locate stream crossings in stable stream reaches.
5. Design bridge abutments to minimize disturbances to streambanks and place abutments outside of the floodplain whenever possible.
6. Avoid road construction across alluvial floodplains, mass wastage areas, or braided stream bottom lands unless site-specific protection can be implemented to ensure protection of soils, water, and associated resources.
7. Avoid side-casting on native surfaces of road materials into streams year-round.
8. Use only native vegetation in stabilization plantings.
9. Maintenance practices should not cause existing problems to become worse.

## **3.0 RIVERINE ACTIVITIES**

### **3.1 Mining** (see Section 5.6 - Marine Mining)

Mining and mineral extraction activities take many forms, such as commercial dredging and recreational suction dredging, placer, area surface removal, and contour operations. Activities include exploration, site preparation, mining, milling, waste management, decommissioning or reclamation, and even mine abandonment (American Fisheries Society [AFS] 2000). Mining and its associated activities have the potential to cause environmental impacts from exploration through post-closure. These impacts may

include adverse effects to EFH. The operation of metal, coal, rock quarries, and gravel pit mining has caused varying degrees of environmental damage in urban, suburban, and rural areas. Some of the most severe damage, however, occurs in remote areas, where some of the most productive fish habitat is often located (Sengupta 1993). Regulations have been designed to control and manage these changes to the landscape to avoid and minimize impacts. These regulations are updated as new technologies are developed to improve mineral extraction, reclaim mined lands, and limit environmental impacts. However, while environmental regulations may avoid, limit, control, or offset many of these potential impacts, mining will, to some degree, always alter landscapes and environmental resources (National Research Council [NRC] 1999).

### **3.1.1 Mineral Mining**

#### **Potential Adverse Impacts**

Potential impacts from mining include (1) adverse modification of hydrologic conditions so as to cause erosion of desirable habitats, (2) removal of substrates that serve as habitat for fish and invertebrates, (3) conversion of habitats, (4) release of harmful or toxic materials, and (5) creation of harmful turbidity levels.

The effects of mineral mining on EFH depend on the type, extent, and location of the activities. Minerals are extracted using several methods. Surface mining involves suction dredging, hydraulic mining, panning, sluicing, strip mining, and open-pit mining (including heap leach mining). Underground mining uses tunnels or shafts to extract minerals by physical or chemical means. Surface mining probably has a greater potential to affect aquatic ecosystems, though specific effects will depend on the extraction and processing methods and the degree of disturbance (Spence et al. 1996). Surface mining has the potential to eliminate vegetation, permanently alter topography, permanently and drastically alter soil and subsurface geological structure, and disrupt surface and subsurface hydrologic regimes (AFS 2000). While mining may not be as geographically pervasive as other sediment-producing activities, surface mining typically increases sediment delivery much more per unit of disturbed area than other activities because of the level of disruption of soils, topography, and vegetation. Erosion from surface mining and spoils may be one of the greatest threats to salmonid habitats in the western United States (Nelson et al. 1991).

Mining and placement of spoils in riparian areas can cause the loss of riparian vegetation and changes in heat exchange, leading to higher summer temperatures and lower winter stream temperatures (Spence et al. 1996). Bank instability can also lead to altered width-to-depth ratios, which further influence temperature (Spence et al. 1996). Mining efforts can also bury productive habitats near mine sites.

Mining operations can release harmful or toxic materials and their byproducts, either in association with actual mining, or in connection with machinery and materials used for mining. Mining can also introduce levels of heavy metals and arsenic that are naturally found within the stream bed sediments. Tailings and discharge waters from settling ponds can result in loss of EFH and life stages of managed species. The impact degrades water quality and levels can become high enough to prove lethal (North Pacific Fishery Management Council [Council] 1999).

Commercial operations may also involve road building (see Section 2.3), tailings disposal (Section 4.2), and leaching of extraction chemicals, all of which may create serious impacts to EFH. Cyanide, sulfuric acid, arsenic, mercury, heavy metals, and reagents associated with such development are a threat to EFH. Improper or in-water disposal of tailings may be toxic to managed species or their prey downstream. Upland disposal of tailings in unstable or landslide prone areas can cause large quantities of toxic compounds to be released into streams or to contaminate groundwater (Council 1999). Indirectly, the



sodium cyanide solution used in heap leach mining is contained in settling ponds from which groundwater and surface waters may become contaminated (Nelson et al. 1991).

Water pollution by heavy metals and acid is also often associated with mineral mining operations, as ores rich in sulfides are commonly mined for gold, silver, copper, iron, zinc, and lead. When stormwater comes in contact with sulfide ores, sulfuric acid is commonly produced (West et al. 1995). Abandoned pit mines can also cause severe water pollution problems.

Recreational gold mining with such equipment as pans, motorized or nonmotorized sluice boxes, concentrators, rockerboxes, and dredges can adversely affect EFH on a local level. Commercial mining is likely to involve activities at a larger scale with much disturbance and movement of the channel involved (Oregon Water Resources Research Institute [OWRRI] 1995).

### Recommended Conservation Measures

The recommended conservation measures for mining material are provided below. These measures are adapted from recommendations in Spence et al. (1996), NMFS (1996), and Washington Department of Fish and Wildlife (WDFW) (1998).

1. Avoid mineral mining in waters and streams containing EFH.
2. Schedule necessary in-water activities when the fewest species/least vulnerable life stages of federally managed species will be present.
3. Use an integrated environmental assessment, management, and monitoring package in accordance with state and federal law. Allow for adaptive operations to minimize adverse effects on EFH.
4. Avoid spills of dirt, fuel, oil, toxic materials, and other contaminants into EFH. Prepare a spill prevention plan and maintain appropriate spill containment and water repellent/oil absorbent cleanup materials on hand.
5. Treat wastewater (acid neutralization, sulfide precipitation, reverse osmosis, electrochemical, or biological treatments) and recycle on site to minimize discharge to streams. Test wastewater before discharge for compliance with federal and state clean water standards.
6. Minimize opportunities for sediments to enter or affect EFH. Use methods such as contouring, mulching, and construction of settling ponds to control sediment transport. Monitor turbidity during operations, and cease operations if turbidity exceeds predetermined threshold levels. Use turbidity/sediment curtains to limit the spread of suspended sediments and minimize the area affected.
7. Reclaim, rather than bury, mine waste that contains heavy metals, acid materials, or other toxic compounds if leachate can enter EFH through groundwater.
8. Restore natural contours and plant native vegetation on site after use to restore habitat function to the extent practicable. Monitor the site for an appropriate period of time to evaluate performance and implement corrective measures if necessary.
9. Minimize the aerial extent of ground disturbance (e.g., through phasing of operations), and stabilize disturbed lands to reduce erosion.

### **3.1.2 Sand and Gravel Mining**

#### Potential Adverse Impacts

Mining of sand and gravel is extensive and occurs by several methods. These include wet-pit mining (i.e., remove material from below the water table), dry-pit mining on beaches, exposed bars and ephemeral streambeds, and subtidal mining. Sand and gravel mining in riverine, estuarine, and coastal environments can create EFH impacts, including (1) turbidity plumes and resuspension effects, (2) removal of spawning habitat, and (3) alteration of channel morphology.

Mechanical disturbance of EFH spawning habitat by mining equipment can also lead to high mortality rates in early life stages. One result is the creation of turbidity plumes (Section 4.1), which can move spawning habitat several kilometers downstream. Sand and gravel mining in riverine, estuarine, and coastal environments can also suspend materials at the sites (Section 5).

Sedimentation may be a delayed effect because gravel removal typically occurs at low flow when the stream has the least capacity to transport fine sediments out of the system. Another delayed sedimentation effect results when freshets inundate extraction areas that are less stable than before. In addition, for species such as salmon, gravel operations can also interfere with migration past the site if they create physical or thermal changes at the work site or downstream from the site (OWRRI 1995).

Additionally, extraction of sand and gravel in riverine ecosystems can directly eliminate the amount of gravel available for spawning if the extraction rate exceeds the deposition rate of new gravel in the system. Gravel excavation also locally reduces the supply of gravel to downstream habitats. The extent of suitable spawning habitat may be reduced where degradation reduces gravel depth or exposes bedrock (Spence et al. 1996).

Mining can also alter channel morphology by making the stream channel wider and shallower. Consequently, the suitability of stream reaches as rearing EFH may be decreased, especially during summer low-flow periods when deeper waters are important for survival. Similarly, a reduction in pool frequency may adversely affect migrating adults that require holding pools (Spence et al. 1996). Changes in the frequency and extent of bedload movement and increased erosion and turbidity can also remove spawning substrates, scour redds (resulting in a direct loss of eggs and young), or reduce their quality by deposition of increased amounts of fine sediments. Other effects that may result from sand and gravel mining include increased temperatures (from reduction in summer base flows and decreases in riparian vegetation), decreased nutrients (from loss of floodplain connection and riparian vegetation), and decreased food production (loss of invertebrates) (Spence et al. 1996).

Examples of using gravel removal to improve habitat and water quality are limited and isolated (OWRRI 1995). Deep pools created by material removal in streams appears to attract migrating adult salmon for holding. These concentrations of fish may result in high losses as a result of increase predation or recreational fishing pressure.

### Recommended Conservation Measures

The recommended conservation measures for sand and gravel mining are provided below. These measures are adapted from NMFS (1996) and OWRRI (1995).

1. Avoid sand/gravel mining in waters containing EFH. Many factors influence site selection for a gravel or sand mining site. Because of the need to incorporate technical, economic, and environmental factors, siting decisions should be considered on a case-by-case basis (U.S. Fish and Wildlife Service [USFWS] 1980).
2. Identify upland or off-channel (where channel will not be captured) gravel extraction sites as alternatives to gravel mining in or adjacent to EFH, if possible.
3. Design, manage, and monitor sand and gravel mining operations to minimize potential direct and indirect impacts to EFH if operations in EFH cannot be avoided. This includes, but is not limited to, migratory corridors, foraging and spawning areas, stream/river banks, intertidal areas, etc.
4. Minimize the areal extent and depth of extraction.
5. Include restoration, mitigation, and monitoring plans in sand/gravel extraction plans.

## 3.2 Debris Removal

### 3.2.1 Organic Debris

Natural occurring flotsam such as LWD and macrophyte wrack (i.e., kelp) is often removed from streams, estuaries, and coastal shores. This debris is removed for a variety of reasons including dam operations, aesthetic concerns, and commercial and recreational uses. Because the debris affects habitat function and provides habitat for aquatic and terrestrial organisms, removing it may change the ecological balance among riverine, estuarine, and coastal ecosystems.

#### Potential Adverse Impacts

LWD and macrophyte wrack promote habitat complexity and structure to various aquatic and shoreline habitats. The structure provides cover for managed species, creates habitats and microhabitats (e.g., pools, riffles, undercut banks, side channels), and retains gravels and can maintain the underlying channel structure in riverine systems (Abbe and Montgomery 1996, Montgomery et al. 1995, Ralph et al. 1994, Spence et al. 1996). The structure removal reduces these habitat functions. Reductions in LWD input to estuaries have reduced the spatially complex and diverse channel systems that provide for productive salmon habitat (NRC 1996). Woody debris also plays a significant role in salt marsh ecology (Maser and Sedell 1994). Reductions in woody debris input to the estuaries may affect the ecological balance of the estuary. LWD also plays a significant role in benthic ocean ecology, where deep-sea wood borers convert the wood to fecal matter, providing terrestrial based carbon to the ocean food chain (Maser and Sedell 1994). Dams and commercial in-river harvest of large woody debris have dwindled the supply of wood, jeopardizing the ecological link between the forest and the sea (Collins et al. 2002, Collins et al. 2003, Maser and Sedell 1994).

Species richness, abundance, and biomass of macrofauna (e.g., sand crabs, isopods, amphipods, and polychaetes) associated with beach wrack are higher compared to beach areas with lower amounts of wrack or that are groomed (Dugan et al. 2000). The input and maintenance of wrack can strongly influence the structure of macrofauna communities, including the abundance of sand crabs (*Emerita analoga*) (Dugan et al. 2000), an important prey species to some EFH-managed species. Beach grooming can substantially alter the macrofaunal community structure of exposed sand beaches (Dugan et al. 2000). In addition, there are concerns that beach grooming efforts to remove wrack may also harm the eggs of the grunion (*Leuresthes tenuis*), an important prey item of EFH-managed species.

#### Recommended Conservation Measures

The recommended conservation measures for organic debris include the following:

1. Remove woody debris only when it presents a threat to life or property. Leave LWD wherever possible. Reposition, rather than remove, woody debris that must be moved.
2. Encourage appropriate federal, state, and local agencies to prohibit or minimize commercial removal of woody debris from rivers, estuaries, and beaches.
3. Encourage appropriate federal, state, and local agencies to aid in the downstream movement of LWD around dams, rather than removing it from the system.
4. Educate landowners and recreationalists about the benefits of maintaining LWD.
5. Localize beach grooming practices and minimize them whenever possible.
6. Conduct beach grooming only above the semilunar high tide as soon as the grunion spawning period begins in the spring, and continue 2 weeks after the last grunion spawning runs are observed in the summer.
7. Familiarize beach maintenance staff with the importance of such practices.

### 3.2.2 Inorganic Debris

Marine debris is a problem along much of U.S. coastal waters, littering shorelines, fouling estuaries, and creating hazards in the open ocean. Marine debris consists of a huge variety of man-made materials, such as general litter, dredged materials, hazardous wastes, and discarded or lost fishing gear. It enters waterways either indirectly through rivers and storm drains or by direct ocean dumping. Marine debris can have serious negative effects on EFH. Although several legislative laws and regulatory programs exist to prevent or control the problem, marine debris continues to severely affect our waters.

Congress has passed numerous legislative acts intended to prevent the disposal of marine debris in U.S. ocean waters. These include the Marine Protection, Research, and Sanctuaries Act, Titles I and II (also known as the Ocean Dumping Act), The Federal Water Pollution Control Act (Clean Water Act), and the Comprehensive Environmental Response, Compensation, and Liability Act (CERCLA). The International Convention for the Prevention of Pollution from Ships, commonly known as MARPOL Annex V (33 CFR 151), is intended to protect the marine environment from various types of garbage by preventing ocean dumping if the ship is less than 25 nautical miles (nm) from shore. Dumping of unground food waste and other garbage is prohibited within 12 nm from shore, and ground non-plastic or food waste may not be dumped within 3 nm of shore. The Ocean Dumping Act implements the Convention on the Prevention of Marine Pollution by Dumping of Wastes and Other Matter (London Dumping Convention) for the United States. Section 311 of the Federal Water Pollution Control Act makes it unlawful for any person to discharge any pollutant into the waters of the United States except as authorized by law. CERCLA stipulates that releases of hazardous substances in reportable quantities must be reported, and the release must be removed by the responsible party. Regulations implementing these acts are intended to control marine debris from ocean sources, including galley waste and other trash from ships, recreational boaters and fishermen, and offshore oil and gas exploration and facilities.

Land-based sources of marine debris account for about 80 percent of the marine debris on beaches and in our waters. Debris from these sources can originate from combined sewer overflows and storm drains, storm-water runoff, landfills, solid waste disposal, poorly maintained garbage bins, floating structures, and general littering of beaches, rivers, and open waters. Typical debris from these land-based sources includes raw or partially treated sewage, litter, hazardous materials, and discarded trash. Legislation and programs that address these land-based sources of pollution include the BEACH Act, the National Marine Debris Monitoring Program (NMDMP), the Shore Protection Act of 1989, and the CWA. The BEACH Act authorizes the EPA to fund state, territorial, Tribal, and local government programs that test and monitor coastal recreational waters near public access sites for microbial contaminants and to assess and monitor floatable debris. The NMDMP is a 5-year study designed to provide statistically valid estimates of marine debris affecting the entire U.S. coastline and to determine the main sources of the debris. The Shore Protection Act contains provisions to ensure that municipal and commercial solid wastes are not deposited in coastal waters during vessel transport from source to the waste receiving station. The CWA requires the EPA to develop and enforce regulations that treat storm water and combined sewer overflows as point source discharges requiring National Pollution Discharge Elimination System (NPDES) permits that prohibit non-storm water discharges into storm sewers.

#### Potential Adverse Impacts

Land- and ocean-based marine debris is a very diverse problem and adverse effects to EFH are likewise diverse. Floating or suspended trash can directly affect fish who consume or are entangled in the debris. Toxic substances in plastics can kill or impair fish and invertebrates that use habitat polluted by these materials which persist in the environment and can bioaccumulate through the food web. Once floatable debris settles to the bottom of estuaries, coastal, and open ocean areas it may continue to cause

environmental problems. Plastics and other materials with a large surface area can cover and suffocate immobile animals and plants, creating large spaces devoid of life. Currents can carry suspended debris to underwater reef habitats where the debris can become snagged, damaging these sensitive habitats. The typical floatable debris from combined sewer overflows includes street litter, sewage containing viral and bacterial pathogens, pharmaceutical by-products from human excretion, and pet wastes. It may contain condoms, tampons, and contaminated hypodermic syringes, all of which can pose physical and biological threats to EFH. Suspended organic matter has a high biological oxygen demand, and its reduction can cause algal blooms and anoxia that are detrimental to productive marine habitats. Pathogens can also contaminate shellfish beds.

### Recommended Conservation Measures

The recommended conservation measures for inorganic debris include the following:

1. Encourage proper trash disposal in coastal and ocean settings.
2. Advocate and participate in coastal cleanup activities.
3. Encourage enforcement of regulations addressing marine debris pollution and proper disposal.
4. Provide resources and technical guidance for development of studies and solutions addressing the problem of marine debris.
5. Provide resources to the public on the impact of marine debris and guidance on how to reduce or eliminate the problem.

### **3.3 Dam Operation**

The construction and operation of dams provide a source of hydropower, a reservoir for water storage, and a means to control flood control. Their operation, however, can affect water quality and quantity in riverine systems.

#### Potential Adverse Impacts

The effects of dam construction and operation on EFH can include (1) migratory impediments, (2) water flow and current pattern shifts, (3) thermal impacts, and (4) limits on sediment and woody debris transport.

One of the major impacts from dam construction and operation is that it impedes or completely creates impassable barriers to anadromous fish migrations in streams and rivers. Unless proper fish passage devices are in place, dams can either prevent access to productive upstream spawning habitat upstream or can alter downstream juvenile movements. The passage of salmon through turbines, sluiceways, bypass systems, and fish ladders also affects the quality of EFH (Pacific Fishery Management Council [PFMC] 1999).

In addition, dam operations also reduce downstream water velocities and change current patterns (PFMC 1999). These modifications can increase migration times (Raymond 1979). Water-level fluctuations, altered seasonal and daily flow regimes, reduced water velocities, and discharge volumes can affect the migratory behavior of juvenile salmonids and reduce the availability of shelter and foraging habitat (PFMC 1999).

Dams can also affect the thermal regimes of streams by raising water temperatures. Changes in water temperature can affect the development and smoltification of salmonids (PFMC 1999) and adult migration (Spence et al. 1996).

Dams also limit or alter natural sediment and LWD transport processes by impeding the high flows needed to scour fine sediments and move woody debris downstream (PFMC 1999). Curtailing these resources will affect the availability of spawning gravels and change channel morphology (Spence et al. 1996).

#### Recommended Conservation Measures (Adapted from PFMC 1999)

The recommended conservation measures for dam operation include the following:

1. Operate facilities to create flow conditions that provide for passage, water quality, proper timing of life history stages, and properly functioning channel conditions, and to avoid strandings and redd dewatering.
2. Develop water and energy conservation guidelines for integration into dam operation plans and into regional and watershed-based water resource plans.
3. Provide mitigation (including monitoring and evaluation) for nonavoidable adverse effects on EFH.

### **3.4 Commercial and Domestic Water Use**

Commercial and domestic water use demands to support the needs of homes, farms, and industries require a constant supply of water. Freshwater is diverted directly from lakes, streams, and rivers by means of pumping facilities, or is stored in impoundments. Because human populations are expected to continue increasing along most of the west coast, it is reasonable to assume that water uses, including water impoundments and diversion, will similarly increase (Gregory and Bisson 1997).

#### Potential Adverse Impacts

The information in this section is adapted from NMFS 1998.

The withdrawal of water can affect EFH by (1) altering natural flows and the process associated with flow rates, (2) affecting shoreline riparian habitats, (3) affecting prey bases, (4) affecting water quality, and (5) entrapping fishes. Water diversions can involve either withdrawals (reducing flow) or discharges (increasing flow). Water withdrawal will alter natural flow and stream velocity and channel depth and width. It can also change sediment and nutrient transport characteristics (Christie et al. 1993, Fajen and Layzer 1993), increase deposition of sediments, reduce depth, and accentuate diel temperature patterns (Zale et al. 1993). Loss of vegetation along stream banks and coastlines due to fluctuating water levels can decrease the availability of fish cover and reduce stability (Christie et al. 1993). Changes in the quantity and timing of stream flow alters the velocity of streams, which, in turn, affects the composition and abundance of both insect and fish populations (Spence et al. 1996). Returning irrigation water to a stream, lake, or estuary can substantially alter and degrade habitat (NRC 1989). Problems associated with return flows include increased water temperature, increased salinity, introduction of pathogens, decreased dissolved oxygen, increased toxic contaminants from pesticides and fertilizers, and increased sedimentation (Northwest Power Planning Council [NPPC] 1986). Diversions can also physically divert or entrap EFH-managed species (see Section 5.3).

#### Recommended Conservation Measures

The recommended conservation measures for commercial and domestic water use include the following:

1. Design projects to create flow conditions adequate to provide for passage, water quality, proper timing of life history stages, and avoidance of juvenile stranding and redd dewatering, as well as to maintain and restore properly functioning channel, floodplain, riparian, and estuarine conditions.

2. Establish adequate instream flow conditions for anadromous fish.
3. Screen water diversions on fish-bearing streams, as needed.
4. Incorporate juvenile and adult fish passage facilities on all water diversion projects (e.g., fish bypass systems).
5. Ensure that mitigation is provided for nonavoidable impacts.

#### **4.0 ESTUARINE ACTIVITIES**

##### **4.1 Dredging**

Dredging navigable waters is a continuous impact primarily affecting benthic and water-column habitats in the course of constructing and operating marinas, harbors, and ports. Routine dredging (i.e., the excavation of soft bottom substrates) is used to create deepwater navigable channels or to maintain existing channels that periodically fill with sediments. In addition, port expansion has become an almost continuous process due to economic growth, competition between ports, and significant increases in vessel size (see Section 4.3). Elimination or degradation of aquatic and upland habitats is commonplace because port expansion almost always affects open water, submerged bottoms, and, possibly, riparian zones.

##### Potential Adverse Impacts

The environmental effects of dredging on EFH can include (1) direct removal/burial of organisms; (2) turbidity/siltation effects, including light attenuation from turbidity; (3) contaminant release and uptake, including nutrients, metals, and organics; (4) release of oxygen consuming substances; (5) entrainment; (6) noise disturbances; and (6) alteration to hydrodynamic regimes and physical habitat.

Many EFH species forage on infaunal and bottom-dwelling organisms. Dredging may adversely affect these prey species at the site by directly removing or burying immobile invertebrates such as polychaete worms, crustacean, and other EFH prey types (Newell et al. 1998, Van der Veer et al. 1985). Similarly, the dredging activity may also force mobile animals such as fish to migrate out of the project area. Recolonization studies suggest that recovery may not be quite as straightforward. Physical factors, including particle size distribution, currents, and compaction/stabilization processes following deposition reportedly can regulate recovery after dredging events. Rates of recovery listed in the literature range from several months for estuarine muds to up to 2 to 3 years for sands and gravels. Recolonization can also take up to 1 to 3 years in areas of strong current but up to 5 to 10 years in areas of low current. Thus, forage resources for benthic feeders may be substantially reduced.

The use of certain types of dredging equipment can result in greatly elevated levels of fine-grained mineral particles or suspended sediment concentration (SSC), usually smaller than silt, and organic particles in the water column. The associated turbidity plumes of suspended particulates may reduce light penetration and lower the rate of photosynthesis for subaquatic vegetation (Dennison 1987) and the primary productivity of an aquatic area if suspended for extended periods of times (Cloern 1987). If suspended sediments loads remain high, fish may suffer reduced feeding ability (Benfield and Minello 1996) and be prone to fish gill injury (Nightingale and Simenstad 2001a).

Sensitive habitats such as submerged aquatic vegetation beds, which provide food and shelter, may also be damaged. Eelgrass beds are critical to nearshore food web dynamics (Wyllie-Echeverria and Phillips 1994, Murphy et al. 2000). Studies have shown seagrass beds to be among the areas of highest primary productivity in the world (Herke and Rogers 1993, Hoss and Thayer 1993). This primary production, combined with other nutrients, provide high rates of secondary production in the form of fish (Herke and Rogers 1993, Good 1987, Sogard and Able 1991).

The contents of the suspended material may react with the dissolved oxygen in the water and result in short-term oxygen depletion to aquatic resources (Nightingale and Simenstad 2001a). Dredging can also disturb aquatic habitats by resuspending bottom sediments and, thereby, recirculate toxic metals (e.g., lead, zinc, mercury, cadmium, copper etc.), hydrocarbons (e.g., polyaromatics), hydrophobic organics (e.g., dioxins), pesticides, pathogens, and nutrients into the water column (EPA 2000). Toxic metals and organics, pathogens, and viruses, absorbed or adsorbed to fine-grained particulates in the material, may become biologically available to organisms either in the water column or through food chain processes.

Direct uptake of fish species by hydraulic dredging at the proposed borrow site is also an issue. Definitive information in the literature shows that elicit avoidance responses to the suction dredge entrainment occurs for both benthic and water column oriented species (Larson and Moehl 1998, McGraw and Armstrong 1990).

Dredging, as well as the equipment used in the process such as pipelines (see Section 4.10), may damage or destroy spawning, nursery, and other sensitive habitats such as emergent marshes and subaquatic vegetation, including eelgrass beds and kelp beds. Dredging may also modify current patterns and water circulation of the habitat by changing the direction or velocity of water flow, water circulation, or dimensions of the water body traditionally used by fish for food, shelter, or reproductive purposes.

#### Recommended Conservation Measures

The recommended conservation measures for dredging include the following:

1. Avoid new dredging to the maximum extent practicable. Activities that would likely require dredging (such as placement of piers, docks, marinas, etc.) should instead be sited in deep water areas or designed to alleviate the need for maintenance dredging. Projects should be permitted only for water dependent purposes and only when no feasible alternatives are available.
2. Incorporate adequate control measures to minimize turbidity where the dredging equipment used is expected to create significant turbidity.
3. Undertake multi-season, pre-, and post-dredging biological surveys to assess impacts to animal and submerged aquatic vegetation communities.
4. Provide appropriate compensation for significant impacts (short-term, long-term, and cumulative) to benthic environments resulting from dredging.
5. Perform dredging during the time frame when impacts due to entrainment of EFH-managed species or their prey are least likely to be entrained. Dredging should be avoided in areas with submerged aquatic vegetation.
6. Reference all dredging latitude-longitude coordinates at the site so that information can be incorporated into a geographical information system (GIS) format. Inclusion of aerial photos may be useful to identify precise locations for long-term evaluation.
7. Test sediments for contaminants as per EPA and USACE requirements.
8. Address cumulative impacts of past and current dredging operations on EFH by considering them as part of the permitting process.
9. Identify excess sedimentation in the watershed that prompts excessive maintenance dredging activities and implement appropriate management agencies to ensure that actions are taken to curtail those causes.
10. Ensure that bankward slopes of the dredged area are slanted to acceptable side slopes (e.g., 3:1) to ensure that sloughing does not occur.
11. Avoid placing pipelines and accessory equipment used in conjunction with dredging operations to the maximum extent possible close to kelp beds, eelgrass beds, estuarine/salt marshes, and other high value habitat areas.



## **4.2 Disposal/Landfills**

The discharge of dredged materials subsequent to dredging operations or the use of fill material in the construction/development of harbors results in sediments (e.g., dirt, sand, mud) covering or smothering existing submerged substrates. Usually these covered sediments are of a soft-bottom nature as opposed to rock or hard-bottom substrates.

### **4.2.1 Disposal of Dredged Material**

#### Potential Adverse Impacts

The disposal of dredged material can adversely affect EFH by (1) affecting or destroying benthic communities, (2) affecting adjacent habitats, and (3) creating turbidity plumes and introducing contaminants and/or nutrients.

Disposing dredged materials result in varying degrees of change in the physical, chemical, and biological characteristics of the substrate. Discharges may adversely affect infaunal and bottom-dwelling organisms at the site by smothering immobile organisms (e.g., prey invertebrate species) or forcing mobile animals (e.g., benthic-oriented fish species) to migrate from the area. Infaunal invertebrate plants and animals present prior to a discharge are unlikely to recolonize if the composition of the discharged material is drastically different.

Erosion, slumping, or lateral displacement of surrounding bottom of such deposits can also adversely affect substrate outside the perimeter of the disposal site by changing or destroying benthic habitat. The bulk and composition of the discharged material and the location, method, and timing of discharges may all influence the degree of impact on the substrate.

The discharge of material can result in greatly elevated levels of fine-grained mineral particles, usually smaller than silt, and organic particles in the water column (i.e., turbidity plumes). These suspended particulates may reduce light penetration and lower the rate of photosynthesis and the primary productivity of an aquatic area if suspended for lengthy intervals. Aquatic vegetation such as eelgrass beds and kelp beds may also be affected. Managed fish species may suffer reduced feeding ability, leading to limited growth and lowered resistance to disease if high levels of suspended particulates persist. The contents of the suspended material may react with the dissolved oxygen in the water and result in oxygen depletion. Toxic metals and organics, pathogens, and viruses absorbed or adsorbed to fine-grained particulates in the material may become biologically available to organisms either in the water column or through food chain processes.

The discharge of dredged or fill material can change the chemistry and the physical characteristics of the receiving water at the disposal site by introducing chemical constituents in suspended or dissolved form. Reduced clarity and excessive contaminants can reduce, change, or eliminate the suitability of water bodies for populations of groundfish, other fish species, and their prey. The introduction of nutrients or organic material to the water column as a result of the discharge can lead to a high BOD, which in turn can lead to reduced dissolved oxygen, thereby potentially affecting the survival of many aquatic organisms. Increases in nutrients can favor one group of organisms such as polychaetes or algae to the detriment of other types.

#### **4.2.2 Fill Material**

##### Potential Adverse Impacts

Adverse impacts to EFH from the introduction of fill material included (1) loss of habitat function and (2) changes in hydrologic patterns.

Aquatic habitats sustain remarkably high levels of productivity and support various life stages of fish species and their prey. Many times these habitats are used for multiple purposes, including habitat necessary for spawning, breeding, feeding, or growth to maturity. The introduction of fill material eliminates those functions and permanently removes the habitat from production.

The discharge of dredged or fill material can modify current patterns and water circulation by obstructing flow, changing the direction or velocity of water flow and circulation, or otherwise changing the dimensions of a water body. As a result, adverse changes can occur in the location, structure, and dynamics of aquatic communities; shoreline and substrate erosion and deposition rates; the deposition of suspended particulates; the rate and extent of mixing of dissolved and suspended components of the water body; and water stratification (NMFS 1998).

##### Recommended Conservation Measures

The recommended conservation measures for fill material include the following:

1. Study all options for disposal of dredged materials, including disposal sites and methods used. Upland dredge disposal sites should be considered as an alternative to offshore disposal sites.
2. The cumulative impacts of past and current fill operations on EFH should be addressed by federal, state, and local resource management and permitting agencies and considered in the permitting process.
3. Disposal of dredge material in EFH should meet or exceed applicable state and/or federal quality standards for such disposal.
4. State and federal agencies should identify the direct and indirect impacts open-water disposal permits for dredged material may have on EFH during proposed project reviews. Benthic productivity should be determined by sampling prior to any discharge of fill material. Sampling design should be developed with input from state and federal natural resource agencies.
5. The areal extent of any disposal site in EFH should be avoided or minimized. In some cases, however, thin layer disposal may be less deleterious. All non-avoidable adverse impacts should be mitigated.
6. All spoil disposal permits should reference latitude-longitude coordinates of the site so information can be incorporated into GIS systems. Inclusion of aerial photos or benthic photos may also be required to identify precise locations and determine long-term effects.
7. Fills in estuaries and bays for development of commercial enterprises should be avoided.
8. Identify and characterize EFH habitat functions/services in the project areas.
9. Adequate compensatory mitigation should be provided for unavoidable impacts.

#### **4.3 Vessel Operations/Transportation/Navigation**

The demand by port districts to increase infrastructure capacity to accommodate additional vessel operations for cargo handling activities and marine transportation is predicted to continue. Population growth and demands for international business trade along the Pacific Rim exert pressure to expand coastal towns and port facilities, resulting in net estuary losses (Kagan 1991, Fawcett and Marcus 1991). Port expansion has become an almost continuous process due to economic growth, competition between

ports, and significant increases in vessel size (Council 1999). In addition, with increased population growth comes the steady demand for providing new and expanded water transit services. Finally, providing additional recreational opportunities by constructing and enlarging recreational marinas is also foreseen.

### Potential Adverse Impacts

The expansion of port facilities, vessel/ferry operations, and recreational marinas can bring additional impacts to EFH. Additional land needed to improve shipping efficiency can only be accommodated by changing land-use operations or adding new land by filling aquatic habitats. New wharves and piers decrease photic penetration in the water and decreases primary production (see Section 4.6). More hard surface increases nonpoint surface discharges (see Section 2.2), adds debris sources, and reduces buffers between land use and the aquatic ecosystem. These will include direct, indirect, and cumulative impacts on shallow subtidal, deep subtidal, eelgrass beds, mudflats, sand shoals, rock reefs, and salt marsh habitats. Such impacts would be site-specific. Some activities affecting these habitats, including new channel deepening and maintenance dredging (see Section 4.1), disposal of dredged material (see Section 4.2), reduced water quality from resuspension of contaminated sediments, ballast water discharge (see Section 4.4), and shading from overwater structures (see Section 4.6), have been addressed in other sections. Additional impacts include vessel groundings, modification of water circulation (breakwaters, channels, and fill), vessel wake generation, pier lighting, anchor scour and prop scour, and the discharge of contaminants and debris.

Potential adverse impacts to EFH can occur during both the construction and operation phases. Direct impacts include permanent or temporary loss of productive forage habitat resulting from new channel deepening and maintenance dredging (see Section 4.1), turbidity-related impacts due to both dredging and disposal of dredged material (see Section 4.2), and reduced water quality from resuspension of contaminated sediments (see Section 4.1). In addition, dredging in tidal wetland areas could result in the spread of nonnative invasive plant species (see Section 4.4).

An increase in the number and size of vessels can generate more wave and surge effects on shorelines. These vessel-wake, wash events can affect shorelines depending on the wake wave energy, the water depth, and the type of shoreline. Vessel wakes can cause a significant increase in shoreline erosion, affect wetland habitat, and increase water turbidity. Vessel prop wash can also damage aquatic vegetation and disturb sediments, which may increase turbidity and suspend contaminants (Klein 1997, Warrington 1999). Changes in prey communities under ferry terminals have been attributed, in part, to prop wash from ferries (Blanton et al. 2001, Haas et al. 2002).

Impacts can also occur from anchor scour. Mooring buoys, when anchored in shallow nearshore waters, can drag the anchor chain across the bottom, destroying submerged vegetation and creating a circular scour hole (Walker et al. 1989, *In* Shafer 2002). A study by Hastings et al. (1995) (*In* Shafer 2002) in Australia found that up to 18 percent of total seagrass cover was lost to mooring buoy scour.

Vessel discharges, engine operations, bottom paint sloughing, boat washdowns, painting, and other vessel maintenance activities can deliver debris, nutrients, and contaminants to waterways and may degrade water quality and contaminate sediments.

Inadequate flushing of marinas also results in water quality problems (USACE 1993, Klein 1997). Poor flushing in marinas in Puget Sound resulted in increases in temperature, increased phytoplankton populations with nocturnal dissolved oxygen level declines resulting in organism hypoxia, and pollutant inputs (Cardwell et al. 1980). An exchange of at least 30 percent of the water in the marina during a tidal change should minimize temperature increases and dissolved oxygen problems (Cardwell et al. 1980).

## Recommended Conservation Measures

The recommended conservation measures for vessel operations, transportation, and navigation include the following:

1. Locate marinas in areas of low biological abundance and diversity, for example, avoiding dense beds of eelgrass or other submerged aquatic vegetation including macroalgae.
2. Excavate uplands to create marina basins rather than converting intertidal or shallow subtidal to deeper subtidal for basin creation.
3. Avoid the disturbance of eelgrass beds, mudflats, and wetlands as part of the project design. In situations where such impacts are unavoidable, appropriate compensatory mitigation should be incorporated into the project with the approval of appropriate regulatory agencies. Specific habitat types such as eelgrass beds need to be mitigated in-kind. For other habitat types where in-kind mitigation is unavailable, the habitat values or functions of these threatened habitats should be calculated and appropriate mitigation be provided to ensure no net loss of habitat functions. This also includes the habitat value of traditional shoreline protection materials (e.g., revetment and breakwaters). Other dredging-related conservation measures are provided in Section 4.1.
4. Leave marine riparian buffers in place to enhance intertidal microclimate and nutrient input.
5. Adequate monitoring on the success of mitigation efforts should be included as part of the project and incorporated into a mitigation and monitoring plan.
6. Conduct preconstruction surveys by qualified biologists/botanists to identify and map areas of invasive plant species existing within potential project construction areas. Eradication of non-native species should be conducted well in advance of construction.
7. Include low-wake vessel technology, appropriate routes, and BMPs for wave attenuation structures as part of the design and permit process. Vessels should be operated at sufficiently low speeds to reduce wake energy, and no-wake zones should be designated near sensitive habitats.
8. Incorporate BMPs to prevent or minimize contamination from ship bilge waters, antifouling paints, shipboard accidents, shipyard work, maintenance dredging and disposal, and nonpoint source contaminants from upland facilities related to vessel operations and navigation.
9. Locate mooring buoys in water deep to avoid grounding and minimize affects of prop wash. Use subsurface floats or other methods to prevent contact of the anchor line with the substrate.
10. Collect and treat runoff from parking lots and other impervious surfaces to remove contaminants prior to delivery to any receiving waters
11. Locate facilities in areas with sufficient water velocities to dissipate fuels and pollutants from vessels and maintain temperature and dissolved oxygen levels within acceptable ranges.
12. Locate marinas where they do not interfere with drift sectors determining the structure and function of adjacent habitats.

### **4.4 Introduction of Exotic Species**

The introductions of exotic species into estuarine and marine habitats has been well documented (Rosecchi et al. 1993, Kohler and Courtenay 1986, Spence et al. 1996) and can be intentional (e.g., for the purpose of stock or pest control) or unintentional (e.g., fouling organisms). Exotic fish, shellfish, pathogens, and plants can enter the environment from industrial shipping (e.g., as ballast), recreational boating, aquaculture (see Section 4.11), biotechnology, and aquariums. The transportation of nonindigenous organisms to new environments can have many severe impacts on habitat (Omori et al. 1994).

## Potential Adverse Impacts

Long-term impacts of the introduction of nonindigenous and reared species can change the natural community structure and dynamics, lower the overall fitness and genetic diversity of natural stocks, and pass and/or introduce exotic lethal disease. Overall, exotic species introductions create five types of negative effects, including (1) habitat alteration, (2) trophic alteration, (3) gene pool alteration, (4) spatial alteration, and (5) introduction of diseases. Habitat alteration includes the excessive colonization of exotic species (e.g., *Spartina* grasses), which preclude the growth of endemic organisms (e.g., eelgrass). The introduction of exotic species may alter community structure by predation on native species or by population explosions of the introduced species. Spatial alteration occurs when territorial introduced species compete with and displace native species. Although hybridization is rare, it may occur between native and introduced species and can result in gene pool deterioration.

Non-native plants and algae can degrade coastal and marine habitats by changing natural habitat qualities. Introduced organisms increase competition with indigenous species or forage on indigenous species, which can reduce fish and shellfish populations. Long-term impacts from the introduction of nonindigenous and reared species can change the natural community structure and dynamics, lower the overall fitness and genetic diversity of natural stocks, and pass and/or introduce exotic lethal diseases. The introduction of exotic organisms also threatens native biodiversity and could lead to changes in relative abundances of species and individuals that are of ecological and economic importance.

The introduction of bacteria, viruses, and parasites is another severe threat to EFH as it may reduce habitat quality. New pathogens or higher concentrations of disease can be spread throughout the environment resulting in deleterious habitat conditions.

## Recommended Conservation Measures

The recommended conservation measures for the introduction of exotic species include the following:

1. Encourage vessels to perform a ballast water exchange in marine waters (in accordance with the U.S. Coast Guard's voluntary regulations) to minimize the possibility of introducing exotic estuarine species into similar habitats. Ballast water taken on in marine waters will contain fewer organisms and these will be less likely to become invasive in estuarine conditions than species transported from other estuaries.
2. Discourage vessels that have not performed a ballast water exchange from discharging their ballast water into estuarine receiving waters.
3. Require vessels brought from other areas over land via trailer to clean any surfaces that may harbor non-native plant or animal species (propellers, hulls, anchors, fenders, etc.). Bilges should be emptied and cleaned thoroughly using hot water or a mild bleach solution. These activities should be performed in an upland area to prevent introduction of non-native species during the cleaning process.
4. Prevent exotic species from aquaculture operations until a thorough scientific evaluation and risk assessment is performed (see Section 4.11).
5. Aquaculture facilities rearing non-native species should be located upland and use closed-water circulation systems whenever possible.
6. Treat effluent from public aquaria displays and laboratories and educational institutes using exotic species prior to discharge to prevent the introduction of viable animals, plants, reproductive material, pathogens, or parasites into the environment.

## **4.5 Pile Installation and Removal**

Pilings are an integral component of many overwater and in-water structures. They provide support for the decking of piers and docks, function as fenders and dolphins to protect structures, support navigation markers, and help in the construction of breakwaters and bulkheads. Materials used in pilings include steel, concrete, wood (both treated and untreated), plastic, or a combination thereof. Piles are usually driven into the substrate using either impact hammers or vibratory hammers. Impact hammers consist of a heavy weight that is repeatedly dropped onto the top of the pile, driving it into the substrate. Vibratory hammers use a combination of a stationary, heavy weight and vibration, in the plane perpendicular to the long axis of the pile, to force the pile into the substrate. The type of hammer used depends on a variety of factors, including pile material and substrate type. Impact hammers can be used to drive all types of piles, while vibratory hammers are generally most efficient at driving piles with a cutting edge (e.g., hollow steel pipe) and are less efficient at driving displacement piles (those without a cutting edge that must displace the substrate). Displacement piles include solid concrete, wood, and closed-end steel pipe. While impact hammers are able to drive piles into most substrates (including hardpan, glacial till, etc.), vibratory hammers are limited to softer, unconsolidated substrates (e.g., sand, mud, gravel). Because vibratory hammers do not use force to drive the piles, the bearing capacity is not known and the piles must often be proofed with an impact hammer. This involves striking the pile a number of times with the impact hammer to ensure that it meets the designed bearing capacity. Under certain circumstances, piles may be driven using a combination of vibratory and impact hammers. The vibratory hammer makes positioning and plumbing of the pile easier; therefore, it is often used to drive the pile through the soft, overlying material. Once the pile stops penetrating the sediment, the impact hammer is used to finish driving the pile to final depth. An additional advantage of this method is that the vibratory hammer can be used to extract and reposition the pile, while the impact hammer cannot.

Overwater structures must often meet seismic stability criteria, requiring that the supporting piles are attached to, or driven into, the underlying hard material. This requirement often means that at least some impact driving is necessary. Piles that do not need to be seismically stable, including temporary piles, fender piles, and some dolphin piles, may be driven with a vibratory hammer, providing the type of pile and sediments are appropriate.

Piles can be removed using a variety of methods, including vibratory hammer, direct pull, clam shell grab, or cutting/breaking the pile below the mudline. Vibratory hammers can be used to remove all types of pile, including wood, concrete, and steel. Old brittle piles, however, may break under the vibrations and necessitate another method. The direct pull method involves placing a choker around the pile and pulling upward with a crane or other equipment. Broken stubs are often removed with a clam shell and crane. In this method, the clam shell grips the pile near the mudline and pulls it out. In other instances, piles may be cut or broken below the mudline, leaving the buried section in place.

### **4.5.1 Pile Driving**

#### Potential Adverse Impacts

Pile driving can generate intense underwater sound pressure waves that may adversely affect the ecological functioning of EFH. These pressure waves have been shown to injure and kill fish (e.g., CalTrans 2001, Longmuir and Lively 2001, Stotz and Colby 2001, Stadler, pers. obs. 2002). Injuries associated directly with pile driving are poorly studied, but include rupture of the swimbladder and internal hemorrhaging (CalTrans 2001; Abbott and Bing-Sawyer 2002; Stadler, pers. obs. 2002). Sound pressure levels (SPL) 100 decibels (dB) above the threshold for hearing are thought to be sufficient to damage the auditory system in many fishes (Hastings 2002).

The type and intensity of the sounds produced during pile driving depend on a variety of factors, including, but not limited to, the type and size of the pile, the firmness of the substrate into which the pile is being driven, the depth of water, and the type and size of the pile-driving hammer. SPLs are positively correlated with the size of the pile, as more energy is required to drive larger piles. Wood and concrete piles appear to produce lower sound pressures than hollow steel piles of a similar size, although it is not yet clear if the sounds produced by wood or concrete piles are harmful to fishes. Hollow steel piles as small as 14 inches in diameter have been shown to produce SPLs that can injure fish (Reyff 2003). Firmer substrates require more energy to drive piles and produce more intense sound pressures. Sound attenuates more rapidly with distance from the source in shallow water than it does in deep water (Rogers and Cox 1988).

Driving hollow steel piles with impact hammers produces intense, sharp spikes of sound that can easily reach levels injurious to fish. Vibratory hammers, on the other hand, produce sounds of lower intensity, with a rapid repetition rate. A key difference between the sounds produced by impact hammers and those produced by vibratory hammers is the responses they evoke in fish. When exposed to sounds that are similar to those of a vibratory hammer, fish consistently displayed an avoidance response (Enger et al. 1993, Dolat 1997, Knudsen et al. 1997, Sand et al. 2000), and they did not habituate to the sound, even after repeated exposure (Dolat 1997, Knudsen et al. 1997). Fishes may respond to the first few strikes of an impact hammer with a startle response. After these initial strikes, the startle response wanes, and the fishes may remain within the field of a potentially harmful sound (Dolat 1997, NMFS 2001). The differential responses to these sounds are due to the differences in the duration and frequency of the sounds. When compared to impact hammers, the sounds produced by vibratory hammers are of longer duration (minutes versus msec) and have more energy in the lower frequencies (15 to 26 Hertz [Hz] versus 100 to 800 Hz) (Wüsig, et al. 2000, Carlson et al. 2001). Studies have shown that fish respond to particle acceleration of 0.01 meters per second squared ( $m/s^2$ ) at infrasound frequencies, that the response to infrasound is limited to the nearfield (less than 1 wavelength), and that the fish must be exposed to the sound for several seconds (Enger et al. 1993, Knudsen et al. 1994, Sand et al. 2000). Impact hammers, however, produce such short spikes of sound with little energy in the infrasound range, that fish fail to respond to the particle motion (Carlson et al. 2001). Thus, impact hammers may be more harmful than vibratory hammers because they produce more intense pressure waves and because the sounds produced do not elicit an avoidance response in fishes, which exposes them to those harmful pressures for longer periods.

The degree to which an individual fish exposed to sound will be affected depends on a number of variables, including (1) species of fish, (2) fish size, (3) presence of a swimbladder, (4) physical condition of the fish, (5) peak sound pressure and frequency, (6) shape of the sound wave (rise time), (7) depth of the water around the pile, (8) depth of the fish in the water column, (9) amount of air in the water, (10) size and number of waves on the water surface, (11) bottom substrate composition and texture, (12) effectiveness of bubble curtain sound/pressure attenuation technology, (13) tidal currents, and (14) presence of predators.

Depending on these factors, effects on fish can range from changes in behavior to immediate mortality. There are little data on the SPL required to injure fish. Short-term exposure to peak SPLs above 190 dB (re: 1  $\mu$ Pa) is thought to impose physical harm on fish (Hastings 2002). However, 155 dB (re: 1  $\mu$ Pa) may be sufficient to temporarily stun small fish (personal communication, J. Miner, Gunderboom, Inc., Anchorage, Alaska, 2002). Stunned fish, while perhaps not physically injured, are more susceptible to predation. Small fish are more prone to injury by intense sound than are larger fish of the same species (Yelverton et al. 1975). For example, a number of surfperches (*Cymatogaster aggregata* and *Embiotoca lateralis*) were killed during impact pile driving (Stadler, pers. obs. 2002). Most of the dead fish were the smaller *C. aggregata* and similar sized specimens of *E. lateralis*, even though many larger *E. lateralis* were in the same area. Dissections revealed that the swimbladder of the smallest fish (80 millimeter

[mm] forklength [FL]) was completely destroyed, while that of the largest individual (170 mm FL) was nearly intact, indicating a size-dependent effect. The SPLs that killed these fish are not yet known. Of the reported fish kills associated with pile driving, all have occurred during use of an impact hammer on hollow steel piles (Longmuir and Lively 2001, NMFS 2001, Stotz and Colby 2001, NMFS 2003).

Systems successfully designed to reduce the adverse effects of underwater SPLs on fish have included the use of air bubbles. Both confined (i.e., metal or fabric sleeve) and unconfined air bubble systems have been shown to attenuate underwater sound pressures up to 28 dB (Wursig et al. 2000, Longmuir and Lively 2001, Christopherson and Wilson 2002, Reyff and Donovan 2003). When using an unconfined air bubble system in areas of strong currents, it is critical that the pile be fully contained within the bubble curtain. To accomplish this when designing the system, adequate air flow and ring spacing, both vertically and in terms of distance from the pile, are factors that should be considered.

#### Recommended Conservation Measures

Install hollow steel piles with an impact hammer at a time of year when larval and juvenile stages of fish species with designated EFH are not present. If this is not possible, then the following measures regarding pile driving should be incorporated to minimize adverse effects:

1. Drive piles during low tide when located in intertidal and shallow subtidal areas.
2. Use a vibratory hammer when driving hollow steel piles. Under those conditions where impact hammers are required for reasons of seismic stability or substrate type, the pile should be driven as deep as possible with a vibratory hammer before using the impact hammer.
3. Monitor peak SPLs during pile driving to ensure that they do not exceed the 190 dB re:1  $\mu$ Pa threshold for injury to fish.
4. Implement measures to attenuate the sound should SPLs exceed the 180 dB re: 1  $\mu$ Pa threshold. If sound pressure levels exceed acceptable limits, implement mitigation measures. Methods to reduce the sound pressure levels include, but are not limited to, the following:
  - a) Surround the pile with an air bubble curtain system or air-filled coffer dam.
  - b) Because the sound produced has a direct relationship to the force used to drive the pile, use a smaller hammer to reduce the sound pressures.
  - c) Use a hydraulic hammer if impact driving cannot be avoided. The force of the hammer blow can be controlled with hydraulic hammers; reducing the impact force will reduce the intensity of the resulting sound.
5. Drive piles when the current is reduced (i.e., centered around slack current) in areas of strong current to minimize the number of fish exposed to adverse levels of underwater sound.

#### **4.5.2 Pile Removal**

##### Potential Adverse Impacts

The primary adverse effect of removing piles is the suspension of sediments, which may result in harmful levels of turbidity and release of contaminants contained in those sediments (see Section 4.1). Vibratory pile removal tends to cause the sediments to slough off at the mudline, resulting in relatively low levels of suspended sediments and contaminants. Vibratory removal of piles is gaining popularity because it can be used on all types of piles, providing that they are structurally sound. Breaking or cutting the pile below the mudline may suspend only small amounts of sediment, providing the stub is left in place and little digging is required to access the pile. Direct pull or use of a clamshell to remove broken piles may, however, suspend large amounts of sediment and contaminants. When the piling is pulled from the substrate using these two methods, sediments clinging to the piling will slough off as it is raised through



the water column, producing a potentially harmful plume of turbidity and/or contaminants. The use of a clamshell may suspend additional sediment if it penetrates the substrate while grabbing the piling.

While there is a potential to adversely affect EFH during the removal of piles, many of those removed are old creosote-treated timber piles. In some cases, the long-term benefits to EFH obtained by removing a consistent source of contamination may outweigh the temporary adverse effects of turbidity.

#### Recommended Conservation Measures

The recommended conservation measures for pile removal include the following:

1. Remove piles completely rather than cutting or breaking them off, if the piles are structurally sound.
2. Minimize the suspension of sediments and disturbance of the substrate when removing piles. Measures to help accomplish this include, but are not limited to, the following:
  - a) When practicable, remove piles with a vibratory hammer, rather than using the direct pull or clamshell method.
  - b) Remove the pile slowly to allow sediment to slough off at, or near, the mudline.
  - c) The operator should first hit or vibrate the pile to break the bond between the sediment and the pile to minimize the potential for the pile to break, as well as to reduce the amount of sediment sloughing off the pile during removal.
  - d) Place a ring of clean sand around the base of the pile. This ring will contain some of the sediment that would normally be suspended.
  - e) Encircle the pile, or piles, with a silt curtain that extends from the surface of the water to the substrate.
3. Complete each pass of the clamshell to minimize suspension of sediment if pile stubs are removed with a clamshell.
4. Fill all holes left by the piles with clean, native sediments if possible.
5. Place piles on a barge equipped with a basin to contain all attached sediment and runoff water after removal. Creosote-treated timber piles should be cut into short lengths to prevent reuse, and all debris, including attached, contaminated sediments, should be disposed of in an approved upland facility.
6. Using a pile driver, drive broken/cut stubs far enough below the mudline to prevent release of contaminants into the water column as an alternative to their removal.

#### **4.6 Overwater Structures**

Overwater structures include commercial and residential piers and docks, floating breakwaters, barges, rafts, booms, and mooring buoys. These structures typically are located in intertidal areas out to about 15 meters below the area exposed by the mean lower low tide (i.e., the shallow subtidal zone). Light, wave energy, substrate type, depth, and water quality are the primary factors controlling the plant and animal assemblages found at a particular site. Overwater structures and associated activities can alter these factors and interfere with key ecological functions such as spawning, rearing, and refugia. Site-specific factors (e.g., water clarity, current, depth, etc.) and the type and use of a given overwater structure determine the occurrence and magnitude of these impacts.

#### Potential Adverse Impacts

Overwater structures and associated developments may adversely affect EFH in a variety of ways, primarily by changes in ambient light conditions, alteration of the wave and current energy regime, and activities associated with the use and operation of the facilities (Nightingale and Simenstad 2001b).

Overwater structures create shade which reduces the light levels below the structure. The size, shape, and intensity of the shadow cast by a particular structure depends upon its height, width, construction materials, and orientation. High and narrow piers and docks produce narrower, more diffuse shadows than do low and wide structures. Increasing the numbers of pilings used to support a given pier enhances the shade pilings cast on the under-pier environment. In addition, less light is reflected underneath structures built with light-absorbing materials (e.g., wood) than under structures built with light-reflecting materials (e.g., concrete or steel). Structures that are oriented north-south produce a shadow that moves across the bottom throughout the day, resulting in a smaller area of permanent shade than those that are oriented east-west.

The shadow cast by an overwater structure affects both the plant and animal communities below the structure. Distributions of plants, invertebrates, and fishes appear severely limited in under-dock environments when compared to adjacent, unshaded, vegetated habitats. Light is the most important factor affecting aquatic plants. Under-pier light levels appear to fall below threshold amounts for the photosynthesis of diatoms, benthic algae, eelgrass, and associated epiphytes and other autotrophs. These photosynthesizers are an essential part of nearshore habitat and the estuarine and nearshore foodwebs that support many species of marine and estuarine fishes. Eelgrass and other macrophytes can be reduced or eliminated, even through partial shading of the substrate, and have little chance to recover.

Fishes rely on visual cues for spatial orientation, prey capture, schooling, predator avoidance, and migration. The reduced-light conditions found under an overwater structure limit the ability of fishes, especially juveniles and larvae, to perform these essential activities. Shading from overwater structures may also reduce prey organism abundance and the complexity of the habitat by reducing aquatic vegetation and phytoplankton abundance (Kahler et al. 2000, Haas et al. 2002). Glasby (1999) found that epibiotic assemblages on pier pilings at marinas subject to shading were markedly different than in surrounding areas. Other studies have shown shaded epibenthos to be reduced relative to that in open areas. These factors are thought to be responsible for the observed reductions in juvenile fish populations found under piers and the reduced growth and survival of fishes held in cages under piers, when compared to open habitats (Able et al. 1998, Duffy-Anderson and Able 1999).

The shadow cast by an overwater structure may increase predation on EFH-managed species by creating a light/dark interface that allows ambush predators to remain in a darkened area (barely visible to prey) and watch for prey to swim by against a bright background (high visibility) (Helfman 1981). Prey species moving around the structure are unable to see predators in the dark area under the structure and are more susceptible to predation. Furthermore, the reduced vegetation (i.e., eelgrass) densities associated with overwater structures decrease the available refugia from predators.

In addition to piscivorous predation, in-water structures (e.g., pilings) also provide perching platforms for avian predators such as double-crested cormorants (*Phalacrocorax auritis*), from which they can launch feeding forays or dry their plumage.

Wave energy and water transport alterations from overwater structures can impact the nearshore detrital foodweb by altering the size, distribution, and abundance of substrate and detrital materials. Disruption of longshore transport can alter substrate composition and present potential barriers to the natural processes that build spits and beaches and provide substrates required for plant propagation, fish and shellfish settlement and rearing, and forage fish spawning.

Pilings can alter adjacent substrates with increased shell deposition from piling communities and changes to substrate bathymetry (see Section 4.5). Changes in substrate type can alter the nature of the flora and fauna native to a given site. In the case of pilings, native dominant communities typically associated with

sand, gravel, mud, and eelgrass substrates are replaced by communities associated with shell hash substrates.

Treated wood used for pilings and docks releases contaminants into saltwater environs. PAHs are commonly released from creosote-treated wood. PAHs can cause a variety of deleterious effects (cancer, reproductive anomalies, immune dysfunction, and growth and development impairment) to exposed fish (Johnson et al. 1999, Johnson 2000, Stehr et al. 2000). Wood also is commonly treated with other chemicals such as ammoniacal copper zinc arsenate (ACZA) and chromated copper arsenate (CCA) (Poston 2001). These preservatives are known to leach into marine waters for a relatively short period of time after installation, but the rate of leaching varies considerably, depending on many factors. Concrete and steel, on the other hand, are relatively inert and do not leach contaminants into the water.

Construction and maintenance of overwater structures often involve driving pilings (see Section 4.5) and dredging navigation channels (see Section 4.1). Both activities may also adversely affect EFH.

While the effect of some individual overwater structures on EFH may be minimal, the overall impact may be substantial when considered cumulatively. The additive effects of these structures increase the overall magnitude of impact and reduce the ability of EFH to support native plant and animal communities.

#### Recommended Conservation Measures

The recommended conservation measures for overwater structures include the following:

1. Use upland boat storage whenever possible to minimize need for overwater structures.
2. Locate overwater structures in deep enough waters to avoid intertidal and shade impacts, to minimize or preclude dredging, to minimize groundings, and to avoid displacement of submerged aquatic vegetation, as determined by a preconstruction survey.
3. Design piers, docks, and floats to be multiuse facilities to reduce the overall number of such structures and to limit impacted nearshore habitat.
4. Incorporate measures that increase the ambient light transmission under piers and docks. These measures include, but are not limited to the following:
  - Maximizing the height of the structure and minimizing the width of the structure to decrease the shade footprint using grated decking material
  - Using solar tubes to direct light under the structure and glass blocks to direct sunlight under the structure
  - Illuminating the under-structure area with metal halide lamps and use of reflective paint or materials (e.g., concrete or steel instead of materials that absorb light such as wood) on the underside of the dock to reflect ambient light
  - Using the fewest number of pilings necessary to support the structures to allow light into under-pier areas and minimize impacts to the substrate
  - Aligning piers, docks, and floats in north-south orientation to allow the arc of the sun to cross perpendicular to the structure and to reduce the duration of light limitation
5. Use floating breakwaters whenever possible and remove them during periods of low dock use. Encourage seasonal use of docks and off-season haul-out.
6. Use waveboards to minimize the effects on littoral drift and benthic habitats.
7. Locate floats in deep water to avoid light limitation and grounding impacts to the intertidal zone.
8. Maintain at least 1 foot of water between the substrate and the bottom of the float.
9. Conduct in-water work when EFH-managed species and prey species are least likely to be impacted.

10. Avoid use of treated wood timbers or pilings to the extent practicable. Use alternative materials such as untreated wood, concrete, or steel.
11. Fit all pilings and navigational aids, such as moorings and channel markers, with devices to prevent perching by piscivorous bird species.
12. Orient night lighting to avoid illumination of the surrounding waters.
13. Mitigate for unavoidable impacts to benthic habitats. Mitigation should be adequate, monitored, and adaptively managed.

#### **4.7 Flood Control/Shoreline Protection**

Protecting riverine and estuarine communities from flooding events can result in varying degrees of change in the physical, chemical, and biological characteristics of existing shoreline and riparian habitat. The use of dikes and berms can also have long-term adverse effects on tidal marsh and estuarine habitats. Tidal marshes are highly variable, but typically have freshwater vegetation at the landward side, saltwater vegetation at the seaward side, and gradients of species inbetween that are in equilibrium with the prevailing climatic, hydrographic, geological, and biological features of the coast. These systems normally drain through highly dendritic tidal creeks that empty into the bay or estuary. Freshwater entering along the upper edges of the marsh drains across the surface and enters the tidal creeks. Structures placed for coastal shoreline protection include, but are not limited to, concrete or wood seawalls, rip-rap revetments (sloping piles of rock placed against the toe of the dune or bluff in danger of erosion from wave action), dynamic cobble revetments (natural cobble placed on an eroding beach to dissipate wave energy and prevent sand loss), vegetative plantings, and sandbags.

##### Potential Adverse Impacts

Dikes, levees, ditches, or other water controls at the upper end of a tidal marsh can cut off all tributaries feeding the marsh, preventing freshwater flushing and annual flushing, annual renewal of sediments and nutrients, and the formation of new marshes. Water controls within the marsh proper intercept and carry away freshwater drainage, block freshwater from flowing across seaward portions of the marsh, increase the speed of runoff of freshwater to the bay or estuary, lower the water table, permit saltwater intrusion into the marsh proper, and create migration barriers for aquatic species. In deeper channels where reducing conditions prevail, large quantities of hydrogen sulfide are produced. These quantities are toxic to marsh grasses and other aquatic life. Acid conditions of these channels can also result in release of heavy metals from the sediments

Long-term effects on the tidal marsh include land subsidence (sometimes even submergence), soil compaction, conversion to terrestrial vegetation, greatly reduced invertebrate populations, and general loss of productive wetland characteristics. Loss of these low-salinity environments reduces estuarine fertility, restricts suitable habitat for aquatic species, and creates abnormally high salinity during drought years. Low-salinity environments form a barrier that prevents the entrance of many marine species, including competitors, predators, parasites, and pathogens.

Armoring of shorelines to prevent erosion and to maintain or create shoreline real estate simplifies habitats, reduces the amount of intertidal habitat, and affects nearshore processes and the ecology of numerous species (Williams and Thom 2001). Hydraulic effects on the shoreline include increased energy seaward of the armoring, reflected wave energy, dry beach narrowing, substrate coarsening, beach steepening, changes in sediment storage capacity, loss of organic debris, and downdrift sediment starvation (Williams and Thom 2001). Installation of breakwaters and jetties can result in community changes from burial or removal of resident biota; changes in cover and preferred prey species; and predator attraction (Williams and Thom 2001). As with armoring, breakwaters and jetties modify

hydrology and nearshore sediment transport, as well as movement of larval forms of many species (Williams and Thom 2001).

#### Recommended Conservation Measures

The recommended conservation measures for flood control and shoreline protection include the following:

1. Minimize the loss of riparian habitats as much as possible.
2. Do not undertake diking and draining of tidal marshlands and estuaries unless a satisfactory compensatory mitigation plan is in effect and monitored.
3. Wherever possible, use soft approaches (such as beach nourishment, vegetative plantings, and placement of large woody debris) to shoreline modifications.
4. Include efforts to preserve and enhance EFH by providing new gravel for spawning areas, removing barriers to natural fish passage, and using weirs, grade control structures, and low-flow channels to provide the proper depth and velocity for fish.
5. Construct a low-flow channel to facilitate fish passage and help maintain water temperature in reaches where water velocities require armoring of the riverbed.
6. Replace in-stream fish habitat by providing rootwads, deflector logs, boulders, and rock weirs and by planting shaded riverine aquatic cover vegetation.
7. Use an adaptive management plan with ecological indicators to oversee monitoring and to ensure that mitigation objectives are met. Take corrective action as needed.

#### **4.8 Water Control Structures**

Many coastal areas of the Pacific Northwest use water control structures (WCSs) such as pumping stations and tidegates to regulate water levels in nearshore and estuary settings. WCSs enable certain agricultural crops to survive through floods, maintain high water tables, and manage the threat of saltwater intrusion. In some cases, infrastructures such as roads, industrial and residential developments, and sewer treatment plants have been built because of the enhanced drainage. These structures have been installed in streams, blind and distributary sloughs, and marsh/wetlands that are within estuarine and nearshore areas.

Tide gates typically have been installed on culverts passing through levees, dikes, and berms to prevent tidal inundation in areas landward of the berms. As the tide backs up and closes the tide gate, fish passage upstream is blocked. As the tide turns and begins to flow out, or the river level drops, a conventional tide gate opens a little but often not enough to allow upstream passage. The tidegate may also open with such velocity that constitutes a complete or partial blockage (Charland 1998). Pump stations are used to maintain more consistent control of water levels in nearshore and estuary settings. Some pumps are also used in conjunction with tide gates; many act as dams by stopping tidal or river stage levels, thus extending the capacity of the drainage system. While the design and operation of these structures vary, they generally pump surface water from the drainage system to the respective receiving body.

#### Potential Adverse Impacts

Adverse effects to EFH from the installation and operation of WCSs can occur through (1) partially or completely blocked habitat, (2) altered water chemistry composition through suppressed mixing of fresh and saltwater, (3) decreased sediment and nutrient delivery, and (4) degraded water quality through thermal loading.

Various life stages of some EFH-managed species use nearshore and estuarine habitats, and food produced from these areas in the form of small fish and other aquatic organisms are important for overall food web function (PFMC 1998, PFMC 2003). WCSs can limit or eliminate habitat access to areas that may be important for food sources and refuge from predators of these species.

Depending on their location, WCSs alter the normal circulation and mixing of fresh and saltwater. Estuaries are biologically rich and productive areas, partly because of the complex gradient of the fresh and salt water mixing process. Estuaries accumulate nutrients such as potassium and nitrogen, which are concentrated and recycled in a repeating interactive process by which the incoming tidal water resuspends nutrients at the fresh-saltwater interface while moving them back up the estuary to meet the seaward-moving, land-based nutrients (Day 1989). Estuarine food chains are extremely complex and are sensitive to alterations in the physical and chemical range of stresses (Day 1989). Loss or disruption of one element can have a cascading effect on species presence and productivity. The inhibition of the gradual mixing of salt and fresh water and nutrients over the original volume of habitat can decrease the overall productivity of the estuary and may cause prey community changes.

Often WCSs impound water for various amounts of time, which can lead to premature sediment and nutrient deposition and cause a subsequent need to dredge behind the structure. Sediment deposition within estuarine and nearshore areas is important for beach nourishment, and sediments often serve as absorptive surfaces for nutrients.

Impounded water can result in increased thermal loading which, in turn, can interfere with physiological processes, cause behavioral changes, and enhance disease enhancement (Bell 1986). Increased thermal loading can also cause increased microbial activity and vegetative growth, which in turn can deplete levels of dissolved oxygen (Waldichuk 1993, Spence et al. 1996). These impacts may combine to affect entire aquatic systems by changing primary and secondary productivity, community respiration, species composition, biomass, and nutrient dynamics (Hall et al. 1978). These effects, while perhaps more acute in the regulated watercourse, can nonetheless be manifested in the receiving body as well, particularly in areas where much of the historic estuary habitat is regulated by WCSs.

### Recommended Conservation Measures

The recommended conservation measures for water control structures include the following:

1. Avoid installing new WCSs. In some cases, tidegates that replace dams or pump stations (those which completely block habitat) can improve habitat conditions by enhancing fish passage and water circulation.
2. Design WCSs to enhance habitat access and water circulation.
3. Assess habitat potential or value behind the WCS by investigating current and potential aquatic vegetation, the volume and depth of the water body, the amount and timing of freshwater inflow, the presence of upland rearing and spawning habitat, and the relative salinity of the water body.
4. Assess the hydrology of the regulated land's tolerance for increased water exchange. The assessment should account for active management of the WCS to allow increased water exchange during critical periods. Existing programs that compensate landowners for lost production of land can be investigated (such as the Conservation Reserve Enhancement Program administered by the United States Department of Agriculture [UDSA]) if appropriate.
5. Design WCSs to mimic natural water exchange velocities. This can be done by maximizing the conveyance of water through increased width, thus reducing flow velocities when the gates are open.
6. Use WCS materials that are nontoxic and noncorrosive. Do not use treated wood.
7. Stabilize associated banks through bioengineering, minimizing the use of riprap and incorporating native materials as appropriate.

8. Install WCS during low-flow periods and low-tide stages; incorporate appropriate erosion and sediment control BMPs, and have an equipment spill and containment plan and appropriate materials on site.
9. Monitor WCS operations to assess impacts on water temperatures, dissolved oxygen, and other applicable parameters. Use adaptive management to minimize impacts.

#### **4.9 Log Transfer Facilities/In-water Log Storage**

Rivers, estuaries, and bays were historically the primary ways to transport and store logs in the Pacific Northwest. Log storage within the bays and estuaries remains an issue in several Pacific Northwest bays. Using estuaries and bays and nearby uplands for storage of logs is common in Alaska, with most LTFs found in Southeast Alaska and a few located in Prince William Sound.

##### Potential Adverse Impacts

Log handling and storage in the estuary and intertidal zones of rivers can result in water quality degradation and modifications to habitat. An LTF is a facility that is wholly or partly constructed in waters of the United States and that is used to transfer commercially harvested logs to or from a vessel or log raft, including the formation of a log raft (EPA 2000). LTFs may include a crane, A-frame structure, conveyor, or a slide or ramp; they are used move logs into the water. Logs can also be placed in the water at the site by helicopters and barges. The physical adverse impacts from these structures are similar in many ways to those of floating docks and other “over-water” structures (Section 4.6).

EFH may also be physically impacted by activities associated with LTFs. Bark and wood debris may impact EFH as a result of the abrasion of log surfaces from transfer equipment. After the logs have entered the water, they usually are bundled into rafts and hooked to a tug for shipment. In the process, bark and other wood debris can pile up on the ocean floor. The piles can smother clams, mussels, some seaweed, kelp, and grasses, with the bark sometimes remaining for decades. Accumulation of bark debris in shallow and deep water environments has resulted in locally decreased epifaunal macrobenthos richness and abundance (Kirkpatrick et al. 1998, Jackson 1986), which can ultimately impact various life-stages of groundfish.

Log storage may also result in a significant release of soluble, organic compounds. Log bark may affect groundfish by significantly increasing oxygen demand within the area of accumulation (Pacific Northwest Pollution Control Council [PNPCC] 1971). High oxygen demand can lead to an anaerobic zone where toxic sulfide compounds are generated, particularly in brackish and marine waters. Leaching of soluble organic compounds also leads to cumulative oxygen demand and reduced visibility. Reduced oxygen levels, anaerobic conditions, and the presence of toxic sulfide compounds are presumed to lead to reduced production of groundfish species and their forage base. Anaerobic areas reduce available habitat. In addition, soils at onshore facilities where logs are decked are often contaminated with gasoline, diesel fuel, solvents, etc., from trucks and heavy equipment. These contaminants can leach into nearshore EFH.

The physical, chemical, and biological impacts of LTF operations can be substantially reduced by adherence to appropriate siting and operational constraints. In 1985, the Alaska Timber Task Force (ATTF) developed guidelines to “delineate the physical requirements necessary to construct a log transfer and associated facilities, and in context with requirements of applicable law and regulations, methods to avoid or control potential impacts from these facilities on water quality, aquatic and other resources.” Since 1985, the ATTF guidelines have been applied to new LTFs through the requirements of NPDES permits and other state and federal programs (EPA 1996). Adherence to guidelines such as the ATTF operational and siting guidelines and BMPs in the NPDES General Permit will reduce the (1) amount of bark and wood debris that enters the marine and coastal environment, (2) the potential for displacement or

harm to aquatic species, and (3) the accumulation of bark and wood debris on the ocean floor. The following conservation measures reflect those guidelines.

#### Recommended Conservation Measures

The recommended conservation measures for log transfer facilities and in-water log storage include the following:

1. Storage and handling of logs should be restricted or eliminated from waters where state and federal water quality standards cannot be met at all times.
2. Minimize potential impacts of log storage by employing effective bark and wood debris controls, collection, and disposal methods at log dumps, raft building areas, and mill-side handling zones; avoiding free-fall dumping of logs; using easy let-down devices for placing logs in the water; and bundling logs before water storage (bundles should not be broken except on land and at millside).
3. Storage of logs should not take place where they will ground at any time or shade aquatic vegetation.
4. Avoid siting log-storage areas and LTFs in sensitive habitat and areas important for specified species.
5. Site log-storage areas and LTFs in areas with good currents and tidal exchanges.
6. Recommend land-based storage sites with the goal of eliminating in-water storage of logs.
7. For the Alaska region, also see the following link: Log Transfer Facility (LTF) Guidelines: [http://www.fs.fed.us/r10/TLMP/F\\_PLAN/APPEND\\_G.PDF](http://www.fs.fed.us/r10/TLMP/F_PLAN/APPEND_G.PDF).

#### **4.10 Utility Line/Cables/Pipeline Installation**

With the continued development of coastal regions comes greater demand for the installation of cables, utility lines for power and other services, and pipelines for water, sewage, etc. The installation of pipelines, utility lines, and cables can have direct and indirect impacts on the offshore, nearshore, estuarine, wetland, beach, and rocky shore coastal zone habitats. The coastal zone can be as narrow as a few feet in some areas to hundreds of miles inland in others, and it is not just development in the nearshore coastal regions that can cause impacts. Many of the primary and direct impacts occur during the construction phase of installation, such as with the ground disturbance in the clearing of the right-of-way, access roads, and equipment staging areas. Indirect impacts can include increased turbidity, saltwater intrusion, accelerated erosion, and the introduction of urban and industrial pollutants.

#### Potential Adverse Impacts

Adverse effects on EFH from the installation of pipelines, utility lines, and cables can occur through (1) destruction of organisms and habitat, (2) turbidity impacts, (3) resuspension of contaminants, and (4) changes in hydrology.

Destruction of organisms and habitats can occur in pipeline or cable right-of-way. This destruction can lead to long-term or permanent damage depending on the degree and type of habitat disturbance and the mitigation measures employed. Shallow-water environments, rocky reefs, nearshore and offshore rises, salt and freshwater marshes (wetlands), and estuaries are more likely to be adversely impacted than open-water habitats. This is due to their higher sustained biomass and lower water volumes, which decrease their ability to dilute and disperse suspended sediments (Gowen 1978).

Because vegetated coastal wetlands provide forage for and protection of commercially important invertebrates and fish, marsh degradation due to plant mortality, soil erosion, or submergence will eventually decrease productivity. Vegetation loss and reduced soil elevation within pipeline construction corridors should be expected with the continued use of current double-ditching techniques (Polasek 1997).



Increased water turbidity from higher than normal sediment loading can result in decreased primary production. Depending on the time of year of the construction, adverse impacts can occur, such as during highly productive spring phytoplankton blooms or times when organisms are already under stressed conditions. Changes in turbidity can temporarily alter phytoplankton communities. Depending upon the severity of the turbidity, these changes in water clarity can affect the EFH habitat functions of species higher in the food chain.

Another impact is resuspension of contaminants such as heavy metals and pesticides from the sediment, which can have lethal effects (Gowen 1978). Spills of petroleum products, solvents, and other construction-related material can also adversely affect habitat.

Pipeline canals have the potential to change the hydrology of coastal areas by (1) facilitating rapid drainage of interior marshes during low tides or low precipitation, (2) reducing or interrupting freshwater inflow and associated littoral sediments, and (3) allowing saltwater to move farther inland during periods of high tides (Chabreck 1972). Saltwater intrusion into freshwater marshes often causes loss of salt-intolerant emergent and submerged aquatic plants (Chabreck 1972, Pezeshki 1987), erosion, and net loss of soil organic matter (Craig et al. 1979).

#### Recommended Conservation Measures

The recommended conservation measures for utility line, cables, and pipeline installation include the following:

1. Align crossings along the least environmentally damaging route. Sensitive habitats such as hard-bottom (e.g., rocky reefs), submerged aquatic vegetation, oyster reefs, emergent marsh, sand, and mud flats should be avoided. If unavoidable, compensatory mitigation should be implemented.
2. Use horizontal directional drilling where cables or pipelines would cross salt marsh, vegetated inter-tidal zones, or steep erodible bluff areas adjacent to the inter-tidal zone to avoid surface disturbances.
3. Avoid construction of permanent access channels since they disrupt natural drainage patterns and destroy wetlands through excavation, filling, and bank erosion.
4. Store and contain excavated material on uplands. If storage in wetlands or waters cannot be avoided, alternate stockpiles should be used to allow continuation of sheet flow. Stockpiled materials should be stored on construction cloth rather than bare marsh surfaces, sea grasses, or reefs.
5. Backfill excavated wetlands with either the same or comparable material capable of supporting similar wetland vegetation. Original marsh elevations should be restored. Topsoil and organic surface material such as root mats should be stockpiled separately and returned to the surface of the restored site. Adequate material should be used so that the proper preproject elevation is attained following settling and compaction of the material. If excavated materials are insufficient to accomplish this, similar particle-size material should be used to restore the trench to the required elevation. After backfilling, erosion protection measures should be implemented where needed.
6. Use existing rights-of-way whenever possible to lessen overall encroachment and disturbance of wetlands.
7. Bury pipelines and submerged cables where possible. Unburied pipelines, or pipelines buried in areas where scouring or wave activity eventually exposes them, run a much greater risk of damage leading to leaks or spills.
8. Remove inactive pipelines and submerged cables unless they are located in sensitive areas (e.g., marsh, reefs, sea grass, etc.) or located in areas that present no safety hazard. If allowed to remain in place, pipelines should be properly pigged, purged, filled with seawater, and capped before abandonment in place.
9. Use silt curtains or other type barriers to reduce turbidity and sedimentation if sea grass or oyster reefs occur at or near the project site. These silt barriers should extend at least 100 feet beyond the

- limits of the sea grass beds or oyster reefs. If sea grasses and oyster reefs cannot be avoided, pre- and post-construction surveys should be completed to determine project impacts and mitigation needs.
10. Access for equipment should be limited to the immediate project area. Tracked vehicles are preferred over wheeled vehicles. Consideration should be given to the use of mats and boards to avoid sensitive areas. Equipment operators should be cautioned to avoid sensitive areas. Sensitive areas should be clearly marked to ensure that equipment operators do not traverse them.
  11. Limit construction equipment to the minimum size necessary to complete the work. Shallow-draft equipment should be used to minimize effects and to eliminate the necessity for temporary access channels. The size of the pipeline trench proper should also be minimized. The push-ditch method, in which the trench is immediately backfilled, reduces the impact duration and should, therefore, be used when possible.
  12. Conduct construction during the time of year when it will have the least impact on sensitive habitats and species.
  13. Suspend transmission lines beneath existing bridges or conduct directional boring under streams to reduce the environmental impact. If transmission lines span streams, site towers at least 200 feet from streams.

#### Activities on the Continental Shelf

14. Shunt drill cuttings through a conduit and discharge near the sea floor, or transport ashore.
15. Locate drilling and production structures, including pipelines, at least one mile from the base of a hard-bottom habitat.
16. Conduct the following mitigation measures:
  - a) Bury pipelines at least 3 feet beneath the sea floor, whenever possible. Particular considerations (i.e., currents, ice scour) may require deeper burial or weighting to maintain adequate cover. Buried pipeline and cables should be examined periodically for maintenance of adequate earthen cover.
  - b) Where burial is not possible, such as in hard-bottomed areas, pipelines and cables should be attached to substrate to avoid unnecessary conflicts with fishing gear. Wherever possible, mark the route by using lighted buoys and/or lighted ranges on platforms to reduce the risk of damage to fishing gear and the pipelines.
  - c) Locate alignments along routes that will minimize damage to marine and estuarine habitat. Avoid laying cable over high-relief bottom habitat and across live bottom habitats such as coral and sponge. If coral or sponge habitats are encountered, NMFS is interested in position and description information.
  - d) Where user conflicts are likely, consult and coordinate with fishing stakeholder groups through the appropriate FMC during the route-planning process in order to minimize conflict.
17. Avoid all natural reefs and banks, as well as artificial reef areas. Hard-bottom areas should be avoided to permit cable or pipeline burial. If unavoidable, compensatory mitigation should be mitigated.

#### **4.11 Commercial Utilization of Habitat**

Productive embayments are often used for commercial culturing and harvesting operations. These locations provide warmer water temperatures and protected waters, serving as excellent growout sites for oyster and mussel culturing. These operations may occur in areas of productive eelgrass beds. The commercial harvest of nearshore giant kelp is another habitat type that is used. Giant kelp forest canopies serve as nurseries, feeding grounds, and/or shelter to a variety of groundfish species and their prey (Cross and Allen 1993, Feder et al. 1974, Foster and Schiel 1985). In addition, when kelp plants are naturally

broken free of their holdfasts, drift kelp is produced. Kelp detritus supports high secondary production and prey for many fishes (Vetter 1995).

#### Potential Adverse Impacts

Adverse impacts to EFH by operations that directly or indirectly use habitat include (1) discharge of organic waste/contaminants, (2) impacts to the seafloor bed, (3) risk of introducing undesirable species, (4) impacts on estuarine food webs, and (5) impacts on kelp forest communities.

The culture of estuarine and marine species in estuarine areas can reduce or degrade habitats used by native species, depending on the location and operation of these facilities. A major concern of culture operations is the discharge of organic waste. The introduction of antibiotics and other drugs in medicated feeds is also a concern. Wastes are composed primarily of feces and excess feed. The buildup of waste products into the receiving waters will depend on water depths and circulation patterns. The release of these wastes can introduce nutrients or organic materials into the surrounding water body and lead to high BOD, lowering dissolved oxygen levels and, thereby, potentially affecting the survival of many aquatic organisms in the area. Nutrient overloads at the discharge site can also induce changes in community composition and structure, potentially favoring one group of organisms to the detriment of other.

In the case of cage mariculture operations for grow-out operations, impacts to the seafloor below the cages or pens can occur. The build-up of organic materials on the sea floor can impact the composition and diversity of the bottom-dwelling community (e.g., prey organisms for EFH species). Growth of submerged aquatic vegetation, which can provide shelter and nursery habitat for a number of fish species and their prey, can be inhibited by shading effects. Disruption of eelgrass habitat by management activities (e.g., dumping of shell with spawn on eelgrass beds, damage to eelgrass due to subsequent water or wind shear against the sharp oyster shells, and repeated mechanical raking or trampling) associated with this category are also of concern, though few studies have documented impacts. It is known that hydraulic dredges used to harvest oysters in coastal bays with eelgrass habitat can cause long-term adverse impacts to eelgrass beds, reducing or eliminating the beds (Phillips 1984).

The rearing of non-native, ecologically undesirable species may pose a risk of escape or accidental release into areas adversely affecting the ecological balance. Escape or other release into the environment can result in competition with native, wild fish for food, mates, spawning sites, which, if followed by successful interbreeding with wild stocks, can result in genetic dilution. Escapees can also pose a risk of transmission of disease to wild stocks.

Concern has also been expressed about extensive shellfish culture in estuaries and its impact on estuarine food webs. Oysters are efficient filter feeders and can change the trophic structure by removal of the microalgae and zooplankton that are also the food source for salmon prey species. The extent of this effect, if any, is unknown, especially in light of the fact that native oysters were once present in large quantities and co-existed with other species. Some effects might also be offset by the structures that oyster shells create, which offer shelter for a diverse biota.

Kelp is harvested for several reasons, which include directly obtaining its byproducts and indirectly as a food source in abalone culturing and a substrate in the Pacific herring fishery. Harvesting can have a variety of possible impacts on the habitat functions provided by kelp canopies. For example, giant kelp provides refuge to prey resources used by some EFH species. The kelp canopy also serves as habitat for canopy-dwelling invertebrates and can enhance fish recruitment and abundance. Removal of the canopy may affect some species by potentially displacing young-of-the-year or juvenile rockfishes, for example (Miller and Geibel 1973).

## Recommended Conservation Measures

The recommended conservation measures for commercial utilization of habitat include the following:

1. Site mariculture operations away from subaquatic vegetation areas. Facilities should be isolated close-circuited and located in upland areas as often as possible. Tidally influenced wetlands should not be enclosed or impounded for mariculture purposes, including hatchery and grow-out operations. Siting of facilities should also take into account the size of the facility, proximity of wild fish stocks, migratory patterns, competing uses, hydrographic conditions, and upstream uses.
2. Determine benthic productivity by sampling prior to any operations. Areas of high productivity should be avoided to the maximum extent possible. Sampling design should be developed with input from local, state, Tribal, and federal resource agencies.
3. Investigate water depths and circulation patterns where cage mariculture operations are undertaken to ensure that conditions are adequate to preclude the buildup of waste products, excess feed, and chemical agents.
4. Undertake a thorough scientific review and risk assessment before any non-native species are introduced. Any net pen structure should have small enough webbing to prevent entanglement by prey species. Mitigation should be provided for the areas impacted by the facility.
5. Encourage research into the timing of fish recruitment to help canopies and the response of canopy-dwelling juvenile groundfish to help harvesting operations to minimize potential adverse impacts to canopy habitat function.
6. Encourage development of harvesting methods to minimize impacts on plant communities such as the destruction of canopy-dwelling invertebrates and the loss of food and/or habitat to fish populations during harvesting operations.
7. Mitigation for unavoidable, extensive, or permanent loss of plant communities should be provided.

## **5.0 COASTAL/MARINE ACTIVITIES**

### **5.1 Point Source Discharge**

Point-source discharges from municipal sewage treatment facilities or storm water discharges are controlled through the EPA's mandated regulations under the CWA and by state water regulations. The primary concerns associated with municipal point-source discharges involve treatment levels needed to attain acceptable nutrient inputs and overloading of treatment systems due to rapid development of the coastal zone. Storm drains are contaminated from communities using settling and storage ponds, street runoff, harbor activities, and honey buckets. Annually, wastewater facilities introduce large volumes of untreated excrement and chlorine through sewage outfall lines, as well as releasing treated freshwater into the nation's waters. This can significantly alter pH levels of marine waters (Council 1999).

#### Potential Adverse Impacts

There are many potential impacts from point-source discharge, but point-source discharges and resulting altered water quality in aquatic environments do not necessarily result in adverse impacts to either marine resources or EFH. Because most point-source discharges are regulated by the state or EPA, effects to receiving waters are generally considered in those cases. Point-source discharges can adversely affect EFH by (1) reducing habitat functions necessary for growth to maturity, (2) modifying community structure, (3) bioaccumulation, and (4) modifying habitat.

At certain concentrations, point-source discharges can alter the following properties of ecosystems and associated communities: diversity, nutrient and energy transfer, productivity, biomass, density, stability, connectivity, and species richness and evenness. Pollution effects may be related to changes in water

flow, pH, hardness, dissolved oxygen, and other parameters that affect individuals, populations, and communities. Sewage, fertilizers, and de-icing chemicals (e.g., glycols, urea) are examples of common urban pollutants that decompose with high biological or chemical oxygen demand (Council 1999).

Point-source discharges, at certain concentrations, can alter the following characteristics of finfish, shellfish, and related organisms: growth, visual acuity, swimming speed, equilibrium, feeding rate, response time to stimuli, predation rate, photosynthetic rate, spawning seasons, migration routes, and resistance to disease and parasites. Additionally, zones of low dissolved oxygen resulting from their decomposition can retard growth of salmon eggs, larvae, and juveniles and may delay or block smolt and adult migration. Sewage and fertilizers also introduce nutrients into urban drainages that drive algal and bacterial blooms, which may smother incubating salmon or produce toxins as they grow and die. Thermal effluents from industrial sites and removal of riparian vegetation from streambanks can degrade salmon habitat by allowing solar warming of water. Heavy metals, petroleum hydrocarbons, chlorinated hydrocarbons, and other chemical wastes can be toxic to salmonids and their food, and they can inhibit salmon movement and habitat use in streams (Council 1999).

Elevated salinity levels from desalination plants also have to be considered. While studies have shown that elevated salinity levels may not produce toxic effects (Bay and Greenstein 1994), peripheral effects of pollution may include forcing rearing fish into areas of high predation. Conversely, an influx of treated freshwater from municipal wastewater plants may force rearing fish into habitat with less than optimal salinity for growth (Council 1999).

Point discharges may affect the growth, survival, and condition of EFH-managed species and prey species if high levels of contaminants (e.g., chlorinated hydrocarbons, trace metals, PAHs, pesticides, and herbicides) are discharged. If contaminants are present, they may be absorbed across the gills or concentrated through bioaccumulation as contaminated prey is consumed (Raco-Rands 1996). Many heavy metals and persistent organic compounds such as pesticides and polychlorinated biphenyls tend to adhere to solid particles discharged from outfalls. As the particles are deposited, these compounds or their degradation products (which may be equally or more toxic than the parent compounds) can enter the EFH foodchain by bioaccumulating in benthic organisms at much higher concentrations than in the surrounding waters (Stein et al. 1995). Due to burrowing, diffusion, and other upward transport mechanisms that move buried contaminants to the surface layers and eventually to the water column, pelagic and nektonic biota may also be exposed to contaminated sediments through mobilization into the water column.

Discharge sites may also modify habitat by creating adverse impacts to sensitive areas such as freshwater shorelines and wetlands, emergent marshes, sea grasses, and kelp beds if located improperly. Extreme discharge velocities of effluent may also cause scouring at the discharge point, as well as entraining particulates and thereby creating turbidity plumes. These turbidity plumes of suspended particulates can reduce light penetration and lower the rate of photosynthesis and the primary productivity of an aquatic area while elevated turbidity persists. The contents of the suspended material can react with the dissolved oxygen in the water and result in oxygen depletion, or smother submerged aquatic vegetation sites including eelgrass beds and kelp beds. Accumulation of outfall sediments may also alter the composition and abundance of infaunal or epibenthic invertebrate communities (Ferraro 1991). Pollutants, either suspended in the water column (e.g., nitrogen, contaminants, fine sediments) or settled on the bottom, can affect habitat. Many benthic organisms are quite sensitive to grain size, and accumulation of sediments can also submerge food organisms (Section 4.2.2).

## Recommended Conservation Measures

The recommended conservation measures for point source discharges include the following:

1. Locate discharge points in coastal waters well away from shellfish beds, sea grass beds, coral reefs, and other similar fragile and productive habitats.
2. Reduce potentially high velocities by diffusing effluent to acceptable velocities.
3. Determine benthic productivity by sampling prior to any construction activity related to installation of new or modified facilities. Outfall design (e.g., modeling concentrations within the predicted plume or likely extent of deposition along a productive nearshore) should be developed with input from appropriate resource and Tribal agencies.
4. Provide for mitigation when degradation or loss of habitat occurs from placement and operation of the outfall structure and pipeline.
5. Institute source-control programs that effectively reduce noxious materials to avoid introducing these materials into the waste stream.
6. Ensure compliance with pollutant discharges regulated through discharge permits which set effluent discharge limitations and/or specify operation procedures, performance standards, or BMPs. These efforts rely on the implementation of BMPs to control polluted runoff (EPA 1993).
8. Discharges should be treated to the maximum extent practicable, including implementation of up-to-date methodologies for reducing discharges of biocides (e.g., chlorine) and other toxic substances.
9. Use land-treatment and upland disposal/storage techniques where possible. Use of vegetated wetlands as natural filters and pollutant assimilators for large-scale discharges should be limited to those instances where other less damaging alternatives are not available, and the overall environmental and ecological suitability of such actions has been demonstrated.
10. Avoid siting pipelines and treatment facilities in wetlands and streams. Since pipelines and treatment facilities are not water-dependent with regard to positioning, it is not essential that they be placed in wetlands or other fragile coastal habitats. Avoiding placement of pipelines within streambeds and wetlands will also reduce inadvertent infiltration into conveyance systems and retain natural hydrology of local streams and wetlands.

### **5.2 Fish Processing Waste—Shoreside and Vessel Operation**

Seafood processing facilities are either shore-based facilities discharging through stationary outfalls or mobile vessels engaged in the processing of fresh or frozen seafood (Science Applications International Corporation [SAIC] 2001). Discharge of fish waste from shoreside and vessel processing has occurred in marine waters since the 1800s (Council 1999). With the exception of fresh market fish, some form of processing involving butchering, evisceration, precooking, or cooking is necessary to bring the catch to market. Precooking or blanching facilitates the removal of skin, bone, shell, gills, and other materials. Depending on the species, the cleaning operation may be manual, mechanical, or a combination of both (EPA 1974). Seafood processing facilities generally consist of mechanisms to offload the harvest from fishing boats; tanks to hold the seafood until the processing lines are ready to accept them; processing lines, process water, and waste collection systems; treatment and discharge facilities; processed seafood storage areas; and necessary support facilities such as electrical generators, boilers, retorts, water desalinators, offices, and living quarters. In addition, marinas that cater to patrons who fish a large amount can produce an equally large quantity of fish waste at the marina from fish cleaning.

### Potential Adverse Impacts

Generally, seafood processing wastes consist of biodegradable materials that contain high concentrations of soluble organic material. Seafood processing operations have the potential to adversely affect EFH

through (1) direct and/or nonpoint source discharge, (2) particle suspension, and (3) increased turbidity and surface plumes.

Seafood processing operations have the potential to adversely affect EFH through the direct and/or nonpoint source discharge of nutrients, chemicals, fish by-products, and “stickwater” (water and entrained organics originating from the draining or pressing of steam-cooked fish products). EPA investigations show that impacts affecting water quality are direct functions of the receiving waters. In areas with strong currents and high tidal ranges, waste materials disperse rapidly. In areas of quieter waters, waste materials can accumulate and result in shell banks, sludge piles, dissolved oxygen depressions, and associated aesthetic problems (Stewart and Tangarone 1977). If adequate disposal facilities are not available at marinas that generate a large amount of fish waste, there is a potential for disposal of fish waste in areas without enough flushing to prevent decomposition and the resulting dissolved oxygen depression (EPA 1993).

Processors discharging fish waste are required to have EPA-issued NPDES permits. Various water quality standards, including those for BOD, total suspended solids (TSS), fecal coliform bacteria (FC), oil and grease, pH, and temperature, are all considerations in the issuance of such permits. Although fish waste, including heads, viscera, and bones, is biodegradable, fish parts that are ground to fine particles may remain suspended for some time, thereby overburdening habitats from particle suspension (Council 1999). Such pollutants have the potential to adversely impact EFH. The wide differences in habitats, types of processors, and seafood processing methods define those impacts and can also prevent the effective use of technology-based effluent limits.

In certain areas such as Alaska, seafood processors are allowed to deposit fish parts in a Zone of Deposit (ZOD) (EPA 2001). This can remove benthic habitat from the environment, reduce locally associated invertebrate populations, and lower dissolved oxygen levels in overlying waters. Impacts from accumulated processing wastes are not limited to the area covered by the ZOD. Severe anoxic and reducing conditions occur adjacent to effluent piles (EPA 1979). Examples of localized damage to benthic environment include several acres of bottom-driven anoxic by piles of decomposing waste up to 26 feet deep. Juvenile and adult stages of flatfish are drawn to these areas for food sources. One effect of this attraction may lead to increased predation on juvenile fish species by other flatfishes, diving seabirds, and marine mammals drawn to the food source (Council 1999). However, due to the difficulty in monitoring these areas, impacts to species can go undetected.

Scum and foam from seafood waste deposits can also occur on the water surface and/or increase turbidity. Increased turbidity decreases light penetration into the water column, reducing primary production. Reduced primary production decreases the amount of food available for consumption by higher trophic level organisms. In addition, stickwater takes the form of a fine gel or slime that can concentrate on surface waters and move onshore to cover intertidal areas.

#### Recommended Conservation Measures

The recommended conservation measures for fish processing waste include the following:

1. Base effluent limitations on site-specific water quality EFH concerns to the maximum extent practicable.
2. Avoid the practice of discharging untreated solid and liquid waste directly into the environment. Use of secondary or wastewater treatment systems should be encouraged where possible.
3. Designation of new ZODs should not be allowed. Options to eliminate or reduce ZODs at existing facilities should be explored.
4. Control stickwater by physical or chemical methods.

5. Promote sound fish waste management through a combination of fish-cleaning restrictions, public education, and proper disposal of fish waste.
6. Encourage the alternative use of fish processing wastes (e.g., fertilizer for agriculture and animal feed).
7. Options for additional research should be explored. There is not much current research on which to base management decisions about habitat. Some improvements in waste processing have occurred, but the technology-based effluent guidelines have not changed in 20 years.
8. Locate new plants outside rearing and nursery habitat. Monitor both biological and chemical changes to the site.

### **5.3 Water Intake Structures/Discharge Plumes**

The withdrawal of riverine, estuarine, and marine waters by water intake structures is a common aquatic activity. Water may be withdrawn and used to cool coastal power generating stations, used as a source of water for agricultural purposes, and, more recently, used as a source of potable water for desalinization plant operations. In the case of power plants and desalinization plants, the subsequent discharge of heated and/or chemically-treated discharge water can also occur.

#### Potential Adverse Impacts

Adverse impacts to EFH from water intake structures and effluent discharges can interfere or disrupt EFH functions in the source or receiving waters by (1) entrainment, (2) impingement, (3) discharge, (4) operation and maintenance, and (5) construction-related impacts.

Entrainment is the withdrawal of aquatic organisms along with the cooling water into the cooling system. These organisms are usually the egg and larval stages of managed species and their prey. Entrainment can subject these life stages to adverse conditions resulting from the effects of increased heat, antifouling chemicals, physical abrasion, rapid pressure changes, and other detrimental effects. Consequently, diverting water without adequate screening prevents that portion of the EFH from providing important habitat functions necessary for the early life stages of managed living marine resources and their prey. Long-term water withdrawal may adversely affect fish and shellfish populations by adding another source of mortality to the early life stage, which often determines recruitment and year-class strength (Travnichek et al. 1993).

Impingement occurs to organisms that are too large to pass through in-plant screening devices and instead become stuck or impinged against the screening device or remain in the forebay sections of the system until they are removed by other means (Grimes 1975, Hanson et al. 1977, Helvey and Dorn 1987, Helvey 1985, Langford et al. 1978, Moazzam and Rizvi 1980). The organisms cannot escape due to the water flow that either pushes them against the screen or prevents them from exiting the intake tunnel. Similar to entrainment, the withdrawal of water can entrapped particular species, especially when visual acuity is reduced (Helvey 1985). This condition reduces the suitability of the source waters to provide normal EFH functions necessary for subadult and adult life stages of managed living marine resources and their prey.

Thermal effluents in inshore habitat can cause severe problems by directly altering the benthic community or killing marine organisms, especially larval fish. Temperature influences biochemical processes of the environment and the behavior (e.g., migration) and physiology (e.g., metabolism) of marine organisms (Blaxter 1969). Further, the proper functioning of sensitive areas may be affected by the action of intakes as selective predators, resulting in cascading negative consequences as observed by the overexploitation of local fish populations in coral-reef fish communities (Carr et al. 2002).



Other impacts to aquatic habitats can result from construction related activities (e.g., dewatering, dredging, etc.) (see Section 4.1), as well as routine operation and maintenance activities. A broad range of impacts associated with these activities depend on the specific design and needs of the system. For example, dredging activities can cause turbidity, degraded water quality, noise, and substrate alterations. Many of these impacts can be reduced or eliminated through the use of various techniques, procedures, or technologies, but some may not be fully eliminated except by eliminating the activity itself.

Power plants using once-through cooling biocides, such as sodium hypochlorite and sodium bisulfate, may be used periodically to clean the intake and discharge structures. Chlorine is extremely toxic to aquatic life.

#### Recommended Conservation Measures

The recommended conservation measures for water intake structures and discharge plumes include the following:

1. Locate facilities that rely on surface waters for cooling in areas other than estuaries, inlets, heads of submarine canyons, rock reefs, or small coastal embayments where EFH species or their prey concentrate. Discharge points should be located in areas that have low concentrations of living marine resources. The discharge points should also incorporate cooling towers to control temperature and employ sufficient safeguards to ensure against release of blow-down pollutants into the aquatic environment in concentrations that reduce the quality of EFH.
2. Design intake structures to minimize entrainment or impingement. Velocity caps that produce horizontal intake/discharge currents should be employed and intake velocities across the intake screen should not exceed 0.5 foot per second.
3. Design power plant cooling structures to meet the best technology available requirements (BTAs) as developed pursuant to Section 316(b) of the CWA. Use of alternative cooling strategies, such as closed cooling systems (e.g., dry cooling), should be used to completely avoid entrainment/impingement impacts in all industries that require cooling water. When alternative cooling strategies prove infeasible, other BTAs may include, but are not limited to, fish diversion or avoidance systems, fish return systems that convey organisms away from the intake and mechanical screen systems that prevent organisms from entering the intake system, and habitat restoration measures.
4. Regulate discharge temperatures (both heated and cooled effluent) so they do not appreciably alter the temperature that could cause a change in species assemblages and ecosystem function in the receiving waters. Strategies should be implemented to diffuse the heated effluent.
5. Avoid the use of biocides (e.g., chlorine) to prevent fouling where possible. The least damaging antifouling alternatives should be implemented.
6. Mitigate for impacts related to power plants and other industries requiring cooling water. Mitigation should compensate for the net loss of EFH habitat functions from placement and operation of the intake and discharge structures. Mitigation should be provided for the loss of habitat from placement of the intake structure and delivery pipeline, the loss of fish larvae and eggs that may be entrained by large intake systems, and the degradation or loss of habitat from placement of the outfall structure and pipeline, as well as the treated water plume.
7. Treat all discharge water from outfall structures to meet state water quality water standards at the terminus of the pipe. Pipes should extend a substantial distance offshore and be buried deep enough to not affect shoreline processes. Buildings and associated structures should be set well back from the shoreline to preclude the need for bank armoring.

## 5.4 Oil/Gas Exploration/Development/Production

Offshore exploration, development, and production of natural gas and oil reserves have been, and continues to be, an important aspect of the U.S. economy. As demand for energy resources grows, the debate over trying to balance the development of oil and gas resources and the protection of the environment will also continue. Projections indicate that U.S. demand for oil will increase by 1.3 percent per year between 1995 and 2020. Gas consumption is projected to increase by an average of 1.6 percent during the same time frame (Waisley 1998). Much of the 1.9 billion acres within the offshore jurisdiction of the United States remain unexplored Oil and Gas Technologies for the Arctic and Deepwater [OGTAD] 1985). It is also expected that some of the older oil and gas platforms in operation will reach the end of their productive life in the near future. The question of decommissioning is also an issue.

### Potential Adverse Impacts

Offshore oil and gas operations can be classified into exploration, development, and production activities. Petroleum exploration/development/production occurs in varying water depths and usually over soft-bottom substrates, although hard-bottom habitats may be present in the general vicinity. These areas are subject to an assortment of physical, chemical, and biological disturbances. These disturbances include (1) noise from seismic surveys, vessel traffic, and construction of drilling platforms or islands; (2) physical alterations to habitat from the construction, presence, and eventual decommissioning and removal of facilities, such as islands or platforms, storage and production facilities, and pipelines to onshore common carrier pipelines, storage facilities, or refineries; (3) waste discharges, including well drilling fluids, produced waters, surface runoff and deck drainage, domestic waste waters generated from the offshore facility, solid-waste from wells (drilling muds and cuttings) and other trash and debris from human activities associated with the facility; (4) oil spills; and (5) platform storage and pipeline decommissioning (Council 1999, Helvey 2002).

Noise sources may generate sound pressure that can disrupt or damage marine life. Oil and gas activities may generate noise from drilling activities, construction, production facility operations, seismic exploration, and supply vessel and barge movements (Section 4.5). The impacts of oil exploration-related seismic energy releases may interrupt and cause fish to disperse from the acoustic pulse with possible disruption to their feeding patterns. It is known that noise in the marine environment may adversely affect marine mammals by causing them to change behavior (movement, feeding), interfere with echolocation and communication, or may result in injury to hearing organs (Richardson et al. 1995).

Activities such as vessel anchoring, platform or artificial island construction, pipeline laying (Section 4.10), dredging, and pipeline burial can alter bottom habitat by altering substrates used for feeding or shelter. Disturbances to the associated epifaunal communities, which may provide feeding or predator escape habitat, can also result. Benthic organisms, especially prey species, may recolonize disturbed areas, but this may not occur if the composition of the substrate is drastically changed or if facilities are left in place after production ends. Dredging, trenching, and pipelaying generate spoils that may be disposed of on land or the marine environment where sedimentation may smother benthic habitat and organisms. Most of these activities associated with oil and gas operations, however, are conducted under permits and regulations that require companies to minimize impacts or to avoid construction or other disturbances in sensitive marine habitats (Section 4.2.2).

The discharge of drilling muds and cuttings can result in varying degrees of change on the sea floor and affect feeding, nursery, and shelter habitat for various life stages of managed species. Drilling muds and cuttings may adversely affect bottom-dwelling organisms at the site by burial of immobile forms or forcing mobile forms to migrate. Exploratory and construction activities may also result in resuspension of fine-grained mineral particles, usually smaller than silt, in the water column. These suspended

particulates can reduce light penetration and lower the rate of photosynthesis and the primary productivity of the aquatic area especially if suspended for lengthy intervals. Groundfish and other fish species can suffer reduced feeding ability leading to limited growth if high levels of suspended particulates persist. The contents of the suspended material can react with the dissolved oxygen in the water and result in oxygen depletion. In addition, the discharge of oil drilling muds can change the chemical and physical characteristics of benthic sediments at the disposal site by introducing toxic chemical constituents. Changes in the clarity and the addition of contaminants can reduce or eliminate the suitability of water bodies as habitat for fish species and their prey (NMFS 1998).

Oil spills are a serious potential source of contamination to the marine environment from oil and gas development. Offshore oil and gas development will inevitably result in some oil entering the environment. Most spills are expected to be of small size, although there is a potential for large spills to occur. Many factors determine the degree of damage from a spill, including the type of oil, size and duration of the spill, geographic location of the spill, and the season. Oil is toxic to all marine organisms at high concentrations, but certain species are more sensitive than others. In general, the early life stages (eggs and larvae) are most sensitive, juveniles are less sensitive, and adults are least sensitive (Rice et al. 2000).

Lost oil can affect habitats and living marine resources in both large and small quantities. Accidental discharge of oil can occur during almost any stage of exploration, development, or production on the outer continental shelf (OCS) or in nearshore coastal areas. Oil spills can occur from many possible sources, including equipment malfunction, ship collisions, pipeline breaks, other human error, or severe storms. Oil spills can also be attributed to support activities associated with product recovery and transportation. In addition to crude oil spills, chemical, diesel, and other contaminant spills, accidental discharge can also occur with OCS activities (Council 1999).

Chronic small oil spills are a potential problem because residual oil can build up in sediments and affect living marine resources. Low levels of petroleum components (e.g., PAHs) from such chronic pollution can accumulate in salmon tissues and cause lethal and sublethal effects, particularly at the embryo stage. Effects on fish from low-level chronic exposure may increase embryo mortality, reduce marine growth (Heintz et al. 2000), or increase straying away from natal streams by returning adults (Wertheimer et al. 2000).

It is possible for a major oil spill (i.e., 50,000 barrels) to produce a surface slick covering up to several hundred square kilometers of surface area. If the oil spill moves toward land, habitats and species could be affected by the loading of oil into the near shore environment. In the initial hours after a large spill, aromatic hydrocarbons would generally be at toxic levels to some organisms. Beneath and surrounding the surface slick, there would be some oil-contaminated waters. Physical and biological forces act to reduce oil concentrations with depth and distance (Council 1999); generally the lighter fraction aromatic hydrocarbons evaporate rapidly, particularly during periods of high wind and wave activity. Heavier oil fractions may settle through the water column. Suspended sediment can adsorb and carry oil to the seabed. Hydrocarbons may be solubilized by wave action, which may enhance adsorption to sediments, which then sink to the seabed, contaminating benthic sediments. Carls et al. (2003) demonstrated that tides and the resultant hydraulic gradients provide a mechanism for groundwater transport of soluble and slightly soluble contaminants (such as oil) from beaches surrounding streams into the hyporheic zone where pink salmon eggs incubate. Oil may reach nearshore areas and affect productive nursery grounds or areas containing high densities of fish eggs and larvae. An oil spill near an especially important habitat (e.g., a gyre where fish or invertebrate larvae are concentrated) could also result in a disproportionately high loss of a population of marine organisms. Other aquatic biota at risk would be eggs, larvae, and other planktonic organisms in the upper seawater column. Because they cannot actively avoid exposure, their small size means they absorb contaminants quickly, and their proximity to the seasurface means they

may be vulnerable to photo-enhanced toxicity effects, which can increase the toxicity of hydrocarbons several fold (Barron et al. 2003). In addition, oil spills may interrupt commercial or subsistence fishing activities.

Habitats that are susceptible to damage from spill oil include not just the low-energy coastal bays and estuaries where oil may accumulate but also high-energy cobble environments where oil is driven into sediments through wave action. Many of the beaches in Prince William Sound with the highest persistence of oil following the *Exxon Valdez* oil spill were high-energy environments containing large cobbles overlain with boulders. These beaches were pounded by storm waves that drove the oil into and well below the surface (Michel and Hayes 1999). Oil that mixes into bottom sediments can persist for years. Subsurface oil was still detected in beach sediments of Prince William Sound 12 years after the *Exxon Valdez* oil spill, much of it unweathered and more prevalent in the lower intertidal biotic zone than at higher tidal elevations (Short et al. 2002). Additional concern is the unknown impact of an oil-related event near and within the ice. The water column adjacent to the ice edge is stable. This stabilization (or stratification) would allow relatively quick transport of oil to the sea floor. Additionally, oil trapped in ice could significantly affect habitat after the initial event, months or years later, and even into a different region (Council 1999).

Residual oil from a spill can remain toxic for long periods. Petroleum is a complex mixture of alkanes and aromatic hydrocarbons, of which the alkyl-substituted and multi-ring PAHs are the most toxic and persistent. Following weathering, the aromatic fraction of oil is dominated by PAHs as the lighter aromatic components evaporate or are degraded. Because of low solubility in water, the large PAH concentrations probably contribute little to acute toxicity of oil-water solutions. Lipophilic PAH, however, may cause physiological injury if it accumulates in tissues after exposure (Carls et al. 1999, Heintz et al. 2000). Even when concentrations of oil are sufficiently diluted not to be physically damaging to marine organisms, it still may be detected by them, and it may alter certain behavior patterns.

Oil and gas platforms may be comprised of a lattice-work of pilings, beams, and pipes that support diverse fish and invertebrate populations and are considered de facto artificial reefs (Love and Westphal 1990, Love et al. 1994, Love et al. 1999, Helvey 2002). Because decommissioning includes plugging and abandoning all wells and removing the platforms and associated structures from the ocean, impacts to EFH can result during removal. Impacts during the demolition phase may include underwater sound pressure waves (see Section 4.5.1) and impacts on marine organisms; removal of structures may remove habitat for invertebrates and fish that associate with midwater structures. In some areas of the United States, offshore oil and gas platforms are allowed to remain after decommissioning, thereby providing permanent habitat for some organisms.

The potential disturbances and associated adverse impacts on the marine environment has been reduced through the operating procedures required by regulatory agencies and, in many cases, self imposed by facilities operators. Most of the activities associated with oil and gas operations are conducted under permits and regulations that require companies to minimize impacts or avoid construction in sensitive marine habitats. New technological advancements result in improved operating practices that reduce the potential for impacts. For example, the discharge of muds and cuttings is being phased out of modern oil and gas production programs; generally such byproducts of exploration or development are ground into finer materials and injected into wells that penetrate subsea reservoir strata and do not enter the marine environment.

#### Recommended Conservation Measures

Oil and gas exploration, development, and production can be conducted in a manner that minimizes adverse impacts on the marine environment. Over the past several decades, government agencies and

petroleum production companies have developed operating procedures that reduce potential adverse effects; these procedures are generally required through permits. The following are recommended measures that should be considered in permitting future oil and gas operations:

1. Conduct pre-project biological surveys in consultation with NMFS to determine the extent and composition of biological populations or habitat in the proposed production area. On the basis of the site-specific surveys, a determination will be made whether or not the operations are likely to have an adverse effect on EFH, or that a special biological population/habitat does not exist. Based on the information in the surveys, the following may be recommended:
  - a. Locate the site of operation.
  - b. Redesign facilities to accommodate habitat concerns.
  - c. Operate during those periods of time, as established in consultation with NMFS, that do not adversely affect biological resources.
  - d. Modify operations to ensure that significant biological populations or habitats deserving protection are not affected.
2. Limit the discharge of produced waters into marine and estuaries environments. Re-inject produced waters into the oil formation whenever possible.
3. Avoid discharge of muds and cuttings into the marine and estuarine environment. Use methods to grind and re-inject such wastes down an approved injection well or use onshore disposal wherever possible. When not possible, provide for a monitoring plan to quantitatively assess whether effluent discharges are meeting the needs of EFH.
4. Limit placement of causeways or structures in the nearshore marine environment.
5. Encourage the use of geographic response strategies that identify EFH and environmentally sensitive areas and identify appropriate cleanup methods to include the prestaging of response equipment.
6. Use methods to transport oil and gas that limit the need for handling in environmentally sensitive areas, including EFH.
7. Prohibit drilling of the first development well into the targeted hydrocarbon formations during hazardous or sensitive environmental conditions, such as broken ice.
8. Prohibit drilling of exploration wells into untested formations during hazardous or sensitive environmental conditions.
9. Provide for monitoring and leak detection systems that preclude oil and gas from entering the environment. The following measures should be taken:
  - a. Use systems that detect spills and leaks as rapidly as technologically possible so that action can be taken to avoid or reduce the effect to EFH.
  - b. Use maximum precautions to eliminate pipeline failure caused by external forces.
10. Evaluate impacts to habitat during the decommissioning phase, including impacts during the demolition phase and impacts resulting from permanent habitat losses.

## **5.5 Habitat Restoration/Enhancement**

Habitat loss and degradation are major, long-term threats to the sustainability of fishery resources (NMFS 2002). Viable coastal and estuarine habitats are important to maintaining healthy fish stocks. Good water quality and quantity, appropriate substrate, ample food sources and substantial hiding places are needed to sustain fisheries. Restoration and/or enhancement of coastal and riverine habitat that supports managed fisheries and their prey will assist in sustaining and rebuilding fisheries stocks and recovering certain threatened or endangered species by increasing or improving ecological structure and functions. Habitat restoration/enhancement may include, but is not limited to, improvement of coastal wetland tidal exchange or re-establishment of historic hydrology; dam or berm removal; fish passage barrier removal/modification; road related sediment source reduction; natural or artificial reef/substrate/habitat creation; establishment or repair of riparian buffer zones and improvement of freshwater habitats that support anadromous fishes; planting of native coastal wetland and submerged aquatic vegetation; creation

of oyster reefs; and improvements to feeding, shade or refuge, spawning and rearing areas that are essential to fisheries.

### Potential Adverse Impacts

The implementation of restoration/enhancement activities may have localized and temporary adverse impacts on EFH. Possible impacts can include (1) localized nonpoint source pollution such as influx of sediment or nutrients, (2) interference with spawning and migration periods, (3) temporary or permanent removal feeding opportunities, and (4) indirect effects from actual construction portions of the activity.

Unless proper precautions are taken, upland related restoration projects can contribute to nonpoint source pollution. Such concerns should be addressed as part of the planning process (Section 2.1). Particular in-water projects may interfere with spawning periods or impede migratory corridors and should be addressed accordingly. Projects may also have an affect on the feeding behavior of managed species. For instance, if dredging is involved, benthic food resources may be affected (Section 4.1). Impacts can occur from individuals conducting the restoration, especially at staging areas, as part of accessing the restoration site, or the actual restoration techniques employed. Particular impacts can result from water quality impacts from individuals conducting the restoration, excessive foot traffic, diving techniques, equipment handling, boat anchoring, and planting techniques.

The use of artificial reefs is a popular form of habitat enhancement, but it can also affect the aquatic environment through the loss of habitat upon which the reef material is placed or the use of inappropriate materials in construction. Usually, reef materials are set upon flat sand bottoms or “biological deserts,” which end up burying or smothering bottom-dwelling organisms at the site or even preventing mobile forms (e.g., benthic-oriented fish species) from utilizing the area as habitat. Some materials may be inappropriate for the marine environment (e.g., automobile tires; compressed incinerator ash) and can serve as sources of toxic releases or physical damage to existing habitat when breaking free of their anchoring systems (Collins et al. 1994).

### Recommended Conservation Measures

The recommended conservation measures for habitat restoration and enhancement include the following:

1. Use BMPs to minimize and avoid all potential impacts to EFH during restoration activities. This conservation measure requires the use of BMPs during restoration activities to reduce impacts from project implementation. BMPs should include, but are not limited to, the following:
  - a. Measures to protect the water column—turbidity curtains, haybales, and erosion mats should be used.
  - b. Staging areas—areas used for staging will be planned in advance and kept to a minimum size.
  - c. Buffer areas around sensitive resources—rare plants, archeological sites, etc., will be flagged and avoided.
  - d. Invasive species—invasive plant and animal species should be removed from the proposed action area prior to commencement of work. Only native plant species should be planted. Measures to ensure native vegetation or revegetation success will be identified and implemented (Section 4.4).
  - e. Ingress/egress areas—temporary access pathways will be established prior to restoration activities to minimize adverse impacts from project implementation.
2. Avoid restoration work during critical fish windows to reduce direct impacts to important ecological functions such as spawning, nursery, and migration. This conservation measure requires scheduling projects when managed species are not expected in the area. These periods should be determined prior to project implementation to reduce or avoid any potential impacts.

3. Provide adequate training and education to volunteers and project contractors to ensure minimal impact to the restoration site. Volunteers should be trained in the use of low-impact techniques for planting, equipment handling, and any other activities associated with the restoration. Proper diving techniques need to be used by volunteer divers.
4. Conduct monitoring before, during, and after project implementation to ensure compliance with project design and restoration criteria. If immediate post-construction monitoring reveals that unavoidable impacts to EFH have occurred, appropriate coordination with NMFS should occur to determine appropriate response measures, possibly including mitigation.
5. Mitigate fully any unavoidable damage to EFH during project implementation and accomplish within reasonable period of time after the impacts occurred.
6. Remove and, if necessary, restore any temporary access pathways and staging areas used in the restoration effort.
7. Determine benthic productivity by sampling prior to any construction activity in the case of subtidal enhancement (e.g., artificial reefs). Areas of high productivity should be avoided to the maximum extent possible. Sampling design should be developed with input from state and federal resource agencies. Prior to construction, an evaluation of the impact resulting from the change in habitat (sand bottom to rocky reef, etc.) should be performed. Post-construction monitoring should examine the effectiveness of the structures for increasing habitat productivity.

## **5.6 Marine Mining**

Mining activity, which is also described in Section 3.1.1 and Section 3.1.2, can lead to the direct loss of EFH for certain species. Offshore mining, such as the extraction of gravel and gold in the Bering Sea (EBS) and the mining gravel of gravel from beaches, can increase turbidity of water and, thus, the resuspension of organic materials could affect less motile organisms (i.e., eggs and recently hatched larvae) in the area. Benthic habitats could be damaged or destroyed by these actions. Mining of large quantities of beach gravel can significantly affect the removal, transport, and deposition of sand and gravel along the shore, both at the mining site and down current (Council 1999). Neither the future extent of this activity nor the effects of such mortality on the abundance of marine species is known.

### **Potential Adverse Impacts**

Mining practices that can affect EFH include physical impacts from intertidal dredging and chemical impacts from the use of additives such as flocculants (Council 1999). Impacts include the removal of substrates that serve as habitat for fish and invertebrates; creation (or conversion) of habitats to less productive or uninhabitable sites, such as anoxic holes or silt bottom; burial of productive habitats, such as in near shore disposal sites (as in beach nourishment); release of harmful or toxic materials either in association with actual mining, or in connection with machinery and materials used for mining; creation of harmful turbidity levels; and adverse modification of hydrologic conditions so as to cause erosion of desirable habitats. Submarine disposal of mine tailings can also alter the behavior of marine organisms. Submarine mine tailings may not provide suitable habitat for some benthic organisms. In laboratory experiments, benthic dwelling flatfishes (Johnson et al. 1998a) and crabs (Johnson et al. 1998b) strongly avoided mine tailings.

During beach gravel mining, water turbidity increases and the resuspension of organic materials can affect less motile organisms (i.e., eggs and recently hatched larvae) in the area. Benthic habitats can be damaged or destroyed by these actions. Changes in bathymetry and bottom type may also cause alteration in population and migrations patterns (Hurme and Pullen 1988).

## Recommended Conservation Measures

The recommended conservation measures for marine mining include the following:

1. Avoid mining in waters containing EFH.
2. Minimize the areal extent and depth of extraction to minimize recolonization times.
3. Limit sand mining and beach nourishment in areas with EFH.
4. Monitor turbidity during operations and cease operations if turbidity exceeds predetermined threshold levels. Use sediment or turbidity curtains to limit the spread of suspended sediments and minimize the area affected.
5. Monitor the number of individual mining operations to avoid and minimize cumulative impacts. For instance, three mining operations in an intertidal area could impact EFH, whereas one may not. Also, disturbance of previously contaminated mining areas threaten an additional loss of EFH.

### **5.7 Persistent Organic Pollutants (North Pacific and Alaska Specific)**

The single biggest pollution threat to marine waters in Alaska is the deposition of persistent pollutants from remote sources. North Pacific and Alaskan marine waters are perceived as pristine because most of Alaska's 6,640 miles of coastline are devoid of point-source pollution, unlike much of North America. Effluents from pulp mills, marinas and boat harbors, municipal outfalls, and other industrial activities are generally considered to be the primary sources of contamination in Alaskan waters, so most efforts at monitoring and mitigation have been focused on the local level. The only major regional pollution event was the *Exxon Valdez* oil spill in 1989, a contaminant threat that has abated considerably over the last 14 years. However, there is an increasing body of evidence suggesting that the greatest contaminant threat in Alaska comes from atmospheric and marine transport of contaminants from areas quite distant from Alaska.

The geography of Alaska makes it particularly vulnerable to contaminants volatilized from Asia. During winter, the Aleutian low pressure cell steers air from Southeast Asia into the EBS and northern Gulf of Alaska (GOA), bringing precipitation along the way. When this air meets the mountains along Alaska's southern coast, more precipitation occurs, bringing entrained contaminants from the atmosphere into the marine ecosystem or coastal/interior ecosystems. Thus, pesticides applied in to crops in Southeast Asia can be volatilized into the air, bound to suspended particulates, transported in the atmosphere to Alaska, and deposited in snow or rain directly into marine ecosystems or indirectly from freshwater flow to nearshore waters. Revolatilization of these compounds is inhibited by the cold temperatures associated with Alaskan latitudes, resulting in a net accumulation of these compounds in northern habitats. This same distillation process also transfers volatilized contaminants from the atmosphere to the Pacific at lower latitudes, and ocean currents also deliver the contaminants to Alaska. Concentrations will be very low, but there will extensive geographical marine or land areas to act as "cold" deposit zones.

### Potential Adverse Impacts

The effect of these transport mechanisms has been the appearance of persistent organic contaminants in northern latitudes, despite the absence of local sources. A good demonstration of global transport into northern latitudes is the presence of dichloro-diphenyl-trichloroethanes (DDTs) in the blubber of ring seals in the western Canadian Arctic (Addison and Smith 1996). DDT and its congeners were first observed in these seals during the early 1970s. The persistence DDTs in these seals through the 1990s, despite North American bans on DDT use in the 1970s, is evidence of continued deposition of DDT from countries still using this pesticide.



The existence of organic contaminants in biological tissues means these contaminants are being transported within the food webs in Alaskan fish habitats. For example, Ewald et al. (1998) found detectable levels of polychlorinated biphenyls (PCBs), DDTs, and other pesticides in the tissues of adult sockeye salmon returning to the Copper River. These fish apparently concentrated these contaminants in their tissues during their migration in the northern GOA and delivered them to their spawning habitats in the interior of Alaska. Obviously, avian and mammalian predators of these fish would further distribute these contaminants.

#### *Distribution of Contaminants in Marine Habitats*

A large variety of contaminants can be found in Alaska's marine environment, including persistent organic pollutants (POPs) and heavy metals. POPs are characterized as those with half-lives over 2 months, bioaccumulation factors greater than 5,000, potential for long-range transport, and capable of toxic effects. Currently, 12 classes of compounds are considered POPs and are regulated by Stockholm Convention on Persistent Organic Pollutants (Table 3). In addition to POPs, heavy metals are present in Alaskan habitats, including mercury (Hg), cadmium (Cd), chromium (Cr), arsenic (As), lead (Pb), and silver (Ag). Contaminants found in Alaskan marine mammals sampled between southeastern Alaska and the Aleutian and Pribilof Islands include PCBs, DDT, chlordanes, hexachlorocyclohexanes (HCHs), hexachlorobenzene (HCB), dieldrin, butyltins, arsenic, mercury, cadmium, and lead (Barron and Heintz in press). With over 100,000 chemicals on the market and an additional 1,000 to 2,000 new ones introduced annually, there are likely other toxic compounds in the environment whose concentrations are increasing. In addition, combustion and industrial processes result in the inadvertent production of unregulated chemicals (Arctic Monitoring and Assessment Program [AMAP] 2002).

There have been few large-scale evaluations of the spatial or temporal patterns to contamination in Alaska's marine environment. Most effort at monitoring contaminant loads in Alaskan waters has focused on Arctic habitats where there is evidence that PCBs and DDTs have declined over the last 25 years (AMAP 2002). Recently, Beckmen et al. (2001) reported on the concentrations of PCBs in sea lion scats collected from around the GOA. These data suggest that sea lion prey in the eastern Aleutian Islands (AI) have greater PCB loads than prey near Kodiak, Cook Inlet, and Prince William Sound. Prey from the latter three locations also have lower PCB loads than those from southeastern Alaska. Some of

**Table 3.** The Twelve Persistent Organic Pollutants Regulated by the POPs Treaty

	Common Name	Effect on Organisms
Pesticides	Dieldrin	Reproductive impairment; renal and liver damage
	Aldrin	Neurological damage; reproductive impairment
	Chlordane	Altered hormone function
	DDT/DDE	Neurological damage; hormonal disruption; reproductive impairment
	Endrin	Developmental abnormalities
	Heptachlor	Liver damage; hormonal changes
	Hexachlorobenzene	Reduced embryo weights in herring gulls
	Mirex	Kidney lesions in fish
Polychlorinated biphenyls	Toxaphene	"Broken-back" syndrome in fish
	PCBs	Poor reproductive success Impaired immune function
	Dioxins	Immune suppression; hormonal dysfunction; developmental impairment
Industrial and Incineration Byproducts	Furans	Developmental impairment; increased abortions

Source: World Federation of Public Health Associations 2000. Persistent organic pollutants and human health. Washington, DC.

the relatively high values observed in the eastern Aleutians may reflect the addition of PCB point-source inputs at specific sites (Barron and Heintz in press), but it would seem unlikely that a few point sources could account for the general elevated state of PCB loads in the entire Aleutians.

Temporal studies provide little information because they are quite limited in the number of locations evaluated and samples collected. The mechanism, however, by which contaminants are delivered to the Alaskan marine environment guarantees that as long as the contaminants are released, they will be found in Alaskan waters (Wania and Mackay 1999). For example, the types of PCBs found in seals from sites near the Russian coast are consistent with those used in Russian electrical equipment (Muir and Norstrom 2000). Contributions of contaminants by marine transport will continue for some time. More water soluble organic contaminants like HCHs are slower to accumulate in Arctic and subarctic food webs and appear to be increasing (Wania and Mackay 1999). Mercury appears to be higher in more recent samples (mid 1990s) than in the 1980s and 1970s, and rates of Hg accumulation also appear to be higher than 10 to 20 years ago (Muir et al. 1999). Polybrominated diphenyl ethers (PBDEs) also appear to be increasing in marine mammals (Ikonomou et al. 2002) and may surpass PCBs as the most prevalent POP in arctic habitats.

#### *Factors Leading to Higher Contaminant Loads*

The trophic structure of Alaskan marine food webs coupled with the tendency of contaminants to accumulate in Alaskan habitats causes apex predators to concentrate significant amounts of POPs in their tissues. Organisms occupying the top trophic levels in a food web bioaccumulate the highest concentrations of contaminants (Ruus et al. 2002). For example, the total PCB concentration in seal-eating killer whales sampled near Kenai Fjords National Monument was one to two orders of magnitude greater than fish-eating killer whales, indicating the significance of their trophic position (Ylitalo et al. 2001a). Further, seal-eating killer whale PCB loads were greater than the loads typically associated with belugas from the St. Lawrence River, while those of resident, fish-eating killer whales were consistent with loads observed in harbor seals in Puget Sound (Ylitalo et al. 2001a). The few data available on organisms at lower trophic levels in Alaska's marine habitats indicate these species experience relatively low contaminant loads (de Brito et al. 2002, Aono et al. 1997, Kawano et al. 1986). Thus, Alaskan killer whales are likely accumulating loads of contaminants from remote sources that are consistent with those of marine mammals living near heavily contaminated urban areas as a result of their high trophic position. While this interpretation fails to account for differences in life stage, sex, or analytical method, it illustrates the need for more detailed information about this region.

This issue is particularly relevant when the contaminant loads experienced by Alaskan natives subsisting on foods derived from marine habitats are considered. In one study, the total PCB concentration (not lipid adjusted) in serum collected from Aleutian males aged 45 to 54 averaged 8.7 parts per billion (Alaska Division of Public Health 2003). By comparison, the concentrations in similarly aged males from around the Great Lakes that also consumed large amounts of fish (more than 52 meals per year) averaged 4.8 parts per billion (Hanrahan et al. 1999). Reference males in the latter study were demographically similar, ate less fish, and averaged 1.5 parts per billion. The relatively high level for the Alaskan natives is likely the result of their trophic position relative to that of the Great Lakes fishers. Alaskan natives living in the Aleutians with subsistence lifestyles probably consume seals and fish, leading to a trophic position above that of Great Lakes fishers who likely consume more grains and plant materials than Aleutian natives.

A second contributing factor to increased contaminant loads among apex predators in Alaska is their relatively long life. Contaminant loads increase with age in fish (Vuorinen et al. 2002), Steller sea lions (O'Hara 2001, Ylitalo et al. 2001b), and humans (Alaska Division of Public Health 2003). Female pinnipeds in the EBS and northern GOA typically begin reproducing at 5 years of age, (Riedman 1990)

allowing time for significant accumulation of contaminants, especially because pinnipeds eat relatively large (i.e., old) prey. For example, the pollock consumed by Steller sea lions average 393 mm and Atka mackerel 323 mm (Zeppelin et al. 2003), which translate to fish ages of approximately 3 to 5 years old. These sizes, however, were at the low end of the size distribution, indicating that sea lions can eat much older prey. Vuorinen et al. (2002) reported a sevenfold increase in POP loads of sprat between ages 2 and 10, demonstrating the increased potential for exposure associated with consuming older prey.

#### *Significance of Contaminant Loads*

It is not clear if the levels of contaminants in Alaskan waters are causing deleterious effects to populations, because research in this area is still in its infancy. Relatively small and spotty contaminant surveys have established that POPs are present in Alaskan waters, forage, and predators. No comprehensive geographical and temporal studies have been done to date to examine trends or sources of variation. The potential for the problem has been exposed; the extent and significance remain to be determined.

The potential for significant effects is most likely greatest among apex predators. Contamination is probably wide spread among forage species at low levels, but apex predators are likely to be the most affected as a result of their longevity, lipid storage, and the relatively high concentrations they bear. In mammals, it is most likely that lipophilic contaminants would have the greatest impacts on first-born young. The accumulation of contaminants in females increases with age, but decreases after females reach reproductive age. This is the result of their transfer of contaminants to their offspring in milk. This process has been reported for sea lions, fur seals (Beckmen et al. 1999), and humans (Yang et al. 2002). This process occurs repeatedly for each offspring, consequently, the first-born offspring receives adult level contaminant loads during its most sensitive developmental stage. Beckmen et al. (1999) reported that first-born northern fur seal pups of primiparous mothers had higher PCB levels in their blood than pups of multiparous mothers. This higher load was correlated with a reduced ability to form antibodies to tetanus, along with reduced concentrations of thyroxine and vitamin A in their blood. Barron and Heintz (in press) compared reported PCB loads in juvenile Steller sea lions with loads known to cause deleterious effects in other pinnipeds and concluded some sea lions in the mid-1980s likely experienced immunological impairment. Assessing impacts on humans is more difficult and controversial. While the acute impacts of contaminants on humans are known, the long-term impacts following neonatal exposure have not been explored.

Recent declines in apex predator populations in the EBS and northern GOA may be related to contaminant loading in the region. Over the course of the last 25 years, the populations of Steller sea lions, harbor seals, northern fur seals, and many birds have declined. The reasons underlying these declines are likely complex and may not be the same for all species. For example, the decline in Steller sea lions is presumed to have resulted from nutritional stress, but more recent evidence suggests other factors, including contaminants, may be limiting their recovery (De Master et al. 2001). Contaminants are unlikely to be causing acutely toxic effects in the regions. Sublethal impacts of contaminants, however, could be working indirectly to impair populations through reduced immune function (Beckmen 2001) or reproduction (Reinjders 1986). Both of these characters are displayed by Steller sea lion populations from the affected region. York et al. (1996) attributed continuing declines in affected populations to a failure to recruit offspring to maturity. Zenteno-Savin et al. (1997) reported elevated levels of haptoglobin, an acute phase reaction protein in the blood of Steller sea lions and harbor seals from affected populations relative to levels observed in stable or increasing populations. This protein is indicative of non-specific stressors that could include injury, disease, or toxicity. Thus, a recent panel was unable to reject contaminants as a factor contributing to the failed recovery of Steller sea lion populations (Barron and Heintz 2001).

Impacts may also occur at lower trophic levels, but there has been even less research in this area. Atlantic salmon in the Baltic Sea and salmonids in the Great Lakes have both experienced a common syndrome variously named M74 or early mortality syndrome. The syndrome is characterized by low thiamine content in eggs, resulting in near complete mortality of affected brood years. While cause for the reduced thiamine content in spawning adults remains unknown, increased levels of PCB and dibenzofurans and dibenzo-dioxins were correlated with the onset of the disease in Baltic salmon (Vuorinen et al. 2002).

It is important to remember that the impacts of persistent contaminants on populations in Alaskan waters are not likely to be acute. The impacts are more likely be expressed as sublethal impacts in apparently healthy animals. These sublethal impacts ultimately lead to reduced reproductive fitness or decreased survival to maturity and, therefore, they manifest themselves indirectly. Science is certain that the physical properties of these compounds couple with global climate patterns to ensure that they will be deposited in Alaska habitats, while maintaining their toxicity and perfusing through Alaskan food webs, which include some of the most valuable fisheries on the planet. What is uncertain is how these compounds impact the health of organisms deriving sustenance from those food webs and how those impacts might feed back into the food web.

### Recommended Conservation Measures

No mitigation strategies are proposed at this time relative to contaminants. There are too many unknowns. It is known that POP contaminants are present in Alaska waters and forage species, and in predators up through apex predators, but the significance of the present loads is not known. Also, the relative concentrations in forage species (Pollock for example) from the EBS, near Russia, or Northern GOA is not known. Comprehensive studies on a geographical, temporal, or on a widespread species scale to determine first if there is relationship between contaminant loads and population changes have not been conducted. It is suspected that POP contaminants may contribute to poor recovery in some species such as Steller sea lions, but mitigation strategies, whether they would be changes in fishing regulations or international regulation to curb contaminant releases, will likely need a better research foundation to support changes.

### **REFERENCES**

- Abbe, T.B. and D.R. Montgomery. 1996. "Large woody debris jams, channel hydraulics, and habitat formation in large rivers." *Regulated Rivers: Research and Management*. 12:201-221.
- Abbott, R. and E. Bing-Sawyer. 2002. Assessment of pile driving impacts on the Sacramento blackfish (*Othodon microlepidotus*). Draft report prepared for Caltrans District 4. October 10, 2002.
- Able, K.W., J.P. Manderson, and A.L. Studholme. 1998. "The distribution of shallow water juvenile fishes in an urban estuary: the effects of man-made structures in the lower Hudson River." *Estuaries*. 21:731-744.
- Addison, R.F. and T.G. Smith. 1996. "Trends in organochlorine residue concentrations in ringed seal (*Phoca hispida*) from Holman, Northwest Territories 1972-1991." *Arctic*. 51:253-561.
- Alaska Division of Public Health. 2003. PCB blood test results from St. Lawrence Island recommendations for consumption of traditional foods. State of Alaska Epidemiology Bulletin. 7(1): 1-5.
- American Fisheries Society. 2000. AFS Policy Statement #13: Effects of Surface Mining on Aquatic Resources in North America (Revised). (Abbreviated) ([http://www.fisheries.org/Public\\_Affairs/Policy\\_Statements/ps13ashtml](http://www.fisheries.org/Public_Affairs/Policy_Statements/ps13ashtml)).

- Aono, S, S. Tanabe, Y. Fujise, H. Kato, and R. Tatsukawa. 1997. "Persistent organochlorines in minke whale (*Balaenoptera acutorostrata*) and their prey species from the Antarctic and the north Pacific." *Env. Poll.* 98:81-89.
- Arctic Monitoring and Assessment Programme (AMAP). 2002. Arctic pollution 2002: Persistent organic pollutants, heavy metals, radioactivity, human health, changing pathways. Arctic Monitoring and Assessment Programme. Oslo Norway. pp. iii - 111.
- Arkoosh, M.R., E. Casillas, E. Clemons, P. Huffman, A.N. Kagley, T. Collier, and J.E. Stein. 2001. "Increased susceptibility of juvenile chinook salmon (*Oncorhynchus tshawytscha*) to vibriosis after exposure to chlorinated and aromatic compounds found in contaminated urban estuaries." *Journal of Aquatic Animal Health.* 13:257-268.
- Barron, M.G., M.G. Carls, J.W. Short, and S.D. Rice. 2003. Photoenhanced toxicity of aqueous phase and chemically dispersed weathered Alaska North Slope crude oil to Pacific herring eggs and larvae. *Environmental Toxicology and Chemistry* 22(3):650-660.
- Barron, M.G. and R.A. Heintz. 2001. Workshop to assess contaminant impacts on Steller sea lions in Alaska. Anchorage, AK September 5-6, 2001. National Marine Fisheries Service, Auke Bay Laboratory. Juneau, AK.
- Barron, M.G. and R.A. Heintz. *In press.* "Contaminant exposure and effects in pinnipeds: implications for steller sea lion declines in Alaska." *Sci. Tot. Env.*
- Bay, S. and D. Greenstein. 1994. Toxic effects of elevated salinity and desalination waste brine. In J. Cross, ed. Southern California Coastal Water Research Project, Annual Report 1992-93, pp. 149-153. SCCWRP, Westminster, CA.
- Beckmen, K.B. 2001. Blood organochlorines, immune function and health of free-ranging northern fur seal pups (*Callorhinus ursinus*). PhD. Dissertation. University of Alaska, Fairbanks. Fairbanks, Alaska. 151 pp.
- Beckmen, K.B., G.M. Ylitalo, R.G. Towell, M.M. Krahn, T.M. O'Hara, and J.E. Blake. 1999. "Factors affecting organochlorine contaminant concentrations in milk and blood of northern fur seal (*Callorhinus ursinus*) dams and pups from St. George Island, Alaska." *Sci. Total Environ.* 231:183-200.
- Beckmen, K.B., K.W. Pitcher, G.M. Ylitalo, M.M. Krahn, and K.A. Burek. 2001. Contaminants in Free-ranging Steller Sea Lions, 1998-2001: Organochlorines in Blood, Blubber, Feces and Prey. Workshop to Assess Contaminant Impacts on Steller Sea Lions in Alaska. Anchorage, AK, September 5-6, 2001. Sponsored by Auke Bay Laboratory.
- Beechie, T., E. Beamer, and L. Wasserman. 1994. "Estimating coho rearing habitat and smolt production losses in a large river basin, and implications for habitat restoration." *North American Journal of Fisheries Management.* 14(4):797-811.
- Belford, D.A. and W.R. Gould. 1989. "An evaluation of trout passage through six highway culverts in Montana." *North American Journal of Fisheries Management.* 9(4):437-445.
- Bell, M.C. 1986. Fisheries handbook of engineering requirements and biological criteria. U.S. Army, Corps of Engineers, Office of Chief Engineers, Fish Passage Development and Evaluation Program, Portland, Oregon.
- Benfield, M.C. and T. J. Minello. 1996. "Relative effects of turbidity and light intensity on reactive distance and feeding of an estuarine fish." *Environmental Biology of Fishes.* 46:211-216.
- Beschta, R.L., M.R. Pyles, A.E. Skaugset, and C.G. Surfleet. 2000. "Peakflow response to forest practices in the western Cascades of Oregon, U.S.A." *Journal of Hydrology.* 233:102-120.

- Beschta, R.L., R.E. Bilby, G.W. Brown, L.B. Holtby, and T.D. Hofstra. 1987. Stream temperature and aquatic habitat: fisheries and forestry interactions. In Salo, E., T. Cundy, eds. Streamside management: forestry and fishery interactions. Contribution 57. Seattle: University of Washington, College of Forest Resources: 191-232. 437-445.
- Bilby, R.E. and J.W. Ward. 1991. "Characteristics and function of large woody debris in streams draining old-growth, clear-cut, and second-growth forests in southwestern Washington." *Canadian Journal of Fisheries and Aquatic Sciences*. 48:2499-2508.
- Bilby, R.E., K. Sullivan, and S.H. Duncan. 1989. "The generation and fate of road-surface sediment in forested watersheds of southwestern Washington." *Forest Science*. 35:453-468.
- Bisson, P.A., J.L. Nielsen, and J.W. Ward. 1988. "Summer production of coho salmon stocked in Mount St. Helens streams 3-6 years after the 1980 eruption." *Transactions of the American Fisheries Society*. 117(4):322-335.
- Blanton, S.L., R.M. Thom, and J.A. Southard. 2001. Documentation of ferry terminal shading, substrate composition, and algal and eelgrass coverage. Letter Report prepared for University of Washington, School of Aquatic and Fishery Sciences, Seattle, Washington by Battelle Marine Sciences Laboratory, Sequim, Washington. 17 pp.
- Blaxter, J. H. S. 1969. Development: Eggs and larvae. In W. S. Hoar., and D. J. Randall. eds. Fish Physiology. Vol. 3. Academic Press, Inc. New York, NY. pp. 177-252.
- Bond, A.B., J.S. Stephens, Jr, D. Pondella, II, M.J. Allen, and M. Helvey. 1999. "A method for estimating marine habitat values based on fish guilds, with comparisons between sites in the Southern California Bight." *Bull. Mar. Science*. 64:219-242.
- Botkin, D., K. Cummins, T. Dunne, H. Regier, M. Sobel, L. Talbot, and L. Simpson. 1995. Status and Future of salmon of Western Oregon and Northern California: Findings and Options. Report #8. The Center for the Study of the Environment, Santa Barbara, California. 300 pp.
- Brady, N.C. and R.R. Weil. 1996. The nature and properties of soils. 11th ed. Prentice-Hall, Inc. Upper Saddle River, New Jersey 07458. 740 pp.
- Burroughs, E.R. Jr., G.R. Chalfant, and M.A. Townsend. 1976. Slope stability in road construction: a guide to the construction of stable roads in western Oregon and northern California. Portland, OR: U.S. Department of the Interior, Bureau of Land Management. 102 pp.
- Caltrans. 2001. Fisheries Impact Assessment, Pile Installation Demonstration Project for the San Francisco - Oakland Bay Bridge, East Span Seismic Safety Project, August 2001. 59 pp.
- Cardwell, R.D., M.I. Carr, and E.W. Sanborn. 1980. Water quality and flushing of five Puget Sound marinas. Technical Report No. 56. Washington Department of Fisheries Research and Development. Olympia, Washington. 77 pp.
- Carls, M.G., R.E. Thomas, and S.D. Rice. 2003. "Mechanism for transport of oil-contaminated water into pink salmon redds." *Mar. Ecol. Prog. Ser.* 248:245-255.
- Carls, M.G., S.D. Rice, and J.E. Hose. 1999. "Sensitivity of fish embryos to weathered crude oil: Part 1. Low level exposure during incubation causes malformations and genetic damage in larval Pacific herring (*Clupea pallasii*)." *Environmental Toxicology and Chemistry*. 18:481-493.
- Carlson, T.J., G. Ploskey, R.L. Johnson, R.P. Mueller, M.A. Weiland, and P.N. Johnson. 2001. Observations of the behavior and distribution of fish in relation to the Columbia River navigation channel and channel maintenance activities. Prepared for the U.S. Army, Corps of Engineers, Portland District by Pacific Northwest National Laboratory, U.S. Department of Energy, Richland, WA. 35 pp. + appendices.

- Carr, M.H., T.W. Anderson, and M.A. Hixon. 2002. Biodiversity, population regulation, and the stability of coral-reef fish communities. *Proceedings of the National Academy of Sciences*. 99:11241-11245.
- Castro, J. 2003. Geomorphologic impacts of culvert replacement and removal: avoiding channel incision. U.S. Fish and Wildlife Service—Oregon Fish and Wildlife Office, Portland, Oregon.
- Cederholm, C.J. and L.M. Reid. 1987. Impact of forest management on coho salmon (*Oncorhynchus kisutch*) populations of the Clearwater Rivers, Washington: A project summary. In *Streamside management: forestry and fishery interactions*. Salo, E.O, and T.W. Cundy, eds. College of Forest Resources, University of Washington, Seattle, Washington. University of Washington, Institute of Forest Resources. Contribution No. 57.
- Chabreck, R.H. 1972. Vegetation, water and soil characteristics of the Louisiana coastal region. Louisiana State University Agriculture Experiment Station. Baton Rouge, LA.
- Chapman, D.W. 1988. "Critical review of variables used to define effects of fines in redds of large salmonids." *Transactions of the American Fisheries Society*. 117(1):1-21.
- Charland, J. 1998. Tide Gate Modification for Fish Passage and Water Quality Enhancement. Tillamook Bay National Estuary Project. (503) 322-2222.
- Christie, D., E. Allen, and G. J. Jobsis. 1993. Diversions. In Bryan, C.F. and D.A. Rutherford eds. *Impacts on warmwater streams: Guidelines for evaluation*. pp. 181-186. Southern Division, American Fisheries Society, Little Rock, AR.
- Christopherson, A. and J. Wilson. 2002. Technical Letter Report Regarding the San Francisco-Oakland Bay Bridge East Span Project Noise Energy Attenuation Mitigation. Peratrovich, Nottingham & Drage, Inc. Anchorage, Alaska. 27 pp.
- Clancy, C.G. and D.R. Reichmuth. 1990. "A detachable fishway for steep culverts." *North American Journal of Fisheries Management*. 10(2):244-246.
- Clayton, J.L. 1983. Evaluating slope stability prior to road construction. Res. Pap. INT-307. Ogden, UT. U.S. Department of Agriculture, Forest Service, Intermountain Forest and Range Experiment Station. 6 pp.
- Cloern, J.E. 1987. "Turbidity as a control on phytoplankton biomass and productivity in estuaries." *Continental Shelf Research*. 7:1367-1381.
- Collins, B.D., D.R. Montgomery, and A.D. Haas. 2002. "Historical changes in the distribution and functions of large wood in Puget Lowland rivers." *Canadian Journal of Fisheries and Aquatic Sciences*. 59:66-76.
- Collins, B.D., D.R. Montgomery, and A.J. Sheikh. 2003. Reconstructing the historic riverine landscape of the Puget Lowland. pp 79-128. In Bolton, S, D.R. Montgomery, and D. Booth, eds. *Restoration of Puget Sound Rivers*. University of Washington Press. Seattle, WA.
- Collins, K.J., A.C. Jensen, A.P.M. Lockwood, and S.J. Lockwood. 1994. "Coastal structures, waste materials and fishery enhancement." *Bulletin of Marine Science*. 55(203):1240-1250.
- Corbett, C.W., M. Wahl, D.E. Porter, D. Edwards, and C. Moise. 1997. "Nonpoint source runoff modeling: A comparison of a forested watershed and an urban watershed o the South Carolina coast." *Journal of Experimental Marine Biology and Ecology*. 213(1):133-149.
- Council (North Pacific Fisheries Management Council). 1999. Environmental Assessment for Amendment 55 to the Fishery Management Plan for the Groundfish Fishery of the Bering Sea and Aleutian Islands Area; Amendment 55 to the Fishery Management Plan for Groundfish of the Gulf of Alaska; Amendment 8 to the Fishery Management Plan for the King and Tanner Crab

- Fisheries in the Bering Sea/Aleutian Islands; Amendment 5 to the Fishery Management Plan for Scallop Fisheries off Alaska; Amendment 5 to the Fishery Management Plan for the Salmon Fisheries in the EEZ off the Coast of Alaska, Essential Fish Habitat. 605 West 4th Ave, Suite 306, Anchorage, AK 99501-2252. 20 January.
- Craig, N.J., R.E. Turner, and J.W. Day, Jr. 1979. "Land loss in coastal Louisiana." *Environmental Management*. 3:134-144
- Cross, J. N. and L.G. Allen. 1993. Fishes. In Dailey, M.D, D.J. Reish, and J.W. Anderson, eds. Ecology of the Southern California Bight, Univ. Calif. Press, Berkeley, CA. pp.459-540.
- Culp, J.M. and R.W. Davies. 1983. An assessment of the effects of streambank clear cutting on macroinvertebrate communities in a managed watershed. Canadian Technical Report of Fisheries and Aquatic Sciences. No. 1208. 116 pp.
- Day, J.W, C.A.S. Hall, W.M. Kemp, and A. Yanez-Arancibia. 1989. *Estuarine Ecology*. John Wiley and Sons, New York. 558 pp.
- de Brito, A.P.X., D. Ueno, S. Takahasi, and S. Tanabe. 2002. "Contamination by organochlorine compounds in walleye pollock (*Theragra chalcogramma*) from the Bering Sea, Gulf of Alaska, and the Japan Sea." *Marine Pollution Bulletin*. 44:164-177.
- DeLorenzo, M.E., G.I. Scott, and P.E. Ross. 2001. "Toxicity of pesticides to aquatic microorganisms: A review." *Environ. Toxicol. Chem*. 20:84-98.
- DeMaster, D., S. Atkinson, and R. Dearborn. 2001. Is it food? II Workshop. Alaska Sea Life Center, Seward, Alaska. May 30-31. Sponsored by National Marine Fisheries Service and Alaska Sea Grant Office.
- Dennison, W.C. 1987. "Effect of light on seagrass photosynthesis, growth and depth distribution." *Aquatic Botany*. 27:15-26.
- Dolat, S.W. 1997. Acoustic measurements during the Baldwin Bridge demolition (final, dated March 14, 1997). Prepared for White Oak Construction by Sonalysts, Inc, Waterford, CT. 34 pp. + appendices.
- Duffy-Anderson, J.T., and K.W. Able. 1999. "Effects of municipal piers on the growth of juvenile fishes in the Hudson River estuary: a study across a pier edge." *Mar. Biol*. 133:409-418.
- Dugan, J.E., D.M. Hubbard, D.L. Martin, J.M. Engle, D.M. Richards, G.E. Davis, K.D. Lafferty, and R.F. Ambrose. 2000. Macrofauna communities of exposed sandy beaches on the Southern California mainland and Channel Islands. pp 339-346. In Brown, D.R., K.L. Mitchell, and H.W. Chang, eds. Proceedings of the Fifth California Islands Symposium. Minerals Management Service Publication # 99-0038.
- Ebeling, A.W., R.J. Larson, and W.S. Alevizon. 1980. Annual variability of reef-fish assemblages in kelp forest off Santa Barbara, California. U.S. Natl. Mar. Fish. Serv. Fish. Bull. 78:361-377.
- Enger, P.S., H.E. Karlsen, F.R. Knudsen, and O. Sand. 1993. Detection and reaction of fish to infrasound. Fish Behaviour in Relation to Fishing Operations., 1993, pp. 108-112, ICES marine science symposia. Copenhagen. Vol. 196.
- EPA (U.S. Environmental Protection Agency). 2002. National Water Quality Inventory: 2000 Report to Congress. EPA-841-R-02-001. EPA Office of Water, Washington, D.C.
- EPA. 2001. Reissuance of the NPDES General Permit for Seafood Processors Operating Throughout Alaska in Waters of the United States (NPDES General Permit No. AK-G52-0000) Federal Register: July 27, 2001. U.S. EPA, Region X.



- EPA. 2000. Environmental Screening Checklist and Workbook for the Water Transportation Industry. August 2000.
- EPA, Region 10. 2000. Authorization to discharge under the National Pollutant Discharge Elimination System (NPDES) for Section 402 modifications of Section 404 permits for log Transfer Facilities which received a Section 404 permit prior to October 22, 1985. NPDES Permit Number AK-G70-0000. March 2000. 1200 Sixth Avenue, OW-130 Seattle, Washington 98101.
- EPA, Region 10. 1996. Authorization to discharge under the National Pollutant Discharge Elimination System (NPDES) for Section 402 modifications of Section 404 permits for log Transfer Facilities which received a Section 404 permit prior to October 22, 1985. NPDES Permit Number AK-G70-1000. EPA Response to Comments from September 1996 Public Notice <http://info.dec.state.ak.us/DECPermit/water31rtc.pdf>.
- EPA. 1995. National Water Quality Inventory: 1994 Report to Congress. EPA-841-R-95-005. EPA Office of Water, Washington, D.C.
- EPA. 1993. Guidance for specifying management measures for sources of nonpoint pollution in coastal waters. EPA Office of Water. 840-B-92-002. 500+ pp.
- EPA. 1979. Impact of Seafood Cannery Waste on the Benthic Biota and Adjacent Waters at Dutch Harbor, Alaska.
- EPA. 1974. Development Document for Effluent Limitations Guidelines and Standards of Performance for the Catfish, Crab, Shrimp, and Tuna segments of the Canned and Preserved Seafood Processing Industry Point Source Category. Effluent Guidelines Division, Office of Water and Hazardous Material, Washington, D.C. EPA-44011-74-020-a. 389 pp.
- Evans, W.A., B. Johnston. 1980. Fish migration and fish passage: a practical guide to solving fish passage problems. Rev. ed. EM-7100-2. Washington DC. U.S. Department of Agriculture, Forest Service. 163 pp.
- Everest, F.H., R.L. Beschta, J.C. Scrivener, K.V. Koski, J.R. Sedell, and C.J. Cederholm. 1987. Fine sediment and salmonid production: a paradox. In: Salo, E.; Cundy, T. eds. Streamside management: forestry and fishery interactions: proceedings of a symposium held at the University of Washington, February 12-14, 1986. Contribution 57. Seattle: University of Washington, Institute of Forestry Resources. pp. 98-142.
- Ewald, G., P. Larsson, H. Linge, L. Okla, and N. Szarzi. 1998. "Biotransport of organic pollutants to an inland Alaska lake by migrating sockeye salmon (*Oncorhynchus nerka*)."  
*Arctic*. 51:40-47.
- Fajen, O.F. and J.B. Layzer. 1993. Agricultural practices. In Bryan, C.F. and D.A. Rutherford, eds. Impacts on warmwater streams: Guidelines for evaluation. pp. 257-267. Southern Division, American Fisheries Society. Little Rock, AR.
- Fawcett, J.A. and H.S. Marcus. 1991. Are port growth and coastal management compatible? Coastal Management. Vol. 19. pp. 275-295.
- Feder, H.M., C.H. Turner, and C. Limbaugh. 1974. Observations on fishes associated with kelp beds in southern California. California Department of Fish and Game, Fish Bull. 160. 144 pp.
- Ferraro, S.P., R.C. Swartz, F.A. Cole, and D.W. Schults. 1991. "Temporal changes in the benthos along a pollution gradient: discriminating the effects of natural phenomena from sewage-industrial wastewater effects." *Estuarine Coastal Shelf Science*. 33:383-407.
- Flanders, L.S. and J. Cariello. 2000. Tongass road condition survey report. Alaska Department of Fish and Game, Southeast Regional Office of the Habitat and Restoration Division. Alaska Department of Fish and Game, 802 3rd Street, Douglas, AK 99824. Technical Report 00-7. 48 pp.

- Foster, M.S. and D.R. Schiel. 1985. "The ecology of giant kelp forests in California: A community profile." *U.S. Fish Wildl. Serv. Biol. Rep.* 85(7.2) 152 pp.
- Franco, J., S. Schad, and C.W. Cady. 1994. "California's experience with a voluntary approach to reducing nitrate contamination of groundwater: The Fertilizer Research and Education Program (FREP)." *J. Soil and Water Conser.* 49: S76.
- Fulton, M.H., G.I. Scott, A. Fortner, T.F. Bidleman, and B. Ngabe. 1993. "The effects of urbanization on small high salinity estuaries of the southeastern United States." *Archives for Environmental Contamination and Toxicology.* 25(4):476-484.
- Furniss, M.J., T.D. Roelofs, and C.S. Yee. 1991. Road construction and maintenance. In Meehan W.R., ed. Influences of forest and rangeland management on salmonid fishes and their habitats. Special Publication 19. Bethesda, MD. American Fisheries Society. pp. 297-323.
- Glasby, T.M. 1999. "Effects of shading on subtidal epibiotic assemblages." *J. Exp. Mar. Biol. Ecol.* 234(1999). pp. 275-290.
- Good, J.W. 1987. "Mitigating estuarine development impacts in the Pacific Northwest: from concept to practice." *Northwest Environmental Journal.* Vol. 3. No. 1.
- Gowen, A.W. 1978. The Environmental Effects of OCS Pipelines. Initial Findings. New England River Basins Commission. Boston, MA. 4:24-43
- Greenberg, C.H., S.H. Crownover, and D.R. Gordon. 1997. "Roadside soils: a corridor for invasion of xeric scrub by nonindigenous plants." *Natural Areas Journal.* 17(2):99-109.
- Gregory, S.V. and P.A. Bisson. 1997. Degradation and loss of anadromous salmonid habitat in the Pacific Northwest. In Stouder, J.D., P.A. Bisson, and R.J. Naiman, eds. Pacific Salmon and Their Ecosystems: Status and Future Options, pp. 277-314. Chapman and Hall, New York.
- Grimes, C.B. 1975. "Entrapment of fishes on intake water screens at a steam electric generating station." *Chesapeake Science.* 16:172-177.
- Haas, M.A., C.A. Simenstad, J.R. Cordell, D.A. Beauchamp, and B.S. Miller. 2002. Effects of large overwater structures on epibenthic juvenile salmon prey assemblages in Puget Sound, Washington. Final Research Report No. WA-RD 550.1. Prepared for the Washington State Transportation Commission, Washington State Department of Transportation and the U.S. Department of Transportation, Federal Highway Administration. 114 pp.
- Hall, C., R. Howarth, B. Moore, and C. Vorosmarty. 1978. "Environmental impacts of industrial energy systems in the coastal zone." *Ann. Rev. Energy.* 3:395-475.
- Hammond, C.J., S.M. Miller, and R.W. Prellwitz. 1988. Estimating the probability of landslide failure using Monte Carlo simulation. In Proceedings of the 24th symposium on engineering geology and soils engineering. February 29, 1988. Coeur d'Alene, ID. Utah State University, Department of Civil and Environmental Engineering, Logan, UT. pp. 319-331.
- Hanrahan, L., C. Falk, H.A. Anderson, L. Draheim, M. S. Kanarek, J. Olson, and The Great Lakes Consortium. 1999. "Serum PCB and DDE levels of frequent Great Lakes sport fish consumers - a first look." *Env. Res. Section A.* 80:S26-S37.
- Hanson, C.H., J.R. White, and H.W. Li. 1977. "Entrapment and impingement of fishes by power plant cooling water intakes: an overview." *Marine Fisheries Review.* 39:7-17.
- Hastings, K., P. Hesp, and G. Kendrick. 1995. "Seagrass loss associated with boat moorings at Rottnest Island, Western Australia." *Ocean and Coastal Management.* 26:225-246. In Shafer, D.J. 2002. Recommendations to minimize potential impacts to seagrasses from single-family residential dock structures in the Pacific Northwest. U.S. Army Corps of Engineers, Seattle, WA. 28 pp.

- Hastings, M.C. 2002. Clarification of the meaning of sound pressure levels and the known effects of sound on fish. Document in support of Biological Assessment for San Francisco-Oakland Bay Bridge East Span Seismic Safety Project. August 26, 2002. Revised August 27, 2002. 8 pp.
- Heady, H.F. and R.D. Child. 1994. Rangeland ecology and management. Westview Press, Inc., Boulder, Colorado.
- Heifetz, J., M.L. Murphy, and K.V. Koski. 1986. "Effects of logging on winter habitat of juvenile salmonids in Alaskan streams." *North American Journal of Fisheries Management*. 6:52-58.
- Heintz, R.A., S.D. Rice, A.C. Wertheimer, R.F. Bradshaw, F.P. Thrower, J.E. Joyce, and J.W. Short. 2000. "Delayed effects on growth and marine survival of pink salmon *Oncorhynchus gorbuscha*, after exposure to crude oil during embryonic development." *Mar. Ecol. Prog. Ser.* 208:205-216.
- Helfman, G.S. 1981. The advantage to fishes of hovering in shade. *Copeia*. 1981(2):392-400.
- Helvey, M. 1985. "Behavioral factors influencing fish entrapment at offshore cooling-water intake structures in southern California." *Marine Fisheries Review*. 47:18-26.
- Helvey, M. 2002. "Are southern California oil and gas platforms essential fish habitat?" *ICES Journal of Marine Science*. 59:S266-S271.
- Helvey, M. and P.B. Dorn. 1987. "Selective removal of reef fish associated with an offshore cooling-water intake structure." *J. Applied Ecology* 24:1-12.
- Herke, W.H. and B.D. Rogers. 1993. Maintenance of the estuarine environment. Pages 263-286 in C.C. Kohler and W.A. Hubert, editors. Inland fisheries management in North America. American Fisheries Society, Bethesda, Maryland.
- Hicks, B.J., J.D. Hall, P.A. Bisson, and J.R. Sedell, J.R. 1991. Responses of salmonids to habitat changes. In: Meehan, W.R. (Ed.) 1991. Influences of forest and rangeland management on salmonid fishes and their habitats. Special Publication 19. Bethesda, MD: American Fisheries Society: 483-518.
- Hoagland, K.D., J.P. Carder, and R.L. Spawn. 1996. Effects of organic toxic substances. pp. 469-496. In Stevenson, R.J., M.L. Bothwell, and R.L. Lowe, eds. *Algal Ecology: Freshwater Benthic Ecosystems*. Academic Press. New York, NY.
- Holler, J.D. 1990. "Nonpoint source phosphorus control by a combination wet detention/filtration facility in Kissimmee, Florida." *Florida Scientist*. 53(1):28-37.
- Hoss, D.E. and G.W. Thayer. 1993. The importance of habitat to the early life history of estuarine dependent fishes. American Fisheries Society Symposium. 14:147-158.
- Hurme, A.K. and E.J. Pullen. 1988. Biological effects of marine sand mining and fill placement for beach replenishment: Lesson for other use. *Marine Mining*. Vol. 7.
- Ikonomou, M.G., S. Rayne, and R.F. Addison. 2002. "Exponential increases of the brominated flame retardants, polybrominated diphenyl ethers, in the Canadian Arctic from 1981 to 2000." *Environ. Sci. Technol.* 36:1886-1892.
- Jackson, R.G. 1986. "Effects of bark accumulation on benthic infaunal at a log transfer facility in Southeast Alaska." *Marine Pollution Bulletin*. 17(6):258-262.
- Johnson, K.L. 1992. Management for water quality on rangelands through best management practices: the Idaho approach. pp. 415-441. In Naiman, R.J., ed. *Watershed management: Balancing sustainability and environmental change*. Springer-Verlag, New York.

- Johnson, L. 2000. An analysis in support of sediment quality thresholds for polycyclic aromatic hydrocarbons (PAHs) to protect estuarine fish. White Paper from National Marine Fisheries Service, Northwest Fisheries Science Center, Seattle, Washington. 29 p.
- Johnson, L., S.Y. Sol, G.M. Ylitalo, T. Hom, B. French, O.P. Olson, and T.K. Collier. 1999. "Reproductive injury in English sole (*Pleuronectes vetulus*) from the Hylebos Waterway, Commencement Bay, Washington." *Journal of Aquatic Ecosystem Stress and Recovery*. 6:289-310.
- Johnson, S.W., S.D. Rice, and D.A. Moles. 1998a. Effects of submarine mine tailings disposal on juvenile yellowfin sole (*Pleuronectes asper*): a laboratory study. *Marine Pollution Bulletin*. 36:278-287.
- Johnson, S.W., R.P. Stone, and D.C. Love. 1998b. Avoidance behavior of ovigerous Tanner crabs (*Chionoecetes bairdi*) exposed to mine tailings: a laboratory study. *Alaska Fish. Res. Bull.* 5:39-45.
- Jordan, T.E. and D.E. Weller. 1996. "Human contributions to terrestrial nitrogen flux: Assessing the sources and fates of anthropogenic fixed nitrogen." *BioScience*. 46:655.
- Kagan, R.A. 1991. The dredging dilemma: economic development and environmental protection in Oakland Harbor. *Coastal Management*, Vol. 19, pp. 313-341.
- Kahler, T., M. Grassley, and D. Beauchamp. 2000. A summary of the effects of bulkheads, piers, and other artificial structures and shorezone development on ESA listed salmonids in lakes. Final Report to the City of Bellevue, Washington. 74 pp.
- Kauffman, J.B. and W.C. Krueger. 1984. "Livestock impacts on riparian ecosystems and streamside management implications—a review." *Journal of Range Management*. 37:430-438.
- Kawano, M., S. Matsushita, T. Inoue, H. Tanaka, and R. Tatsukawa. 1986. "Biological accumulation of chlordanes compounds in marine organisms from the northern North Pacific and Bering Sea." *Mar. Poll. Bull.* 17:512-516.
- Kennish, M.J. 1998. *Pollution Impacts on Marine Biotic Communities*. CRC Press, New York, NY. 310 pp.
- Kirkpatrick, B., T.C. Shirley, and C.E. O'Clair. 1998. Deep-water bark accumulation and benthos richness at log transfer and storage facilities. *Alaska Fishery Research Bulletin* 5(2):103-115.
- Klein, R. 1997. The effects of marinas and boating activities upon tidal waters. Community and Environmental Defense Services. Owings Mills, Maryland. 23 pp.
- Knudsen, F.R., C.B. Schreck, S.M. Knapp, P.S. Enger, and O. Sand. 1997. "Infrasound produces flight and avoidance responses in Pacific juvenile salmonids." *Journal of Fish Biology*. 51:824-829.
- Knudsen, F.R., P.S. Enger, and O. Sand. 1994. "Avoidance responses to low frequency sound in downstream migrating Atlantic salmon smolt, *Salmo salar*." *Journal of Fish Biology*. 45:227-233.
- Kohler, C.C. and W.R. Courtenay, Jr. 1986. "Introduction of aquatic species." *Fisheries*. 11(2):39-42. Proceedings of the Seventh International Zebra Mussel and Aquatic Nuisance Species Conference. 1997.
- Koski, K.V. 1981. The survival and quality of two stocks of chum salmon (*Oncorhynchus keta*) from egg deposition to emergence. *Rapp. P.-v. Reun. Conx. int. Explor. Mer*, 178:330-333.
- Langford, T.E., N.J. Utting, and R.H.A. Holmes. 1978. Factors affecting the impingement of fishes on power station cooling-water intake screens. *In Physiology and Behaviour of Marine Organisms*. D.S. McLusky and A.J. Berry, eds. pp. 281-288. Pergamon Press, Oxford and New York.

- Larimore, R.W. and P.W. Smith. 1963. The fishes of Champaign County, Illinois, as affected by 60 years of stream changes. *Illinois Natural History Survey Bulletin* 28:299-382.
- Larsen, M.C. and J.E. Parks. 1997. How wide is a road? The association of roads and mass-wasting in a forested montane environment. *Earth Surface Processes and Landforms* 22:835-848.
- Larson, K. and C. Moehl. 1990. Entrainment of anadromous fish by hopper dredge at the mouth of the Columbia River. In C.A. Simenstad, ed. *Effects of dredging on anadromous Pacific coast fishes*. University of Washington Sea Grant. pp. 102-112.
- Lee, G.F. and S. Taylor. 2000. Review of the water quality characteristics of Upper Newport Bay, Orange County CA and its Tributaries. *California Coastal Conservancy, Oakland, CA* 94612.
- Longmuir, C. and T. Lively. 2001. Bubble curtain systems for use during marine pile driving. Report by Fraser River Pile & Dredge Ltd., New Westminster, British Columbia. 9 pp.
- Lonsdale, W.N. and A.M. Lane. 1994. "Tourist vehicles as vectors of weed seeds in Dadoed National Park, northern Australia." *Biological Conservation*. 69(3):277-283.
- Love, M.S. and W. Westphal. 1990. Comparison of fishes taken by a sportfishing party vessel around oil platforms and adjacent natural reefs near Santa Barbara, California. *Fishery Bulletin, U.S.* 88:599-605.
- Love, M., J. Hyland, A. Egeling, T. Herrlinger, A. Brooks, and E. Imamura. 1994. "A pilot study of the distribution and abundance of rockfishes in relation to natural environmental factors and an offshore oil and gas production platform off the coast of southern California." *Bulletin Marine Science*. 55(2-3): 1062-1085.
- Love, M.S., M. Nishimoto, D. Schroeder, and J. Caselle. 1999. The ecological role of natural reefs and oil and gas production platforms on rocky reef fishes in southern California: Final interim report. U.S. Geological Survey, Biological Resources Division. USGS/BRD/CR-1999-007. 208 pp.
- Luce, A. and M. Crowe. 2001. "Invertebrate terrestrial diversity along a gravel road on Barrie Island, Ontario, Canada." *Great Lakes Entomologist*. 34(1):55-60 SPR-SUM.
- MacDonald, L.H., R.W. Sampson, and D.M. Anderson. 2001. "Runoff and road erosion at the plot and road segment scales, St. John, U.S. Virgin Islands." *Earth Surface Processes and Landforms*. 26:251-272.
- Madej, M.A. 2001. "Erosion and sediment delivery following removal of forest roads." *Earth Surface Processes and Landforms*. 26:175-190.
- Marcus, M.D., M.K. Young, L.E. Noel, and B.A. Mullan. 1990. Salmonid-habitat relationships in the western United States: a review and indexed bibliography. General Technical Report RM-GTR-188. U.S. Department of Agriculture, Forest Service, Rocky Mountain Forest and Range Experiment Station, Fort Collins, CO.
- Maser, C. and J.R. Sedell. 1994. *From the Forest to the Sea: the Ecology of Wood in Streams, Estuaries and Oceans*. St. Lucie Press, Delray Beach, FL. 200 pp.
- McGraw, K. and D. Armstrong. 1990. Fish entrainment by dredges in Grays Harbor, Washington. In C.A. Simenstad, ed. *Effects of dredging on anadromous Pacific coast fishes*. University of Washington Sea Grant. pp. 113-131.
- Meehan, W.R., W.A. Farr, D.M. Bishop, and J.H. Patric. 1969. Some effects of clearcutting on salmon habitat of two southeast Alaska streams. U.S. Department of Agriculture, Forest Service. General Technical Report PNW-82. Institute of Northern Forests, Pacific Northwest Forest and Range Experiment Station.

- Michel, J. and M.O. Hayes. 1999. "Weathering patterns of oil residues eight years after the Exxon Valdez oil spill." *Marine Pollution Bulletin*. 38(10):855-863.
- Miller, D.J. and J.J. Geibel. 1973. Summary of blue rockfish and lingcod life histories; A reef ecology study; and giant kelp, *Macrocystis pyrifera*, experiments in Monterey Bay, California. California Dept. of fish and Game, Fish Bull. 158.
- Moazzam, M. and S.H.N. Rizvi. 1980. "Fish entrapment in the seawater intake of a power plant at Karachi coast." *Environ. Biology of Fishes*. 5:49-57.
- Moles, A. and N. Hale. *in press*. Use of physiological responses in *Mytilus trossulus* as integrative bioindicators of sewage pollution. *Marine Pollution Bulletin*.
- Montgomery, D.R. 1994. "Road surface drainage, channel initiation, and slope instability." *Water Resources Research*. 30:1925-1932.
- Montgomery, D.R., R.D. Smith, K.M. Schmidt, and G.R. Pess. 1995. "Pool Spacing in Forest Channels." *Water Resources Research*. 31:1097-1105.
- Moore, A. and Waring, C.P. 2001. "The effects of a synthetic pyrethroid pesticide on some aspects of reproduction in Atlantic salmon (*Salmo salar* L)." *Aquatic Toxicology*. 52:1-12.
- Muir, D., B. Braune, B. DeMarch, R. Norstrom, R. Wagemann, L. Lockhart, B. Hargrave, D. Bright, R. Addison, J. Payne, and K. Reimer. 1999. "Spatial and temporal trends and effects of contaminants in the Canadian Arctic marine ecosystem: a review." *Sci Total Environ*. 230:83-144.
- Muir, D.C.G. and R.J. Norstrom. 2000. "Geographical differences and time trends of persistent organic pollutants in the Arctic." *Toxicol. Lett*. 112-113:93-101.
- Murphy, M.L., S.W. Johnson, and D.J. Csepp. 2000. "A comparison of fish assemblages in eelgrass and adjacent subtidal habitats near Craig, Alaska." *AK. Fish. Res. J.* 7:11-21.
- Myren, R.T. and R.J. Ellis. 1984. Evapotranspiration in forest succession and long-term effects upon fishery resources: a consideration for management of our old-growth forests. pp. 183-186. *In* Meehan, M.R., T.R. Merrill, Jr., and T.A. Hanley, eds. Fish and wildlife relationships in old growth forests. American Institute of Fishery Research Biologists.
- Neff, J.M. 1985. Polycyclic aromatic hydrocarbons. *In* G.M. Rand and S.R. Petrocelli. Fundamentals of aquatic toxicology. pp. 416-454. Hemisphere Publishing, Washington, D.C.
- Nelson, R.L., M. McHenry, and W.S. Platts. 1991. Mining. Influences of forest and rangeland management in salmonid fishes and their habitats. pp. 425-457. *In*: Meehan, W. ed. Influences of forest and range management on salmonid fishes and their habitats. AFS Special Publication 19. Bethesda, MD.
- Newell, R.C., L.J. Seiderer, and D.R. Hitchcock. 1998. "The impact of dredging on biological resources of the sea bed." *Oceanography and Marine Biology Annual Review*. 36:127-178.
- Nightingale, B. and C.A. Simenstad. 2001a. Dredging activities: Marine issues. Washington State Transportation Center, University of Washington, Seattle, WA 98105. (Document available through the National Technical Information Service, Springfield, VA 22616).
- Nightingale, B. and C.A. Simenstad. 2001b. Overwater Structures: Marine Issues. White paper submitted to Washington Department of Fish and Wildlife, Washington Department of Ecology and Washington Department of Transportation. [www.wa.gov/wdfw/hab](http://www.wa.gov/wdfw/hab). 133 pp.
- NMFS (National Marine Fisheries Service). 2003. Biological Opinion for the Benicia-Martinez New Bridge Project, Southwest Region, Santa Rosa, California. Admin. Rec. 151422SWR02SR6292.

- NMFS. 2002. Environmental Assessment, NMFS' Restoration Plan for the Community-Based Restoration Program. Prepared by the NOAA Restoration Center, Office of Habitat Conservation. Silver Spring, MD.
- NMFS. 2001. Biological Opinion for the San Francisco-Oakland Bay Bridge East Span Seismic Safety Project. Southwest Region, Santa Rosa, California. Admin. Rec. 151422SWR99SR190.
- NMFS. 1998. Final recommendations: Essential Fish Habitat for Pacific Coast Groundfish. Prepared by: The Core Team for EFH for Pacific Coast Groundfish June 3, 1998. 2725 Montlake Blvd. E. Seattle, WA 98112. [http://www.psmfc.org/efh/groundfish\\_desc.pdf](http://www.psmfc.org/efh/groundfish_desc.pdf).
- NMFS. August 29, 1996. National Gravel Extraction Policy. 1335 East-West Highway, Silver Spring, MD 20910. (<http://swr.nmfs.noaa.gov/hcd/gravel.pdf>).
- NPPC (Northwest Power Planning Council). 1986. Compilation of information on salmon and steelhead losses in the Columbia River Basin. Columbia River Basin and Wildlife Program. Portland, OR.
- NRC (National Research Council), Committee on Hardrock Mining. 1999. Hardrock Mining on Federal Lands. Appendix B. Potential Environmental Impacts of Hardrock Mining. ([http://www.nap.edu/html/hardrock\\_fed\\_land/appB.html](http://www.nap.edu/html/hardrock_fed_land/appB.html)).
- NRC. 1996. Upstream: salmon and society in the Pacific Northwest. Report of the committee on protection and management of Pacific Northwest anadromous salmonids. Board on Environmental Studies and Toxicology, and Commission on Life Sciences. National Academy Press. Washington, D.C. 20055.
- NRC. 1989. Irrigation-induced water quality problems: what can be learned from the San Joaquin Valley experience. National Academy Press, Washington, D.C.
- O'Hara, T. 2001. *Evaluating Environmental Contaminant Trends in Arctic Alaska Marine Mammals: Biological and Experimental Design Considerations*. Workshop to Assess Contaminant Impacts on Steller Sea Lions in Alaska. Anchorage, AK, September 5-6, 2001. Sponsored by Auke Bay Laboratory.
- OGTAD (Oil and Gas Technologies for the Arctic and Deepwater). 1985. U.S. Congress, Office of Technology Assessment, OTA-O-270, May 1985. Library of Congress Catalog Card Number 85-600528. U.S. Government Printing Office, Washington, DC 20402.
- OWRRI (Oregon Water Resource Research Institute). 1995. Gravel disturbance impacts on salmon habitat and stream health, volume 1. summary report. Oregon State University, Corvallis, Oregon. (Also available Vol. II: Technical background report). Available from Oregon Division of State Lands, Salem, Oregon, 503-378-3805.
- Pezeshki, S.R., R.D. Delaune, and W.H. Patrick, Jr. 1987. "Response of the freshwater marsh species, *Panicum hemitomon* Schult., to increased salinity." *Freshwater Biology*. 17:195-200
- PFMC (Pacific Fisheries Management Council). 2003. Essential fish habitat: West coast groundfish. Draft revised appendix. Updated January, 2003. Bruce McCain (ed.). Portland, OR.
- PFMC. 1998. Essential fish habitat: Coastal pelagic species. Coastal Pelagics Species Fishery Management Plan. Amendment 8 to the Northern Anchovy Fishery Management Plan. Portland, OR.
- PFMC. 1999. Appendix A: Identification and description of essential fish habitat, adverse impacts, and recommended conservation measures for salmon. Amendment 14 to the Pacific Coast Salmon Plan. Portland, OR. 146 pp.
- Phillips, R.C. 1984. The ecology of eelgrass meadows in the Pacific Northwest: a community profile. U.S. Fish and Wildlife Service. FWS/OBS-84/24. 85 pp.

- Platts, W.S. 1991. Livestock grazing. pp. 389-424. *In* Meehan, W.R, ed. Influences of forest and rangeland management on salmonid fishes and their habitats. Special Publication 19. American Fisheries Society, Bethesda, MD.
- PNPCC (Pacific Northwest Pollution Control Council). 1971. Log storage and rafting in public waters. Task Force report. p. 56.
- Polasek, L.G. 1997. Assessment of wetland habitat alterations resulting from construction of a pipeline through coastal marshes in Orange County, Texas. Final Report: Texas Parks and Wildlife Department, Port Arthur, TX. 40 pp.
- Poston, T. 2001. Treated wood issues associated with overwater structures in marine and freshwater environments. White Paper submitted to Washington Department of Fish and Wildlife, Washington Department of Ecology and Washington Department of Transportation by Batelle. 85 pp.
- Preston, B.L. 2002. "Indirect effects in aquatic ecotoxicology: implications for ecological risk assessment." *Environmental Management*. 29:311-323.
- Raco-Rands, V.E. 1996. Characteristics of effluents from power generating stations in 1994. *In* Allen, M.J, ed. Southern California Coastal Water Research Project, Annual Report 1994-95. SCCWRP, Westminster, CA. pp. 29-36.
- Ralph, S., G. Poole, L. Conquest, and R. Naiman. 1994. "Stream channel morphology and woody debris in logged and unlogged basins in western Washington." *Can. J. Fish. Aquatic Sciences*. 51:37-51.
- Raymond, H. 1979. "Effects of dams and impoundment on migrations of juvenile chinook salmon and steelhead from the Snake River, 1966 to 1975." *Trans. Amer. Fish. Soc.* 108:505-529.
- Reinjders, P.J.H. 1986. "Reproductive failure in common seals feeding on fish from polluted coastal waters." *Nature*. 324:456-457
- Reiser and Bjornn. 1979. Influence of forest and rangeland management on anadromous fish habitat in the Western United States and Canada. Meehan, W.R., ed. U.S. Department of Agriculture, Forest Service. General Technical Report PNW-96. U.S. Department of Agriculture, Forest Service, Pacific Northwest Forest and Range Experiment Station. Portland, OR.
- Reyff, J.A and P. Donovan. 2003. Benicia-Martinez Bridge Bubble Curtain Test - Underwater Sound Measurement Data. Memo to Caltrans dated January 31, 2003. 3 pp.
- Reyff, J.A. 2003. Underwater sound levels associated with seismic retrofit construction of the Richmond-San Rafael Bridge. Document in support of Biological Assessment for the Richmond-San Rafael Bridge Seismic Safety Project. January 31, 2003. 18 pp.
- Rice, S.D., J.W. Short, R.A. Heintz, M.G. Carls, and A. Moles. 2000. Life-history consequences of oil pollution in fish natal habitat. pp. 1210-1215. *In* Catania, P., ed. *Energy 2000*. Balaban Publishers, Lancaster, England.
- Richardson, W.J., C.R. Greene, C.I. Malme, and D.H. Thomson. 1995. Marine Mammals and Noise. Academic Press, San Diego. 576 p.
- Riedman, M. 1990. *The Pinnipeds*. University of California Press. Berkeley, California. 439 pp.
- Rogers, P.H. and M. Cox. 1988. Underwater sound as a biological stimulus. pp. 131-149. *In* Sensory biology of aquatic animals. Atema, J, R.R. Fay, A.N. Popper, and W.N. Tavolga, eds. Springer-Verlag. New York.



- Ruus, A., K.I. Ugland, and J.U. Skaare. 2002. "Influence of trophic position on organochlorine concentrations and compositional patterns in a marine food web." *Env. Tox. Chem.* 21(11):2356-2364.
- Safavi, H.R. 1996. Quality control of urban runoff and sound management. *Hydrobiologia: Diapause in the Crustacea.* pp. 131 -141.
- SAIC (Science Applications International Corporation). 2001. Information Collection Request for National Pollutant Discharge Elimination System (NPDES) and Sewage Sludge Monitoring Reports. Prepared by SAIC, 11251 Roger Bacon Drive, Reston, VA 20190, for Tetra Tech, Inc., Fairfax, VA, for the U.S. Environmental Protection Agency, Office of Wastewater Management, Washington, D.C. EPA ICR# 0229.15. p. 11.
- Sand, O., P.S. Enger, H.E. Karlsen, F. Knudsen, T. Kvernstuen. 2000. "Avoidance responses to infrasound in downstream migrating European silver eels, *Anguilla anguilla*." *Environmental Biology of Fishes.* 57:327-336.
- Scholz, N.L., N.K. Truelove, B.L. French, B.A. Berejikian, T.P. Quinn, E. Casillas, and T.K. Collier. 2000. "Diazinon disrupts antipredator and homing behaviors in chinook salmon (*Oncorhynchus tsawytscha*)." *Canadian Journal of Fisheries and Aquatic Sciences.* 57:1911-1918.
- Scott G.I., M.H. Fulton, D.W. Moore, E.F. Wirth, G.T. Chandler, P.B. Key, J.W. Daugomah, E.D. Strozier, J. Devane, J.R. Clark, M.A. Lewis, D.B. Finley, W. Ellenberg, and K.J. Karnaky. 1999. "Assessment of risk reduction strategies for the management of agricultural nonpoint source pesticide runoff in estuarine ecosystems." *Toxicology and Industrial Health.* 15:200-213.
- Scrivener, J.T. and M.J. Brownlee. 1989. "Effects of forest harvesting on spawning gravel and incubation survival of chum (*Oncorhynchus kata*) and coho (*O. kisutch*) salmon in Carnation Creek, British Columbia." *Canadian Journal of Fisheries and Aquatic Sciences.* 46(4):681-696.
- SCS (Soil Conservation Service), U.S. Department of Agriculture, Forest Service, and Economic Research Service. 1984. Southeast Washington cooperative river basin study. U.S. Department of Agriculture, Washington, D.C.
- Sengupta, M. 1993. Environmental Impacts of Mining: Monitoring, Restoration, and Control. CRC Press, Inc. 2000 Corporate Blvd., N.W. Boca Raton, FL. 33431. p.1.
- Shafer, D.J. 2002. Recommendations to minimize potential impacts to seagrasses from single-family residential dock structures in the Pacific Northwest. U.S. Army Corps of Engineers, Seattle, WA. 28 p.
- Short, J.W., M.R. Lindeberg, P.M. Harris, J. Maselko, and S.D. Rice. 2002. Vertical Oil Distribution Within the Intertidal Zone 12 Years After the *EXXON VALDEZ* Oil Spill in Prince William Sound, Alaska. Pp. 57-72 In: Proceedings of the Twenty-fifth Arctic and Marine Oilspill Program (AMOP) Technical Seminar. Environment Canada, Ottawa, Ontario.
- Sidle, R.C., A.J. Pearce, and C.L. O'Loughlin. 1985. Hillslope stability and land use. Water Resources Monograph 11. Washington D.C.: American Geophysical Union 140 pp.
- Simenstad, C.A., C.D. Tanner, F. Weinmann, and M. Rylko. 1991. The estuarine habitat assessment protocol. Puget Sound Notes. No. 25. June 1991.
- Sogard, S.M. and K.W. Able. 1991. "A comparison of eelgrass, sea lettuce macroalgae and marsh creeks as habitats for epibenthic fishes and decapods." *Estuarine, Coastal and Shelf Science.* 33, 501-519.
- Spence, B.C., G.A. Lomnický, R.M. Hughes, and R.P. Novitzki. 1996. An ecosystem approach to salmonid conservation. TR-4501-96-6057. ManTech Environmental Research Services Corp., Corvallis, OR. 356 pp. (Available from the NMFS Habitat Branch, Portland, OR).

- Stadler, J.H. 2002. Personal observation of fish-kill occurring during pile driving activity at the Winslow Ferry Terminal, Winslow, WA. October 7, 2002. Fish Biologist, DOC/NOAA/National Marine Fisheries Service/HCD, Lacey, WA.
- Stehr, C.M., D.W. Brown, T. Hom, B.F. Anulacion, W.L. Reichert, and T.K. Collier. 2000. "Exposure of juvenile chinook and chum salmon to chemical contaminants in the Hylebos Waterway of Commencement Bay, Tacoma, Washington." *Journal of Aquatic Ecosystem Stress and Recovery*. 7:215-227.
- Stein, J., T. Hom, T. Collier, D. Brown, and U. Varanasi. 1995. "Contaminant exposure and biochemical effects in outmigrant juvenile chinook salmon from urban estuaries of Puget Sound, WA." *Environ. Toxicol. Chem.* 14:1019-1029.
- Stewart, R K. and D.R. Tangarone. 1977. Water Quality Investigations Related to Seafood Processing Wastewater Discharges at Dutch Harbor, Alaska - October 1975 and October 1976. Region X, U.S. Environmental Protection Agency. Working Paper #EPA 910/8-77-100. 78 pp.
- Stotz, T. and J. Colby. 2001. January 2001 dive report for Mukilteo wingwall replacement project. Washington State Ferries Memorandum. 5 pp. + appendices.
- Swanston, D.N. 1974. The forest ecosystem of southeast Alaska. Soil mass movement. General Technical Report PNW-17. U.S. Department of Agriculture, Forest Service. Pacific Northwest Forest and Range Experiment Station. Portland, OR.
- Thayer, G.W., W.J. Kenworthy, and M.S. Fonseca. 1984. The ecology of eelgrass meadows of the Atlantic coast: a community profile. U.S. Fish and Wildlife Service FWS/OBS-84/02. 147 pp.
- Travnichek, V.H., A.V. Zale, and W.L. Fisher. 1993. "Entrainment of ichthyoplankton by a warmwater hydroelectric facility." *Trans. Amer. Fish. Soc.* 122:709-716.
- Trombulak, S.C. and C.A. Frissell. 2000. "Review of ecological effects of roads on terrestrial and aquatic communities." *Conservation Biology*. 14(10):18-30. February.
- USACE (U.S. Army Corps of Engineers). 1993. Engineering and Design: Environmental; Engineering for Small Boat Basins. EM 1110-2-1206. Dept. of the Army, CECW-EH-W. Washington DC.
- USFWS (U.S. Fish and Wildlife Service). 1980. FWS/OBS-80/09. Gravel Removal Guidelines Manual for Arctic and Subarctic Floodplains. June 1980. Prepared by Woodward Clyde Consultants. Contract Number FWS-14-16-0008-970. Performed for the Water Resources Analysis Project, Office of Biological Services. U.S. Department of the Interior, Washington, D.C. 20240.
- USGS (U.S. Geologic Survey). 1999. The quality of our nation's waters: Nutrients and pesticides. pp. 33-55. USGS, Reston, VA.
- Van der Veer, H., M.J.N. Bergmen, and J.J. Beukema. 1985. "Dredging activities in the Dutch Wadden Sea effects on macrobenthic infauna." *Netherlands Journal for Sea Research*. 19:183-190.
- Van Dolah, R.E., P.P. Maier, M.H. Fulton, and G.I. Scott. 1997. "Comparison of azinphomethyl toxicity to juvenile red drum (*Scianops ocellatus*) and the mummichog (*Fundulus heteroclitus*)." *Environmental Toxicology and Chemistry*. 16:1488-1493.
- Vetter, E.W. 1995. "Detritus based patches of high secondary production in the nearshore benthos." *Mar. Ecol. Prog. Ser.* 120:251-262.
- Vuorinen, P.J., R. Parmanne, T. Vartiainen, M. Keinanen, H. Kiviranta, O. Kotovuori, and F. Hallig. 2002. "PCDD, PCF, PCB and thiamin in Baltic herring (*Clupea harengus* L.) and sprat (*Sprattus sprattus* L.) as a background to the M74 syndrome of Baltic salmon (*Salmo salar* L.)." *ICES J. Mar. Sci.* 59:480-496.

- Waisley, S.L. 1998. Projections for U.S. and Global Supply and Demand for 2010 and 2020. presented at U.S. and China Oil and Gas Industrial Forum, Beijing, People's Republic of China, November 2-4, 1998. Office of Natural Gas and Petroleum Technology, U.S. DOE, Washington, D.C. ([http://www.fe.doe.gov/oil\\_gas/china\\_forum/cl04000.html](http://www.fe.doe.gov/oil_gas/china_forum/cl04000.html)).
- Waldichuk, M. 1993. Fish habitat and the impact of human activity with particular reference to Pacific salmon. pp. 295-337. In Parsons, L.S. and W.H. Lear, eds. Perspectives on Canadian marine fisheries management. Canadian Bulletin of Fisheries and Aquatic Sciences 226.
- Walker, D., R. Lukatelich, R.G. Bastyan and A.J. McComb. 1989. "The effect of boat moorings on seagrass beds near Perth, Western Australia." *Aquatic Botany*. 36:69-77. In Shafer, D.J. 2002. Recommendations to minimize potential impacts to seagrasses from single-family residential dock structures in the Pacific Northwest. U.S. Army Corps of Engineers, Seattle, WA. 28 pp.
- Wania, F. and D. Mackay. 1999. "Global chemical fate of hexachlorocyclohexane. 2. Use of a global distribution model for mass balancing, source apportionment, and trend prediction." *Environ Toxicol Chem*. 18: 1400B1407.
- Warrington, P.D. 1999. Impacts of outboard motors on the aquatic environment. <http://www.nalms.org/bclss/impactsoutboard.htm>.
- WDFW (Washington State Department of Wildlife). 1998. Gold and fish. Rules and regulations for mineral prospecting and mining in Washington State. Draft, February 1998. Olympia, WA.
- Weaver, T.M. and J.J. Fraley. 1993. "A method to measure emergence success of westslope cutthroat trout fry from varying substrate compositions in a natural stream channel." *North American Journal of Fisheries Management*. 13(4):817-822.
- Wemple, B.C., F.J. Swanson, and J.A. Jones. 2001. "Forest roads and geomorphic process interactions, Cascade Range, Oregon." *Earth Surface Processes and Landforms*. 26:191-204.
- Wemple, B.C., J.A. Jones, and G.E. Grant. 1996. "Channel network extension by logging roads in two basins, western Cascades, Oregon." *Water Resources Bulletin*. 32:1195-1207.
- Wertheimer, A.C., R.A. Heintz, J.F. Thedinga, J.M. Maselko, and S.D. Rice. 2000. Straying of adult pink salmon from their natal stream following exposure as embryos to weathered Exxon Valdez crude oil. *Trans. Amer. Fish. Soc.* 129: 989-1004.
- West, C., L. Galloway, and J. Lyon. 1995. Mines, Stormwater Pollution, and You. Mineral Policy Center, Washington, D.C.
- Williams, G.D. and R.M. Thom. 2001. Marine and estuarine shoreline modification issues. White paper submitted to Washington Department of Fish and Wildlife, Washington Department of Ecology and Washington Department of Transportation. [www.wa.gov/wdfw/hab/ahg](http://www.wa.gov/wdfw/hab/ahg). 99 pp.
- Würsig, B., C.R. Greene, Jr., and T.A. Jefferson. 2000. "Development of an air bubble curtain to reduce underwater noise of percussive pile driving." *Marine Environmental Research*. 49:79-93.
- Wyllie-Echeverria, S. and R.C. Phillips. 1994. pp. 1-4. In Wyllie-Echeverria, S., A.M. Olson and M.J. Hershman, eds. Seagrass science and policy in the Pacific Northwest: proceedings of a seminar series (SMA 94-1) EPA 910/R-94-004. 63 pp.
- Yang, J., D. Shin, S. Park, Y. Chang, D. Kim, and M.G. Ikonomou. 2002. "PCDDs, PCDFs and PCBs concentrations in breast milk from two areas in Korea: body burden of mothers and implications for feeding infants." *Chemosphere*. 46(3):419-28.
- Yelverton, J.T., D.R. Richmond, W. Hicks, K. Saunders, and R. Fletcher. 1975. The relationship between fish size and their response to underwater blast. Lovelace Foundation for Medical Education and Research, Albuquerque, NM.

- Ylitalo, G.M. C.O. Matkin, J. Buzitis, M.M. Krahn, L.L. Jones, T. Rowles, and J.E. Stein. 2001a. "Influence of life-history parameters on organochlorine concentrations in free-ranging killer whales (*Orcinus orca*) from Prince William Sound, AK." *Sci. Tot. Env.* 281:183-203.
- Ylitalo, G.M., J.W. Bickham, J. Buzitis, G.K. Yanagida, J.E. Stein, and M.M. Krahn. 2001b. *Contaminant Bioaccumulation and Feeding ecology of Steller Sea Lions from Alaska*. Workshop to Assess Contaminant Impacts on Steller Sea Lions in Alaska. Anchorage, AK, September 5-6, 2001. Sponsored by Auke Bay Laboratory.
- York, A.E., R.L. Merrick, and T.R. Loughlin. 1996. An analysis of the Steller sea lion metapopulation in Alaska. In McCullough, D., ed. *Metapopulations and wildlife conservation and management*. Island Press, Covelo, CA. pp. 259-292.
- Young, M.K., W.A. Hubert, and T.A. Wesche. 1991. "Selection of measures of substrate composition to estimate survival to emergence of salmonids and to detect changes in stream substrates." *North American Journal of Fisheries Management*. 11 (3):339-346.
- Zale, A.V., O.E. Maughan, D.J. Orth, and W. Layher. 1993. Withdrawals. In Bryan, C.F and D.A. Rutherford, eds. *Impacts on warmwater streams: Guidelines for evaluation*. Pp. 271-281. Southern Division, American Fisheries Society, Little Rock, AR.
- Zedler, J.B., C.S. Nordby, and B.E. Kus. 1992. The ecology of the Tijuana Estuary, California: A National Estuarine and Research Reserve. NOAA Office of Coastal Resource Management, Sanctuaries and Reserves Division, Washington, D.C.
- Zenteno-Savin, T., M. Castellini, L.D. Rea, and B.S. Fadely. 1997. "Plasma haptoglobin in threatened Alaskan pinniped populations." *J. Wildlife Dis.* 33:64-71.
- Zeppelin, T.K., K.A. Call, D.J. Tollit, T.J. Orchard, and C.J. Gudmundson. 2003. Estimating the size of walleye pollock and Atka mackerel consumed by the western stock of Steller sea lions. Marine Science in the Northeast Pacific. Sponsored by Exxon Valdez Oil Spill Trustee Council, GLOBEC-Northeast Pacific Program, Steller Sea Lion Investigations, North Pacific Research Board, North Pacific Marine Research Institute and Pollock Conservation Cooperative. Anchorage, Alaska. January 13-17.
- Zielger, A.D., R.A. Sutherland, and T.W. Gaimbelluca. 2001. "Interstorm surface preparation and sediment detachment by vehicle traffic on unpaved mountain roads." *Earth Surface Processes and Landforms*. 26:235-250.

### **Additional Resources**

- American Association of Port Authorities (AAPA). Environmental Management Handbook. [http://www.aapa-ports.org/govrelations/env\\_mgmt\\_hb.htm](http://www.aapa-ports.org/govrelations/env_mgmt_hb.htm)
- Aquatic Invasive Species Research and Outreach: <http://www.nsgo.seagrant.org/research/nonindigenous/>.
- Baird, R.C. 1996. "Toward new paradigms in coastal resource management: Linkages and institutional effectiveness." Spagnolo, RJ; E. Ambrogio, F.J. Rielly, Jr., eds. *Estuaries*. 19(2A): 320-335.
- Beckmen, K.B., K.A. Burek, K.W. Pitcher, G.M. Ylitalo, and L.K. Duffy. *In press*. Mercury concentrations in the fur of Steller sea lions and northern fur seals from Alaska. *Mar. Poll. Bull.*
- Benaka, L., ed. 1999. Fish Habitat: essential fish habitat and rehabilitation. American Fisheries Society, Symposium 22, Bethesda, Maryland, 459 pp.

- Bettinger, P., J. Sessions, and K.N. Johnson. 1998. "Ensuring the compatibility of aquatic habitat and commodity production goals in eastern Oregon with a Tabu search procedure." *Forest Science*. 44(1):96-112.
- Bilby, R.E. and J.W. Ward. 1991. "Characteristics and function of large woody debris in streams draining old-growth, clear-cut, and second-growth forests in southwestern Washington." *Canadian Journal of Fisheries and Aquatic Sciences*. 48:2499-2508.
- Bisson, P.A., J.L. Nielsen, and J.W. Ward. 1988. "Summer production of coho salmon stocked in Mount St. Helens streams 3-6 years after the 1980 eruption." *Transactions of the American Fisheries Society*. 117(4):322-335.
- Blanton, S.L., R.M. Thom, A.B. Borde, H. Diefenderfer, and J.A. Southard. 2002. Evaluation of methods to increase light under ferry terminals. Final Research Report prepared for Washington State Department of Transportation by Battelle, Pacific Northwest Division of Battelle Memorial Institute. PNNL-13714. 26 pp.
- Blanton, S.L., R.M. Thom, J.A. Southard. 2001. Documentation of ferry terminal shading, substrate composition, and algal and eelgrass coverage. Letter Report prepared for University of Washington, School of Aquatic and Fishery Sciences, Seattle, Washington by Battelle Marine Sciences Laboratory, Sequim, Washington. 17 pp.
- Bue, B.G., S. Sharr, and J.E. Seeb. 1998. "Evidence of damage to pink salmon populations inhabiting Prince William Sound, Alaska, two generations after the *Exxon Valdez* oil spill." *Transactions of the American Fisheries Society*. 127: 35-43.
- Cardwell, R.D., M.I. Carr, and E.W. Sanborn. 1980. Water quality and flushing of five Puget Sound marinas. Technical Report No. 56. Washington Department of Fisheries Research and Development. Olympia, WA. 77 pp.
- Castro, J. 2003. Geomorphologic impacts of culvert replacement and removal: avoiding channel incision. USFWS-Oregon Fish and Wildlife Office, Portland, OR.
- Cederholm, C.J. and L.M. Reid. 1987. Impact of Forest Management on Coho Salmon (*Oncorhynchus kisutch*) Populations of the Clearwater Rivers, Washington: A Project Summary. In *Streamside Management: Forestry and Fishery Interactions*. Salo, E.O. and T.W. Cundy, eds. College of Forest Resources, University of Washington, Seattle, Washington. University of Washington, Institute of Forest Resources. Contribution No. 57.
- Cederholm, C.J. and N.P. Peterson. 1985. "The retention of coho salmon (*Oncorhynchus kisutch*) carcasses by organic debris in small streams." *Canadian Journal of Fisheries and Aquatic Sciences*. 42:1222-1225.
- Cederholm, C.J., D.B. Houston, D.L. Cole, and W.J. Scarlett. 1989. "Fate of coho salmon (*Oncorhynchus kisutch*) carcasses in spawning streams." *Canadian Journal of Fisheries and Aquatic Sciences*. 46, 1347-1355.
- Chambers, J.R. 1992. Coastal degradation and fish population losses. pp 45-51. In Stroud, R.H. ed. *Stemming the tide of coastal fish habitat loss*. Proceedings of a symposium on conservation of coastal fish habitat. National Coalition for Marine Conservation, Inc., Savannah GA.
- Coffin, B.A. and R.D. Harr. 1992. Effects of forest cover on volume of water delivery to soil during rain-on-snow. Washington Department of Natural Resources Timber/Fish/Wildlife Program. TFW-SH1-92-001.
- Collins, M.A. 1995. Dredging-induced near-field re-suspended sediment concentration and source strengths. Miscellaneous Paper D-95-2, U.S. Army Engineer Waterways Experiment Station, Vicksburg, MS. NTIS No. AD A299 151.

- Culp, J.M. and R.W. Davies. 1983. "An assessment of the effects of streambank clear cutting on macroinvertebrate communities in a managed watershed." *Canadian Technical Report of Fisheries and Aquatic Sciences*. No. 1208. 116 pp.
- Drinkwater, K.F. and K.T. Frank. 1994. "Effects of river regulation and diversion on marine fish and invertebrates." *Aquatic Conservation.: Mar. Freshwat. Ecosyst.* 4(2):135-151.
- Ellis, D.V. 1982. *Marine Tailings Disposal*. Ann Arbor Science Publishers. Ann Arbor, Michigan. 368 pp.
- EPA (U.S. Environmental Protection Agency). 2000. Scoping Document for the Proposed Pogo Mine Site. U.S. EPA, Region 10, 1200 Sixth Ave., Seattle, WA 98101. 11 August. (<http://www.pogomineeis.com/documents/ScopingDraft.pdf>).
- EPA, Office of Water (<http://www.epa.gov/water/>) and Office of Pesticide Programs (<http://www.epa.gov/pesticides/>).
- EPA, Office of Wetlands, Oceans and Watersheds, Ocean and Coastal Protection Division, Marine Debris Abatement (<http://www.epa.gov/owow/oceans/debris/>)
- EPA. 1975. Development document for effluent limitations guidelines and new source performance standards for the fish meal, salmon, bottom fish, clam, oyster, sardine, scallop, herring, and abalone segment of the canned and preserved fish and seafood processing industry point source category. Effluent Guidelines Division, Office of Water and Hazardous Material, Washington, D.C. EPA 440/1-75/041a. 485 pp.
- EPA. 1992. Turning the Tide on Trash: Marine Debris Curriculum. A Learning Guide on Marine Debris. EPA842-B-92-003 (<http://www.epa.gov/owow/OCPD/Marine/contents.html>)
- EPA. 1993. Guidance specifying management measures for sources of nonpoint pollution in coastal waters. 640-B-92-002. U.S. Environmental Protection Agency, Office of Water, Washington, D.C., January.
- EPA. 1997. Office of Water. Guidance specifying management measures for sources of nonpoint pollution in coastal waters. Chapter 5: Management measure for marinas and recreational boating. (<http://www.p2pays.org/ref/04/03686/index-5.html>).
- Ewing, R.D. 1999. Diminishing returns: salmon decline and pesticides. Oregon Pesticide Education Network, Eugene, Oregon. 52 pp.
- Faris, T.L. and K.D. Vaughan. 1985. Log transfer and storage facilities in Southeast Alaska: a review. USDA Forest Service General Technical Report PNW-174. Portland, OR.
- Farnworth, E.G., M.C. Nichols, C.N. Vann, L.G. Wolfson, R.W. Bosserman, P.R. Hendrix, F.B. Golley, and J.L. Cooley. 1979. Impacts of sediment and nutrients on biota in surface waters of the United States. U.S. Environmental Protection Agency. Athens, GA (USA), Oct 1979. Ecol. Res. Series. 331 pp.
- Flanders, L.S. and J. Cariello. 2000. Tongass Road Condition Survey Report. Alaska Department of Fish and Game, Southeast Regional Office of the Habitat and Restoration Division, Alaska Department of Fish and Game, 802 3rd Street, Douglas, AK., 99824. Technical Report 00-7. 48 pp.
- Fresh, K. 1997. The role of competition and predation in the decline of Pacific salmon and steelhead. In Stouter, D., P. Bisson, and R. Naiman, eds. *Pacific Salmon and Their Ecosystems: Status and Future Options*. Chapman and Hall, New York.

- Furniss, M.J., T.D. Roelofs, and C.S. Yee. 1991. Road construction and maintenance. *In* Meehan W.R., ed. Influences of forest and rangeland management on salmonid fishes and their habitats. Special Publication 19. Bethesda, MD: American Fisheries Society: 297-323.
- Gayaldo, P., S. Wyllie-Echeverria, and K. Ewing. 2001. Transplantation and alteration of submarine environment for restoration of *Zostera marina* (eelgrass); a case study at Curtis Wharf (Port of Anacortes), Washington. Presented at Puget Sound Research 2001, February 12-14, 2001, Bellevue, WA.
- Gibbons, D.R., W.R. Meehan, K.V. Koski, and T.R. Merrell, Jr. 1987. History of fisheries and forestry interactions in southeastern Alaska. *In* Salo, E.O. and T.W. Cundy, eds. Streamside Management: Forestry and Fishery Interactions. Contribution No. 57, University of Washington, College of Forest Resources, Seattle. pp. 297-329.
- Good, J.W. 1987. Mitigating estuarine development impacts in the Pacific Northwest: from concept to practice. Northwest Environmental Journal. Volume 3, Number 1.
- Gregory, R.S. 1993. "Effect of turbidity on the predator avoidance behavior of juvenile chinook salmon (*Oncorhynchus tshawytscha*)."  
*Canadian J. Fish. Aquatic Sciences*. 50:241-246.
- Gucinski, H., M.J. Furniss, R.R. Ziemer, and M.H. Brookes. 2000. Forest Roads: A Synthesis of Scientific Information, USDA Forest Service. June 2000. 117 pp.
- Harvey, B.C. and T.E. Lisle. 1998. "Effects of suction dredging on stream: A review and an evaluation strategy." *Fisheries*. 23:8-17.
- Hastings, K., P. Hesp, and G. Kendrick. 1995. "Seagrass loss associated with boat moorings at Rottnest Island, Western Australia." *Ocean and Coastal Management*. 26:225-246.
- Heifetz, J., M.L. Murphy, and K.V. Koski. 1986. "Effects of logging on winter habitat of juvenile salmonids in Alaskan streams." *North American Journal of Fisheries Management*. 6:52-58.
- Heintz, R.A., J.W. Short, and S.D. Rice. 1999. "Sensitivity of fish embryos to weathered crude oil: part II. incubating downstream from weathered *Exxon Valdez* crude oil caused increased mortality of pink salmon (*Oncorhynchus gorbuscha*) Embryos." *Environmental Toxicol. Chem*. 18:494-503.
- Herke, W.H. and B.D. Rogers. 1993. Maintenance of the estuarine environment. pp. 263-286. *In* Kohler, C.C. and W.A. Hubert, eds. Inland fisheries management in North America. American Fisheries Society, Bethesda, MD.
- Hicks, B.J., J.D. Hall, P.A. Bisson, and J.R. Sedell. 1991. Responses of salmonids to habitat changes. *In* Meehan, W.R., ed. Influences of Forest and Rangeland Management on Salmonid Fishes and their Habitats. American Fisheries Society Special Publication 19, 483-518.
- Hilderbrand, R.H., A.D. Lemly, C.A. Dolloff, and K.L. Harpster. 1998. "Design considerations for large woody debris placement in stream enhancement projects." *North American Journal of Fisheries Management*. 18 (1):161B167.
- Independent Multidisciplinary Science Team. 2002. Recovery of Wild Salmonids in Western Oregon Lowlands. Technical Report 2002-1 to the Oregon Plan for Salmon and Watersheds, Governor's Natural Resources Office, Salem, OR.
- Johnson, S.W., J. Heifetz, and K.V. Koski. 1986. "Effects of logging on the abundance and seasonal distribution of juvenile steelhead in some southeastern Alaska USA streams." *North American Journal of Fisheries Management*. 6, 532-537.
- King County Department of Natural Resources and Parametrix, Inc. 2002. Bioaccumulation and King County secondary treated effluent: data review, method evaluation, and potential on Puget Sound

- aquatic life. Unpublished paper in support of the draft Habitat Conservation Plan-in-progress. 38 pp + appendix.
- Klimley, A.P. and S.C. Beavers. 1998. "Playback of ATOC-type signal to bony fishes to evaluate phonotaxis." *Journal of Acoustic Society of America*. 104:2506-2510.
- Kline, E.R. 1994. Potential biological consequences of submarine mine-tailings disposal: a literature synthesis. U.S. Dept. of the Interior, Bureau of Mines, Juneau, AK, OFR 36-94.
- Koski, K.V. 1975. The survival and fitness of two stocks of chum salmon (*Oncorhynchus keta*) from egg deposition to emergence in a controlled-stream environment at Big Beef Creek. Ph.D. Dissertation, University of Washington, Seattle, WA.
- Koski, K.V. 1992. Restoring stream habitats affected by logging activities. Pages 343-404 in G. W. Thayer (editor) Restoring the nation's marine environment. Publication UM-SG-TS-92-06. Maryland Sea Grant College, College Park, MD.
- Koski, K.V., J. Heifetz, S. Johnson, M. Murphy, and J. Thedinga. 1984. Evaluation of buffer strips for protection of salmonid rearing habitat and implications for enhancement. In Hassler, T.J., ed. Proceedings: Pacific Northwest Stream Habitat Management Workshop, American Fisheries Society, Humboldt State University, Arcata, CA. pp.138-155.
- LaSalle, M.W., D.G. Clarke, J. Homziak, J.D. Lunz, and T.J. Fredette. 1991. A framework for assessing the need for seasonal restrictions on dredging and disposal operations. Technical Report D-91-1, U.S. Army Engineer Waterways Experiment Station, Vicksburg, MS. NTIS No. AD A240 567.
- Lee, D.C., J.R. Sedell, B.E. Rieman, R.F. Thurow, J.E. Williams, D. Burns, J. Clayton, L. Decker, R. Gresswell, R. House, P. Howell, K.N. Lee, K. MacDonald, J. McIntyre, S. McGivney, T. Noel, J.E. O'Connor, C.S. Overtone, D. Perkinson, K. Tu, and P. VanEimeren. 1997. Broad-scale assessment of aquatic species and habitats. Vol III, Chapter 4. In Quigley, T.M. and S. J. Arbelbide, eds. An assessment of ecosystem components in the interior Columbia basin and portions of the Klamath and Great Basins. Gen. Tech. Rep. PNW-405. Portland, OR: U.S. Department of Agriculture, Forest Service, Pacific Northwest Research Station.
- Leonard, J.N. 1994. Ocean outfalls for wastewater discharges – meeting CWA 403C requirements. Marine Technology Soc. '94, Conference Proceedings. Challenges and Opportunities in the Marine Environment, Washington, DC, 7-9 Sept. pp. 115-120.
- Love, M.S., J.E. Caselle, and L. Snook. 1999. "Fish assemblages on mussel mounds surrounding seven oil platforms in the Santa Barbara Channel and Santa Maria Basin." *Bulletin Marine Science*. 55:1062-1085.
- McFadden, J.T. 1969. Dynamics and regulation of salmonid populations in streams. In: Northcote, T.E., ed. Symposium on salmonid populations in streams. Vancouver: University of British Columbia, Institute of Fisheries. pp. 313-329.
- McLusky, D.S., D.M. Bryant, and M. Elliot. 1992. "The impact of land-claim on macrobenthos, fish and shorebirds on the Forth Estuary, eastern Scotland." *Aquat. Conserv.: Mar. Freshwat. Ecosyst.* 2 (3):211-222.
- Meehan, W.R., W.A. Farr, D.M. Bishop, and J.H. Patric. 1969. Some effects of clearcutting on salmon habitat of two southeast Alaska streams. USDA Forest Service General Technical Report PNW-82. Institute of Northern Forests., Pacific Northwest Forest and Range Experiment Station.
- Murphy, M.L. and K.V. Koski. 1989. Input and depletion of woody debris in Alaska streams and implications for streamside management. *North American Journal of Fisheries Management* 9, 427-436.



- Murphy, M.L. and J.D. Hall. 1981. "Varied effects of clear-cut logging on predators and their habitat in small streams of the Cascade Mountains, Oregon." *Canadian Journal of Fisheries and Aquatic Sciences*. 38:137-145.
- Myren, R.T. and R.J. Ellis. 1984. Evapotranspiration in forest succession and long-term effects upon fishery resources: a consideration for management of our old-growth forests. Pages 183-186 IN: M.R. Meehan, T.R. Merrill, Jr. and T.A. Hanley, eds. Fish and wildlife relationships in old growth forests. American Institute of Fishery Research Biologists.
- NOAA (National Oceanic and Atmospheric Administration). 1991. National Status and Trends Program for marine environmental quality. Progress report on secondary summary of data on chemical contaminants in sediments from the National Status and Trends Program. Tech. Mem. NOS OMA 59. NOAA, NOS, Silver Spring, MD. 29 pp.
- NMFS. 2001. Guidelines for salmonid passage at stream crossings. Southwest Region, Santa Rosa, CA. 14 pp. (<http://swr.nmfs.noaa.gov/hcd/NMFSSCG.PDF>).
- NRC (National Research Council). 2003. Cumulative Environmental Effects of Oil and Gas Activities on Alaska's North Slope. Committee on the Cumulative Environmental Effects of Oil and Gas Activities on Alaska's North Slope. The National Academies Press, Washington, D.C. Prepublication Copy. 452 pp. ([www.nap.edu](http://www.nap.edu)).
- Paul, J.F., K.J. Scott, A.F. Holland, S.B. Weisberg, J.K. Summers, and A. Robertson. 1992. "The estuarine component of the U.S. EPA's Environmental Monitoring and Assessment Program." Papers from the First International Ocean Pollution Symposium, 28th April 1991--2nd May 1991. University of Puerto Rico, Puerto Rico, (Part Two). *Chem. Ecol.* 7 (1-4): 93-116.
- Peddicord, R.K. and J. B. Herbich, ed. 1979. Impacts of open-water dredged material discharge. Proceedings of the eleventh dredging seminar. Published by TAMU, College Station, TX (USA). Oct 1979. pp. 24-40. Rep. Texas A&M University Sea Grant Program.
- Reiser and Bjornn. 1979. Influence of forest and rangeland management on anadromous fish habitat in the Western United States and Canada. Meehan, W.R. ed. USDA Forest Service General Technical Report PNW-96. U.S. Department of Agriculture, Forest Service, Pacific Northwest Forest and Range Experiment Station. Portland, OR.
- Rice, S.D., D.A. Moles, J.F. Karinen, S. Korn, M.G. Carls, C.C. Brodersen, J.A. Gharrett, and M.M. Babcock. 1984. Effects of petroleum hydrocarbons on Alaskan aquatic organisms: A comprehensive review of all oil-effects research on Alaskan fish and invertebrates conducted by the Auke Bay Laboratory, 1970 - 81. U.S. Department of Commerce, NOAA Technical Memorandum NMFS F/NWC-67. 128 pp.
- Rozengurt, M.A., I. Haydock, and B.P. Anderson. 1994. Running on entropy: The effect of water diversion on the coastal zone. 37th Conference of the International Association for Great Lakes Research and Estuarine Research Federation: Program and Abstracts., Buffalo, NY (USA). 166 pp.
- RPWAST (Rich Passage Wave Action Study Team). 2001. Rich Passage Wave Action Study Final Report, WAC 197-11-970. Prepared for Washington State Ferries. August 27.
- Schaumburg, F.D. 1970. The influence of log handling on water quality, annual report 1969-1970, Department of Civil Engineering, Oregon State University. Corvallis, Oregon.
- Sea Grant. 2002. National Coastal Ecosystem Restoration Manual. Sea Grant Communications, Oregon State University, Corvallis, OR. Pacific Northwest: <http://seagrants.oregonstate.edu/hot/exotics.html>.

- Sedell, J.R., F.N. Leone, and W.S. Duvall. 1991. Water transportation and storage of logs. *In* Meehan, W.R., ed. Influences of Forest and Rangeland Management on Salmonid Fishes and their Habitats. American Fisheries Society Special Publication 19:325-368.
- Shafer, D.J. 2002. Recommendations to minimize potential impacts to seagrasses from single-family residential dock structures in the Pacific Northwest. U.S. Army Corps of Engineers, Seattle, WA. 28 pp.
- Simenstad, C.A. 1994. pp. 11-19. *In* Wyllie-Echeverria, S., A.M. Olson, and M.J. Hershman, eds. Seagrass science and policy in the Pacific Northwest: proceedings of a seminar series (SMA 94-1) EPA 910/R-94-004. 63 pp.
- Simenstad, C.A., B.J. Nightingale, R.M. Thom, and D.K. Shreffler. 1999. Impacts of Ferry Terminal on Juvenile Salmon Migrating along Puget Sound Shorelines Phase I: Synthesis of State of Knowledge.
- Sogard, S.M. and K.W. Able. 1991. "A comparison of eelgrass, sea lettuce macroalgae and marsh creeks as habitats for epibenthic fishes and decapods." *Estuarine, Coastal and Shelf Science*. 33:501-519.
- Stone, R.P. and S.W. Johnson. 1998. "Prolonged exposure to mine tailings and survival and reproductive success of ovigerous Tanner crabs (*Chionoecetes bairdi*)." *Bull. Environ. Contam. Toxicol.* 61: 548-556.
- Stull, J.K. and C.I. Haydock. 1989. Discharges and environmental responses: the Palos Verdes case. *In* Managing inflows in California's bays and estuaries. The Bay Institute, Sausalito, Calif. pp. 44-49.
- Swanston, D.N. 1974. The forest ecosystem of southeast Alaska. Soil mass movement. General Technical Report PNW-17. U.S. Department of Agriculture. Forest Service. Pacific Northwest Forest and Range Experiment Station. Portland, OR.
- TAPS (Trans-Alaska Pipeline System) Draft Environmental Impact Statement. 2002. Renewal of the Federal Grant for the Trans-Alaska Pipeline System (TAPS) Right-of-Way. U.S. Dept. of the Interior, Bureau of Land Management. 4.3: 50.
- Thayer, G.W., ed. 1992. Restoring the Nation's Marine Environment. Publication UM- SG-TS-92-06. Maryland Sea Grant College, College Park, MD, 716 pp.
- Thorsteinson, F.V. and L.K. Thorsteinson. 1982. Finfish Resources. Chapter 6. *In* Hameedi, M.J., ed. The St. George Basin Environment and Possible Consequences of Planned Offshore Oil and Gas Development. Proceedings of a Synthesis Meeting. U.S. Department of Commerce, NOAA, Office of Marine Pollution Assessment.
- Turek, J.G., T.E. Bigford, and J.S. Nichols. 1987. Influence of freshwater inflows on estuarine productivity. NOAA Tech. Memo. NMFS-F/NEC-46. 26 pp.
- U.S. Army Corps of Engineers, Alaska District. 1999. Final Environmental Impact Statement. Beaufort Sea Oil and Gas Development/Northstar Project. P.O. Box 898, Anchorage, Alaska. February 1999.
- U.S. Department of Agriculture. 2002. Southern Intertie Project Final Environmental Impact Statement. U.S. Dept. of Agriculture. Rural Utilities Service. Washington D.C.
- USDA (U.S. Department of Agriculture). 1989. The second RCA appraisal: Soil, water, and related resources on nonfederal land in the United States, analysis of conditions and trends. Washington, D.C. 280 pp.

- Vernberg F.J., W.B. Vernberg. 2001. The Coastal Zone. Past, Present, and Future. University of South Carolina.
- Washington State Department of Ecology, Water Quality Program, Sand & Gravel General Permit (<http://www.ecy.wa.gov/programs/wq/sand/escp.html>).
- Wicker, K.M., R.E. Emmer, D. Roberts, and J. van Beek. 1989. Pipelines, navigation channels, and facilities in sensitive coastal habitats: an analysis of Outer Continental Shelf impacts, Coastal Gulf of Mexico. Volume I: technical narrative. U.S. Dept. of the Interior, Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA. OCS Report MMS 89-0051.
- Williams, R.N., L.D. Calvin, C.C. Coutant, M.W. Erho Jr., J.A. Lichatowich, W.J. Liss, W.E. McConnaha, P.R. Mundy, J.A. Stanford, R.R. Whitney, D.L. Bottom, and C.A. Frissell. 1996. Return to the river: restoration of salmonid fishes in the Columbia River ecosystem. Development of an alternative conceptual foundation and review and synthesis of science underlying the Columbia River Basin fish and wildlife program of the northwest power planning council by the independent scientific group. Northwest Power Planning Council, Portland, Oregon. 584 pp.
- Williams, J.E., C.A. Wood, and M.P. Dombeck, eds. 1997. Watershed Restoration: Principles and Practices. American Fisheries Society, Bethesda, MD. 561 pp.
- Woltemade, C.J. 2000. "Ability of restored wetlands to reduce nitrogen and phosphorus concentrations in agricultural drainage water." *J. Soil and Water Cons.* 55: 303.
- Young, K.A. 2000. "Riparian zone management in the Pacific Northwest: who's cutting what?" *Environmental Management*. 26, 131-44.

### **Online Resources**

BMPs are often specific to certain geographical locations or pesticide application programs, with the aim of reducing or eliminating pesticide transport to surface waters. An example of a pesticide use reduction strategy for a large city (Seattle) is available at: <http://www.metrokc.gov/hazwaste/ipm/>.

and from the following:

EPA. 1984. Best Management Practices for Agricultural Nonpoint Source Control: IV. Pesticides. EPA Number: 841S84107.

Dredging BMPs: <http://www.spn.usace.army.mil/ltms2001/appi.pdf>

Various IPM strategies can be found at:

U.S. Department of Agriculture. Cooperative State Research, Education, and Extension (CSREES) Program (<http://www.ree.usda.gov/ipm/>).

Federal and State invasive species activities and programs: <http://www.invasivespecies.gov/new/whatsnew.shtml>

Logging effects studies on fish habitat (bibliography):  
<http://www.afsc.noaa.gov/abl/habitat/pdfs/logging.pdf>

The following Links provide detailed standards and guidance:

Best Management Practices: [http://www.fs.fed.us/r10/TLMP/F\\_PLAN/APPEND\\_C.PDF](http://www.fs.fed.us/r10/TLMP/F_PLAN/APPEND_C.PDF)

Stream Process Groups: [http://www.fs.fed.us/r10/TLMP/F\\_PLAN/APPEND\\_D.PDF](http://www.fs.fed.us/r10/TLMP/F_PLAN/APPEND_D.PDF)

Watershed Analysis: [http://www.fs.fed.us/r10/TLMP/F\\_PLAN/APPEND\\_J.PDF](http://www.fs.fed.us/r10/TLMP/F_PLAN/APPEND_J.PDF)

Riparian: [http://www.fs.fed.us/r10/ TLMP/F\\_PLAN/FPCHAP4.PDF](http://www.fs.fed.us/r10/TLMP/F_PLAN/FPCHAP4.PDF)

Transportation: [http://www.fs.fed.us/r10/TLMP/F\\_PLAN/FPCHAP4.PDF](http://www.fs.fed.us/r10/TLMP/F_PLAN/FPCHAP4.PDF)

Beach and Estuary Fringe: [http://www.fs.fed.us/r10/TLMP/F\\_PLAN/FPCHAP4.PDF](http://www.fs.fed.us/r10/TLMP/F_PLAN/FPCHAP4.PDF)

Fish: [http://www.fs.fed.us/r10/TLMP/F\\_PLAN/FPCHAP4.PDF](http://www.fs.fed.us/r10/TLMP/F_PLAN/FPCHAP4.PDF)

Wetlands: [http://www.fs.fed.us/r10/TLMP/F\\_PLAN/FPCHAP4.PDF](http://www.fs.fed.us/r10/TLMP/F_PLAN/FPCHAP4.PDF)

Soils and Water: [http://www.fs.fed.us/r10/TLMP/F\\_PLAN/FPCHAP4.PDF](http://www.fs.fed.us/r10/TLMP/F_PLAN/FPCHAP4.PDF)

## MASTER INDEX

### A

agriculture 2.1.1, 2.1.3, 4.8, 5.2, 5.3  
animal waste 2.1.1, 4.11  
artificial reef 4.11, 5.4, 5.5  
avoidance 4.1, 4.5.1

### B

ballast water discharge 4.3, 4.4  
bark 4.9  
beach nourishment 4.7, 4.8  
bioaccumulation 2.1.1, 3.2.2, 5.7  
biological oxygen demand see oxygen depletion  
boat mooring 4.3  
bridge 2.2, 2.1.2, 4.10  
bubble curtain 4.5.1

### C

cables 4.10  
channel 1.0, 2.1, 2.1.1, 2.2, 2.3, 3.1.1, 3.1.2, 3.2.1, 3.4, 4.1, 4.3, 4.6, 4.7, 4.10,  
channelization 2.1, 2.2  
Clean Water Act 2.1, 2.1.3, 3.2.2, 5.1, 5.3  
commercial utilization of habitat 4.11  
contaminant 4.2.1, 4.6, 4.10, 4.11, 5.1, 5.4, 5.7  
cultural eutrophication 2.1.1  
culvert 2.1.2, 2.3, 4.8

### D

dam operations 3.2.1, 3.3, 3.4, 5.5  
desalinization plant 5.3  
discharge plumes 5.3  
disposal of dredged material 4.2.1, 4.3  
diversion 2.1.1, 2.3, 3.4  
dredging 3.1, 3.1.1, 4.1, 4.2, 4.2.1, 4.3, 4.4, 4.6, 4.8, 4.11, 5.3, 5.4, 5.5, 5.6

### E

economic development 2.2  
eelgrass 1.0, 2.1, 4.1, 4.2.1, 4.3, 4.6, 4.7, 4.8, 4.11, 5.1  
Endangered Species Act 1.0  
entrainment 4.1, 5.3  
entrapment 3.3, 3.4, 4.1, 5.3  
erosion 2.1.1, 2.1.2, 2.3, 3.1, 4.2.1, 4.2.2, 4.3, 4.10, 5.5  
essential fish habitat (EFH) 1.0  
essential fish habitat assessment 1.0  
estuary 1.0, Chapter 4  
exotic species 2.3, 4.3, 4.4, 4.11, 5.5

## **F**

Federal Insecticide, Fungicide, and Rodenticide Act (FIFRA) 2.1.3  
fill material 4.2.2  
Fish and Wildlife Coordination Act 1.0  
fish passage 2.1.2, 2.3, 3.3, 4.7, 4.8, 5.5  
fish processing 5.2  
flood control 4.7

## **G**

gas, natural 3.2.2, 5.4  
gravel 3.1.2, 3.2.1, 5.6  
grazing 2.1.1  
groundfish 1.0, 4.2.1, 4.9, 4.11, 5.4  
groundwater 2.1.1, 2.1.3, 2.2, 3.1.1, 4.7, 5.4

## **H**

habitat enhancement 5.5  
habitat restoration 5.5  
heavy metals 4.7, 5.1

## **I**

impingement 5.3  
impoundment 1.0, 3.4, 4.8  
infiltration 2.1.1, 2.1.2, 2.2, 5.1  
inorganic debris 3.2.2  
invasive species see exotic species  
in-water log storage 4.9  
irrigation 2.1.1, 2.1.3, 3.4

## **J**

## **K**

kelp 1.0, 3.2.1, 4.1, 4.2.1, 4.9, 4.11, 5.1

## **L**

land subsidence 4.7  
large woody debris (LWD) see woody debris  
light penetration 2.1, 2.1.1, 2.3, 4.1, 4.2.1, 5.1, 5.2, 5.4  
livestock 2.1.1  
log transfer facilities 2.1.2, 4.9  
logging 2.1.2, 4.9

## **M**

macrophyte wrack 3.2.1  
Magnuson-Stevens Fishery Conservation and Management Act 1.0  
manure 2.1.1  
marine debris see inorganic debris  
mass wasting 2.1, 2.1.2, 2.2, 2.3  
migration 2.1.2, 2.3, 3.1.2, 3.3, 4.6, 4.7, 5.1, 5.3, 5.5

mineral mining 3.1.1  
mining 3.1, 3.1.1, 3.1.2

## **N**

National Environmental Policy Act (NEPA) 1.0  
net primary productivity see productivity  
nitrogen 2.1.1, 4.8, 5.1  
noise see sound  
nonpoint source 2.1, 2.1.1, 2.1.2, 2.1.3, 2.2, 2.3, 4.3, 4.7, 5.2  
National Pollutant Discharge Elimination System (NPDES) 3.2.2, 4.9, 5.2,  
nursery, plant 2.1.1  
nursery, fish 4.1, 4.8, 4.11  
nutrients 2.2, 3.1.2, 3.3, 4.2.1, 4.8, 5.1

## **O**

oil 2.2, 3.2.2, 4.9, 5.4,  
oil spills 5.4  
organic debris 3.2.1, 4.7, 4.11  
overwater structures 4.3, 4.6, 4.9  
oxygen depletion 2.1.1, 2.1.2, 2.2, 3.2.2, 3.4, 4.1, 4.2.1, 4.3, 4.8, 4.9, 4.11, 5.1, 5.2, 5.4  
oysters 4.11

## **P**

PCBs 5.7  
Persistent Organic Pollutants (POPs) 5.7  
pesticide 2.1.1, 2.1.3, 3.3, 4.1, 4.10, 5.1, 5.7  
petroleum 2.2, 4.10, 5.4 see gas, oil  
phosphorus 2.1.1, 2.2  
pier see overwater structure  
pile driving see pile installation  
pile installation 4.5, 4.5.1  
pile removal 4.5, 4.5.2  
pipeline 4.1, 4.10, 5.1, 5.4  
point source 5.1, 5.2  
polyaromatic hydrocarbons (PAH) 2.2, 4.1, 4.6, 5.4  
port expansion 4.1, 4.3  
power generating station 5.3  
predation 4.6, 5.2, 5.7  
primary productivity see productivity  
productivity 1.0, 2.1, 2.1.3, 4.1, 4.2.1, 4.2.2, 4.3, 4.6, 4.8, 4.9, 4.10, 4.11, 5.1, 5.2, 5.4  
pumping stations 4.8

## **Q**

## **R**

riparian 1.0, 2.1.1, 2.1.2, 2.1.3, 2.2, 2.3, 3.1, 3.1.1, 3.1.2, 3.3, 3.4, 4.1, 4.3, 4.7, 5.5  
road 1.0, 2.1, 2.1.1, 2.1.2, 2.2, 2.3, 3.1, 3.1.1, 4.8, 5.5  
runoff 2.1, 2.1.1, 2.1.2, 2.1.3, 2.2, 2.3, 3.2.2, 4.3, 4.5.2, 4.7, 5.1, 5.4

## **S**

salinity 2.1.1, 3.3, 5.1  
salt marsh see wetlands  
saltwater intrusion 4.7, 4.10  
sand 3.1.2  
sand and gravel mining 3.1.2  
seafood 5.2  
seagrass 4.1, 4.2.1, 5.1  
sediment 2.1.1, 2.2, 3.3, 4.1, 4.2, 4.5.2, 4.8  
sedimentation 1.0, 2.1.1, 2.1.2, 2.3, 3.1.2, 3.4, 4.1, 4.10, 5.4  
sewage 2.2, 3.2.2, 4.8, 4.10, 5.1  
sewage treatment plants 4.8  
shading 1.0, 4.3, 4.6, 4.7, 4.9, 4.11, 5.5  
shoreline protection 2.1, 4.2.1, 2.2, 4.3, 4.7  
silviculture 2.1.2  
soil compaction 2.1.1, 4.1, 4.7  
sound 4.1, 4.5.1, 5.3, 5.4,  
spawning 3.1.2, 4.1, 4.7, 5.5  
storm drains 2.2  
stream crossings 2.1.2  
submerged aquatic vegetation 2.1.1, 2.2, 4.2.1, 4.3, 4.6, 4.10, 4.11, 5.1, 5.5  
substrate 3.1, 4.2.1, 4.2.2, 4.5.1, 4.6, 4.11, 5.3, 5.4, 5.5, 5.6  
suburban development 2.2  
suspended sediment concentration 4.1

## **T**

tailings 3.1  
tailings disposal 3.1  
temperature, water 2.1.1, 2.1.2, 2.2, 2.3, 3.1, 3.1.2, 3.3, 4.3, 4.7, 5.1, 5.3  
tidegates 4.8  
timber harvest 2.1.2, 4.9  
toxic metals 3.1, 4.1, 4.7, 5.7  
transportation 4.3  
turbidity 1.0, 2.1.1, 3.1, 3.1.1, 3.1.2, 4.1, 4.2.1, 4.3, 4.5.2, 4.10, 5.1, 5.2, 5.3, 5.5, 5.6

## **U**

urban development 2.2, 4.8, 4.10  
utility line 4.10

## **V**

vessel operations 4.3, 5.2, 5.4

## **W**

wastewater plants 5.1  
water control structures 4.8  
water intake structures 5.3  
water quality 2.1.1, 2.1.3, 2.3, 3.1  
wave 4.3  
wave energy 4.7



wetlands 2.1.1, 3.2.1, 4.1, 4.3, 4.7, 4.10, 4.11, 5.1, 5.3, 5.5  
withdrawal 5.3  
woody debris 3.2.1, 3.3, 4.7  
wrack see macrophyte wrack

## **X**

## **Y**

## **Z**

Zone of Deposit 5.2